

# The Biology and Pathogenesis of Coronaviruses

H. WEGE\*, ST. SIDDELL\*, AND V. TER MEULEN\*

1	Introduction . . . . .	165
2	Biology . . . . .	166
2.1	Members of the Coronavirus Group and Their Relationships . . . . .	166
2.1.1	Antigenic Relationships . . . . .	166
2.1.2	Nucleic Acid Homologies . . . . .	168
2.2	Host Range and Organ Tropism . . . . .	168
3	Coronaviruses and Disease Spectrum . . . . .	171
3.1	Murine Coronaviruses . . . . .	171
3.1.1	Murine Hepatitis Virus . . . . .	171
3.1.1.1	Hepatitis . . . . .	171
3.1.1.2	Encephalomyelitis . . . . .	173
3.1.1.3	Enteritis . . . . .	174
3.2	Human Coronaviruses . . . . .	175
3.3	Avian Coronaviruses . . . . .	176
3.3.1	Infectious Bronchitis Virus . . . . .	176
3.3.2	Turkey Coronavirus . . . . .	177
3.4	Feline Coronaviruses . . . . .	178
3.4.1	Feline Infectious Peritonitis Virus . . . . .	178
3.5	Other Coronaviruses . . . . .	179
3.5.1	Bovine Coronavirus . . . . .	179
3.5.2	Canine Coronavirus . . . . .	180
3.5.3	Hemagglutinating Encephalomyelitis Virus . . . . .	181
3.5.4	Transmissible Gastroenteritis Virus . . . . .	181
3.5.5	Rat Coronavirus . . . . .	182
4	Pathogenetic Aspects . . . . .	183
4.1	The Role of Resistance in the Development of Disease . . . . .	183
4.1.1	Acute Infections . . . . .	183
4.1.1.1	Murine Hepatitis Virus Type 2 . . . . .	183
4.1.1.2	Murine Hepatitis Virus Type 3 . . . . .	184
4.1.1.3	Murine Hepatitis Virus JHM . . . . .	185
4.1.2	Chronic Infections . . . . .	186
4.2	Pathogenicity Associated with Viral Gene Sequences . . . . .	187
5	Conclusions . . . . .	188
	References . . . . .	189

## 1 Introduction

The coronaviruses were first recognized and morphologically defined as a group by *Tyrrell* and co-workers (1968, 1975, 1978). Biochemical studies have recently provided

\* Institute of Virology and Immunobiology, Versbacher Straße 7, D-8700 Wuerzburg, Federal Republic of Germany

additional information which allows better characterization of these agents. Presently, coronaviruses are defined as being particles which are pleomorphic to rounded with a diameter of 60–220 nm, surrounded by a fringe or layer of typical club-shaped spikes. The virion is composed of about four to six proteins and possesses a lipid bilayer. The genome consists of a single-stranded polyadenylated RNA which is infectious and of positive polarity. During maturation these viruses are released by internal budding into vesicles derived from the endoplasmatic reticulum. These viruses are widespread in nature and are associated with a great variety of diseases with an acute, subacute, or subclinical disease process.

Several reviews have been published describing aspects of the physicochemical and biological properties and the clinical significance of coronaviruses (*McIntosh 1974; Kapikian 1975; Pensaert and Callebaut 1978, Robb and Bond 1979*). During the past years new data on the biology of these viruses and on the pathogenesis of diseases, in particular murine-induced coronavirus diseases, have also become available. These recent findings are the basis for this review.

## 2 Biology

### 2.1 Members of the Coronavirus Group and Their Relationships

Table 1 lists the coronaviruses described to date, their natural hosts, and the predominant disease type as caused by these viruses.

#### 2.1.1 Antigenic Relationships

Our knowledge of the antigenic relationships between the different coronaviruses is incomplete. The relationships shown in Table 2 are based on results obtained by enzyme-linked immunoassay (*Macnaughton 1981; Kraaijeveld et al. 1980a, b*), immunofluorescent and immunoelectron microscopic studies (*Pedersen et al. 1978, Pensaert et al. 1981*), other serological methods (*Reynolds et al. 1980; Gerna et al. 1981*), and the data summarized by *Robb and Bond (1979)*. As shown, the avian and the nonavian coronaviruses each appear to fall into two distinct and unrelated groups. In the case of infectious bronchitis virus (IBV) at least eight different serotypes are at present known (*Hopkins 1974*) and these again fall into two groups by cluster analyses based on neutralization assays (*Darbyshire et al. 1979*). Also, comparison of the protein patterns of IBV isolates suggested that two groups exist which differ in the electrophoretic migration of the virion glycoproteins (*Nagy and Lomniczi 1979; Collins and Alexander 1980*).

The location of antigenic sites on coronavirion structural proteins has been investigated. Coronaviruses basically contain three major antigens, as has been shown by immunodiffusion experiments (*Hajer and Storz 1978; Yaseen and Johnson-Lussenburg 1981*) and by the analysis of monospecific antisera prepared against purified coronavirus structural proteins (*Schmidt and Kenny 1981*). In human, porcine, and murine systems the antigenic sites responsible for the induction of neutralizing antibodies are associated with the surface glycoproteins (peplomers). Immunological studies with subcomponents prepared from purified virions of TGEV (*Garwes et al. 1978*), HCV 229E and MHV-3 (*Macnaughton et al. 1981; Hasony and Macnaughton 1981*), and HCV-OC43 and 229E (*Schmidt*

Table 1. Coronaviruses – designations, natural host and predominant clinical disease

Virus <sup>a</sup>	Designation <sup>b</sup>	Natural host	Predominant clinical disease	First description
Avian infectious bronchitis virus	IBV (AIBV)	Chicken	Respiratory disease	<i>Schalk and Hawn 1931</i>
Bovine coronavirus (neonatal calf diarrhoea coronavirus, enteropathogenic bovine coronavirus, bovine enteric coronavirus)	BCV (NCDCV, EBC, BEC)	Calf	Diarrhoea	<i>Mebus et al. 1973a</i>
Canine coronavirus	CCV	Dog	Diarrhoea	<i>Binn et al. 1975</i>
Feline infectious peritonitis virus	FIPV	Cat	Peritonitis, granulomatous inflammations in multiple organs	<i>Holzworth 1963</i>
Human coronavirus	HCV	Man	Common cold	<i>Tyrrill and Bynoe 1965</i>
Murine hepatitis virus	MHV	Mouse	Encephalomyelitis, hepatitis, diarrhoea	<i>Cheever et al. 1949</i>
Porcine transmissible gastroenteritis virus	TGEV	Pig	Diarrhoea	<i>Doyle and Hutchings 1946</i>
Porcine hemagglutinating encephalomyelitis virus (vomiting and wasting disease virus)	HEV	Pig	Vomiting and wasting, encephalomyelitis	<i>Roe and Alexander 1958</i>
Parker's rat coronavirus <sup>c</sup>	RCV	Rat	Pneumonitis, rhinitis	<i>Parker et al. 1970</i>
Rat sialodacryoadenitisvirus <sup>c</sup>	SDAV	Rat	Adenitis	<i>Jonas et al. 1969</i>
Turkey coronavirus (turkey bluecomb disease coronavirus, turkey coronaviral enteritis virus, coronavirus enteritis of turkeys)	TCV (TBDCV, TCEV, CET)	Turkey	Enteritis	<i>Adams and Hojstad 1971</i>
<i>Questionable or unclassified members</i>				
Foal enteritis coronavirus	FECV	Horse	Diarrhoea	<i>Bass and Sharpee 1975</i>
Human enteric coronavirus <sup>d</sup>	HECV	Man	Diarrhoea	<i>Caul et al. 1975</i>
Isolates SD and SK	SD, SK	Mouse, man	Demyelinating encephalomyelitis in mice	<i>Burks et al. 1980</i>
Parrot coronavirus	-	Parrot	Diarrhoea	<i>Hirai et al. 1979</i>
Porcine CV-777 and other isolates	CV-777	Pig	Diarrhoea	<i>Pensaert and Debouck 1978</i>
Runde tick coronavirus <sup>e</sup>	RTCV	Seabird, tick	No data on disease in natural host	<i>Horváth and Mocsári 1981</i> <i>Travnik et al. 1977</i>

<sup>a</sup> In brackets, synonyms used in literature, <sup>b</sup> Abbreviations used in literature, <sup>c</sup> Both viruses might be serotypes of the rat coronaviruses, <sup>d</sup> Probably serotype(s) of human (respiratory) coronaviruses, <sup>e</sup> Probably a bunyavirus

Table 2. Antigenic cross-reactions between coronaviruses

<i>Mammalian</i>	
Group 1	Group 2
HCV-229 E and other isolates	HCV-OC43 and other isolates
TGEV one serotype	MHV many serotypes, also related to RCV and SDAV
CCV one serotype	BCV one serotype
FIPV one serotype	HEV one serotype
<i>Avian</i>	
Group 3	Group 4
IBV at least 8 serotypes	TCV one serotype
No cross-reactions with other strains	No cross-reactions with other strains
Unclassified isolates: Several isolates of HCV (and HECV), porcine coronavirus CV-777 and others, FECV, RTV	

and *Kenny* 1981) support this conclusion. A similar conclusion was reached by immunoelectron microscopy of bovine coronaviruses (*Storz and Rott* 1981). The surface glycoproteins are also involved in complement fixation and hemagglutinin inhibition.

### 2.1.2 Nucleic Acid Homologies

Some preliminary data on the nucleic acid sequence homology between a few coronaviruses is available. Hybridization with MHV-specific cDNA, representative of the entire genome, shows that a close relationship exists between the murine strains MHV-A59, MHV-3 and JHM. Using the same probe no homology between the murine viruses and the human coronavirus 229E could be detected (*Weiss and Leibowitz* 1981).

Using the technique of T<sub>1</sub> oligonucleotide fingerprinting *Lai and Stohlman* (1981a), *Weiss and Leibowitz* (1981), and *Wege et al.* (1981a) have shown variation in the genome RNA of murine hepatitis viruses of different neurovirulence (Sect. 4.2). This variation seems to be independent of the serological relationships of these strains. In the avian coronavirus group such an analysis also revealed considerable variation within serotypes (*Clewley et al.* 1981). Studies such as these might be useful in characterizing the origin, evolution and spread of both new isolates and live vaccine strains.

## 2.2 Host Range and Organ Tropism

Most coronaviruses cause clinical diseases only in the species from which they were isolated and replicate predominantly in cell lines derived from that host. However, transmission to other species can be achieved either experimentally or for some virus strains by a natural route of infection (Table 3). The natural infection of dogs by the porcine strain transmissible gastroenteritis virus (TGEV) and a single case of diarrhea transmitted from cattle to man may indicate a possibly wider host range for enteric infections. The experimental intracerebral inoculation of several coronaviruses into suckling rats, mice,

Table 3. Host range of coronaviruses

Virus strain	Natural host	Transmissible to	Route of inoculation	Effect on experimental host	References
HCV-OC43	Man	Suckling mice, suckling hamsters	Intracerebral	Encephalitis	<i>McIntosh et al. 1967</i>
TGEV	Pig	Dogs	Oral	Inapparent intestinal infection	<i>McIntosh et al. 1969</i> <i>Larson et al. 1979</i>
HEV-67N	Pig	Suckling mice	Intracerebral	Encephalitis	<i>Kaye et al. 1977</i>
BCV LY-138	Cattle	Man <sup>a</sup>	Oral (natural infection)	Diarrhea	<i>Storz and Rotz 1981</i>
BCV Nebraska	Cattle	Suckling mice <sup>b</sup>	Intracerebral	Encephalitis	<i>Kaye et al. 1975a</i>
BCV Kakegawa	Cattle	Suckling mice, rats, and hamsters	Intracerebral and intracutaneous	Encephalitis	<i>Akashi et al. 1981</i>
CCV	Dog	Piglets	Oral	Inapparent intestinal infection	<i>Woods et al. 1981</i>
FIPV	Cat and other feline species	Newborn mice, rats, and hamsters, piglets	Intracerebral	Inapparent CNS infection, inapparent intestinal infection	<i>Osterhaus et al. 1978a, b.</i> <i>Woods et al. 1981</i>
IBV Massachusetts and Beaudette	Chick	Suckling mice	Intracerebral	Encephalitis	<i>McIntosh et al. 1969</i> <i>Estola 1967</i>
MHV-JHM	Mouse	Monkeys	Intracerebral	Encephalomyelitis	<i>Kersting and Pette 1956</i>
		Rats	Intracerebral	Encephalomyelitis	<i>Cheever et al. 1949</i>
		Hamsters	Intracerebral and intranasal	Encephalomyelitis	<i>Cheever et al. 1949</i> <i>Bailey et al. 1949</i>
MHV-S	Mouse	Suckling rats	Intranasal	Asymptomatic infection	<i>Taguchi et al. 1979a</i>
MHV-A59	Mouse	Suckling rats	Intracerebral	Hydrocephalus and encephalitis	<i>Hirano et al. 1980</i> <i>Takahashi et al. 1980</i>
MHV-2 and 3	Mouse	Adult rats	Intracerebral	Clinically inapparent hepatitis, encephalitis	<i>Wege et al. 1981a</i> <i>(Wege et al. unpublished)</i>
SDAV	Rat	Suckling mice	Intracerebral	Encephalitis	<i>Jonas et al. 1969</i> <i>Bhatt et al. 1972</i>
TGEV	Pig	Dogs, foxes, cats	Oral	Virus shedding	<i>Haeterman 1962</i> <i>Reynolds and Garwes 1979</i>

<sup>a</sup> Only one accidental case report; <sup>b</sup> Other strains not transmissible to mice (*Dea et al. 1980a*)

Table 4. Target organs involved in coronavirus infections

Host species Virus	Avian		Bovine		Canine		Feline		Human		Murine		Porcine		Rat	
	IBV	TCV	BCV	CCV	FIPV	HCV	(HECV)	MHV	TGEV and HEV others	RCV	SDAV					
<i>Target organs</i>																
Central nervous system												++*				
Blood vessels					+							+				
Ependym																
Gonad	+															
Intestine		++	++	++												
Kidney	+				+		(++)									
Liver					++											
Lymphoid organs	+				++											
Pancreas																
Parotid gland																
Peritoneum					++											
Respiratory tract	++	+	+	+	+	++								+	++	+

*Symbols:* ++ main target for infection; + organs less frequently involved; \* involvement in persistent/chronic disease

or hamsters often induces an infection (Table 3). The brain of suckling mice is highly susceptible for viruses of avian, human, and mammalian origin. However, infection under these experimental conditions is not representative for the clinical disease in the natural host.

A survey of the organs involved in coronavirus infections is summarized in Table 4. Some coronaviruses reveal relatively restricted organ tropism leading to diseases of the respiratory system (HCV, IBV, RCV) and gastrointestinal tract only (BCV, CCV, TGEV, TCV). In other coronavirus infections, for example with feline and murine coronaviruses, several organs are involved. The murine coronaviruses represent a group containing many strains with different organ tropism. In addition, feline, murine, and avian coronavirus strains have a strong tendency to establish persistent and chronic diseases.

### 3 Coronaviruses and Disease Spectrum

#### 3.1 Murine Coronaviruses

##### 3.1.1 Murine Hepatitis Virus

The first murine coronavirus described was MHV-JHM, which was isolated from a spontaneously paralyzed mouse (*Cheever et al.* 1949). Subsequently, other strains were isolated from different disease conditions and different organs of mice (Table 5). Murine coronavirus infections are often subclinical or inapparent, but clinical disease can be activated by coinfection with leukemia viruses or protozoal agents. These viruses can be transmitted by feces or urine to susceptible strains (Table 5). Vertical transmission by intrauterine infection can also occur with MHV-JHM (*Katami et al.* 1978) and the respiratory route is important in natural transmission (*Carthew and Sparrow* 1981; *Taguchi et al.* 1979c). The prevalent diseases resulting from MHV infection are hepatitis, encephalomyelitis, and enteritis. A strict classification, of all MHVs into hepatotropic, neurotropic, and enterotropic strains is not possible, however, since under certain conditions several organs are affected (Table 4) and the type of disease varies to a great extent with the age and genetic background of the host (Sect. 4.1). The role of murine coronaviruses as pathogens of the respiratory tract must also be taken into consideration (*Carthew and Sparrow* 1981). Variants which differ in organ tropism are easily selected in tissue culture or by animal passages.

##### 3.1.1.1 Hepatitis

Several murine coronavirus strains replicate predominantly in liver tissue and induce an acute fatal hepatitis by destruction of parenchymal and Kupffer cells (Table 5; *Piazza* 1969; *Hirano et al.* 1981a). The highly virulent strains MHV-2, MHV-3, and MHV-A 59 cause hepatitis in adult mice. MHV-1 and MHV-S are less virulent but lead eventually to a similar disease. MHV-S is enteropathogenic for young mice whereas most of the other strains (Table 5) cause hepatitis only in newborn mice. MHV-N is virulent only for mice which have been immunosuppressed by cortisone treatment. Viruses isolated from nude mice (MHV-NuU, NuA and Nu66) cause chronic hepatitis in athymic mice (Sect. 4.1.2). However, tissue-culture-adapted MHV-Nu66 and NuA are also hepatotropic for normal mice, indicating an increase in virulence.

Table 5. Origin and characteristics of murine coronavirus strains

Strain <sup>a</sup>	First isolation	Conditions of isolation	Predominant effect on host
MHV-1	<i>Gledhill and Andrewes</i> 1951	Spontaneous hepatic disease (albino mouse, Parkes strain)	Hepatitis
MHV-2 (PRI)	<i>Nelson</i> 1952	Associated with mouse leukemia (Princeton strain)	Hepatitis
MHV-3	<i>Dick et al.</i> 1956	Inoculation of human serum into Swiss mice	Hepatitis, ascites
MHV-A59	<i>Manaker et al.</i> 1961	Inoculation of organ suspensions from mice with Moloney leukemia into Balb/c	Hepatitis, encephalitis
MHV-S	<i>Rowe et al.</i> 1963	Acute diarrhoea of newborn CD-1 mice housed with other strains	Hepatitis, enteritis
MHV-NuU, NuA, Nu66, and other isolates	<i>Hirano et al.</i> 1975 <i>Sebesteny and Hill</i> 1974 <i>Tamura et al.</i> 1976 <i>Ward et al.</i> 1977	Wasting syndrome in nude mice	Hepatitis, encephalitis
MHV-N	<i>Hirano et al.</i> 1979	Feces of healthy carrier mice	Hepatitis in mice treated with cortisone
MHV-LV	<i>Sabesin et al.</i> 1972	Latent infection of cultured mouse liver cells (NCTC 1469)	Hepatitis
MHV-JHM	<i>Cheever et al.</i> 1949	Spontaneous paralyzes of Swiss mice	Encephalomyelitis, hepatitis
MHV-S/CDC <sup>b</sup>	<i>Broderson et al.</i> 1976	Fatal diarrhoea in ICR mice	Enteritis
LIVIM	<i>Kraft</i> 1962	Fatal diarrhoea	Enteritis
MHV-DVIM	<i>Sato et al.</i> 1976	Diarrhoea of infant mice	Enteritis
MHV-D	<i>Ishida et al.</i> 1978	Fatal diarrhoea in suckling mice	Enteritis, hepatitis
Unclassified isolates			
Isolate SD	<i>Burks et al.</i> 1980	Balb/c mice inoculated with human brain (multiple sclerosis) 2-6 months before isolation	Demyelinating encephalomyelitis in mice
Isolate SK	<i>Burks et al.</i> 1980	Subcultures of 17 CI-1 cells originally inoculated with human brain (multiple sclerosis)	-

<sup>a</sup> The strains H747, EHF 210 and EHF 120 mentioned in earlier reports (*McIntosh* 1974) have not been described further; <sup>b</sup> MHV-S/CDC and LIVIM are probably the same strain

### 3.1.1.2 Encephalomyelitis

Murine coronaviruses can cause encephalitis in suckling and adult mice (Table 5; Hirano et al. 1981a). The strain MHV-JHM is especially neurotropic (Cheever et al. 1949; Bailey et al. 1949), causing acute and chronic demyelinating diseases. By the natural intranasal route of infection the virus invades the central nervous system via the olfactory nerve (Goto et al. 1977, 1979), initially replicating in the nasal mucosa and spreading within 6 days to the spinal cord. The outcome of experimental intracerebral infection is similar and necrotic lesions are localized in the hippocampus, olfactory lobes, and periependymal tissues. Demyelination is prevalently confined to the brain stem and spinal cord. In mice which do not develop an acute disease involvement of grey matter is minimal and viral antigen is detectable in white matter up to 28 days post infection (p.i.) (Weiner 1973). Electron microscopic studies demonstrated that oligodendrocytes are the main target cells for JHM virus (Lampert 1973; Powell 1975), but especially in young mice virus can also be detected in neurons and ependymal and endothelial cells, indicating the pantropic nature of this infection (Fleury et al. 1980). Infectious virus can be isolated from animals with acute encephalomyelitis at any time during the disease process.

Mice which do not show clinical signs within the first weeks p.i. or which recover from disease can develop a chronic infection of the central nervous system. Herndon et al. (1975, 1977) observed small foci of active demyelination in Balb/c mice surviving JHM infection for 16 months. Their studies on remyelination in these mice indicated that some of the oligodendroglia cells active in remyelination might be newly generated cells. No information is available about the presence of viral antigens in the central nervous system or the isolation of infectious virus from these animals. In recent experiments Stohlman and Weiner (1981) induced a chronic infection by intracerebral inoculation of JHM virus into 3-month-old C57 BL/6 mice. No clinical diseases were observed, but during the first 12 days p.i. infectious virus was recoverable from liver, brain, and spinal cord. Three months p.i. small foci of viral antigen were detectable in 70% of the animals and by electron microscopy demyelinated lesions were found. At this point immunosuppression did not lead to clinical disease and no infectious virus could be activated or isolated. These results are in contrast to earlier studies by Weiner (1973) who showed that immunosuppression shortly after infection modified a nonfatal infection to an acute encephalomyelitis. This indicates, that the virus-host interactions differ significantly between the acute disease and the chronic infection.

Experiments using cloned JHM virus and temperature-sensitive (TS) mutants of this strain were reported by Haspel et al. (1978). This collection of genetically stable mutants was tested for neurovirulence in Balb/c mice infected at an age of 4 weeks. Whereas wild-type virus was lethal for most animals within 6 days, many TS mutants were found to be less neurovirulent. Fatal diseases were caused only after the inoculation of about 10 000 times higher doses of infectious virus than was needed for the wild-type virus. Some of the mutants induced demyelination in the spinal cord of survivors, and only very few animals died of an acute encephalomyelitis. Further studies revealed that the wild-type virus replicates in both neuronal cells and oligodendrocytes, whereas a TS mutant selectively replicates in oligodendrocytes of the spinal cord (Knobler et al. 1981a, b). This selective tropism of mutants within the central nervous system is probably an important parameter for the ability to induce demyelination without resulting in fatal encephalomyelitis. Similar observations of different neurovirulence between wild-type

and mutant viruses, obtained by mutagens or isolated from persistent infections, have been reported by *Robb et al.* (1979) and *Hirano et al.* (1981b).

*Cheever et al.* (1949) described a delayed course of encephalomyelitis with marked demyelination in rats after inoculation of wild-type JHM virus. These original observations have been recently enlarged upon (*Nagashima et al.* 1978a, b; 1979). The infection of outbred rats (strain Thomae/Chbb) with uncloned JHM virus results in acute or subacute to chronic demyelinating encephalomyelitis which is dependent on the age of the animals, the time of infection, and the virus preparation used. In suckling rats an acute panencephalitis characterized by necrotic lesions in all parts of the central nervous system is found. In weanling rats (age 3–4 weeks), however, a subacute demyelinating encephalomyelitis can occur after an incubation time of several weeks. Demyelinating plaques are sharply demarcated and distributed in the white matter of the central nervous system. A similar disease picture is also found in other rat strains (*Sorensen et al.* 1980). Preliminary results indicate that the susceptibility of inbred rat strains is dependent on genetic traits (*Sorensen et al.* 1981).

Kinetic studies during the development of subacute demyelinating encephalomyelitis in weanling rats infected with JHM virus suggest a biphasic course of the disease (*Wege et al.* 1981b). Within 2 weeks p.i. most of the rats develop a clinically silent acute encephalomyelitis in parallel to the replication of the virus in the central nervous system. After this period virus cannot be recovered from these animals, but histologically marked demyelinating lesions are found prior to the development of a subacute encephalomyelitis. By the time a clinically recognizable disease appears JHM virus is again isolatable.

Occasionally remissions after acute disease are observed (*Sorensen et al.* 1980), and surviving rats sometimes develop a late demyelinating encephalomyelitis after an incubation time of up to 8 months (*Nagashima et al.* 1979). Brain sections of these animals reveal viral antigen and with conventional techniques virus can be isolated. These observations indicate a persistent infection of the brain tissue which is responsible for a chronic disease process.

Whereas wild-type JHM virus varies in its ability to induce subacute and late diseases in weanling rats, TS mutants cause high rates of subacute to chronic diseases. Moreover, suckling rats from immunized mothers can also develop chronic demyelinating diseases if inoculated with TS mutants (*Wege et al.* 1981b). These observations suggest that the development of acute or subacute to chronic demyelinating disease is dependent on the virulence of the virus and host factors such as age, immune status, and genetic background.

### 3.1.1.3 Enteritis

Several enteropathogenic strains of murine coronaviruses have been isolated during the last few years (Table 5). The first agent of this type was investigated by *Kraft* (1962) and termed lethal intestinal virus for infant mice (LIVIM). This agent is probably identical with an enterotropic variant of MHV-S described by *Rowe et al.* (1963) which was later designated MHV-S/CDC by *Broderson et al.* (1976). These viruses cause an acute intestinal disease with a high mortality rate during the first 3 weeks of life. Intestinal contents from moribund mice contain typical coronavirus particles and the virus spreads by contact infection via the nasal or oral route in newborn mice. Diseased animals are dehy-

drated by severe diarrhea. Multinucleated giant cells are found especially in the villi of the small intestines (*Biggers et al. 1964*). By electron microscopy large numbers of coronavirus particles are detectable in intestinal epithelial cells and in macrophages of the lamina propria of the lower intestines (*Hierholzer et al. 1979*). The mothers of affected litters are clinically healthy but necrotic foci are found in the liver. Orally infected adult animals do not develop defined clinical signs and shed virus for about 15 days. Intranasal infection by cell-adapted virus leads to a mild diarrhea without mortality. These mice show no evidence of liver or brain disease. Litters from immune mothers are protected against both natural and experimental infection.

Strain MHV-S/CDC is serologically related to other MHV prototype strains, especially to MHV-S, and to the human strain OC43 (*Hierholzer et al. 1979*). Endemics of LIVIM disease were reported by *Carthew (1977)* and a similar virus-designated MHV D was isolated during a natural outbreak of diarrhea (*Ishida et al. 1978; Ishida and Fujiwara 1979*). MHV-D tends to produce a more systemic infection with the involvement of liver, brain, lung, and lymphoid organs. Another isolate, MHV-DVIM, causes diarrhea in infant mice and is remarkable for its ability to agglutinate red blood cells from rats and mice (*Sato et al. 1976; Sugiyama and Amano 1980*).

### 3.2 Human Coronaviruses

Human coronaviruses are often responsible for common colds and are associated with lower respiratory tract diseases and probably enteric diseases. Essentially two groups of isolates can be distinguished. One group grows in tissue cultures of human origin and is related to the prototype strain 229E; the other group can only be maintained in organ cultures, for example strain OC43. The antigenic relationships are summarized in Table 2.

These viruses are distributed worldwide and antibodies are present with high prevalence (*Monto 1974; Kaye et al. 1975; Gerna et al. 1978*). The antibody response to 229E and OC43 appears to have a cycle of several years, with peaks against each strain every 2–3 years. About 15% of common colds are attributed to coronaviruses (*McIntosh et al. 1970; Larson et al. 1980*). In children pneumonia and other respiratory distress can be caused by coronaviruses (*McIntosh et al. 1973, 1974*). Results of a seroepidemiological survey (*Riski and Hovi 1980*) indicate a possible association of coronaviruses with more severe diseases such as pneumonia, pleurodynia, myocarditis, and meningitis. An agent termed Tettnang virus has been isolated by inoculation of cerebrospinal fluid from patients with various neuropathies and fever into suckling mice (*Málková et al. 1980*). This virus and similar isolates probably represent MHV strains which naturally infect these mice (*Bárdos et al. 1980*).

The development of the common cold was studied in human volunteers inoculated with HCV (*Bradburne et al. 1967; Beare and Reed 1976, McIntosh et al. 1978*). Virus inoculated by nasal drops causes predominantly coryza, but in contrast to rhinoviruses no cough or mucopurulent nasal discharge occurs. Virus shedding decreases sharply within 3–4 days p.i. No evidence for involvement of the lower respiratory tract or intestinal organs was found under experimental conditions. The ciliary epithelium is selectively infected and shedding of antigen-containing cells coincide with the coryza. Rechallenge of volunteers with homologous and heterologous virus 8–12 months p.i. revealed that no cross-protection occurs against the heterologous strain but immunity against homologous virus exists (*Larson et al. 1980*).

The systemic immune response against purified HCV 229E and subcomponents have been quantitated by an enzyme-linked immunoassay (*Kraaijeveld et al. 1980b; Macnaughton et al. 1981*) and the results indicate that most of the antibodies produced during infection react with the peplomer protein of the virus. Only small amounts of antibodies recognize the matrix and nucleocapsid protein. No data on the role of local immunity in protection against respiratory disease are available.

In addition to respiratory diseases some human coronaviruses may be associated with enteric infections (*Caul et al. 1975; Caul and Clarke 1975; Moore et al. 1977; Caul and Egglestone 1977; Schnagl et al. 1978; Moscovici et al. 1980*). However, no information exists on the characterization of these agents and their serological relationships to other human coronaviruses (reviewed by *Macnaughton and Davies 1981*). Coronaviruses have also been associated with an endemic nephropathy (*Apostolov et al. 1975*), but there are no studies revealing an etiological link between nephropathy and coronaviruses.

Two coronavirus strains were recently isolated from mice or mouse tissue culture cells during attempts to isolate viruses from patients with multiple sclerosis (*Burks et al. 1980*). The first isolate, designated SD virus, was isolated after intracerebral inoculation of human brain material into weanling Balb/c mice. Within 2–6 months p.i., mice developed neurological signs and died. From these animals a coronavirus was isolated which replicated in the 17 Cl-1 mouse cell line. The second isolate, designated SK virus, was obtained after 12 subcultures of 17 Cl-1 cells which had been incubated with brain material from a second patient. These isolates are antigenically related to both murine and human coronaviruses and reveal related structural polypeptides as shown by immunoprecipitation (*Gerdes et al. 1981a, b*). At the present time it cannot be decided whether these isolates have been derived from human or mouse tissue, since it is known that murine coronaviruses establish latency in mouse colonies as well as in mouse tissue cultures (*Sabesin 1972*). Additionally, serological studies by *Leinikki et al. (1981)* could not demonstrate any correlation between coronavirus antibody titers and patients with multiple sclerosis or other neurological diseases. Further studies are necessary to show if there is an association of coronaviruses SD and SK with multiple sclerosis.

### 3.3 Avian Coronaviruses

#### 3.3.1 Infectious Bronchitis Virus

Avian IBV infects young chickens, causing an acute respiratory disease leading to high mortality and a decrease in yield and quality of egg production. The disease was first described by *Schalk and Hawn (1931)* and is a very common and worldwide infection in poultry flocks. The virus spreads by both air and the fecal-oral route. At least eight IBV serotypes have been described (*Dawson and Gough 1971; Hopkins 1974*). These serotypes fall into two main groups, the Massachusetts and Connecticut types, which differ in their antigenic relatedness (Sect. 2). The virulence of many different isolates and attenuated vaccine strains differs widely.

The primary target tissue for infection is the trachea (*Purcell and McFerran 1972; Darbyshire et al. 1975*). The virus replicates also in bronchial tissue, lung, kidneys, ovaries, and oviduct. A strong tendency to produce a prolonged infection which results in shedding of virus of several months via the feces has been observed (*Alexander and Gough 1977*). Per-

sistent infection in the presence of high antibody titers is often accompanied by severe nephritis and infectious virus can be reisolated from cecal lymph nodes up to 8 months p.i. (Alexander et al. 1978). Both age and genetic factors influence the outcome of the diseases. In addition, infections by bacteria, mycoplasmas, or infectious bursal disease virus (Rosenberger and Gelb 1978) increase the susceptibility of chickens to IBV.

Resistance to natural infection or experimental challenge after vaccination is probably mediated by the local immune response in the trachea, the nasal mucosa, and the Harderian gland. Detailed knowledge of the humoral or cellular immune mechanisms is not yet available (reviewed by Darbyshire 1981). No correlation between serum-neutralizing antibodies and resistance to reinfection has been shown, but protection seems to be correlated to resistance of the tracheal epithelium to challenge virus and the presence of high titers of hemagglutinin-inhibiting serum antibodies (Gough and Alexander 1979). Secretion of local antibodies can also be demonstrated in organ cultures (Gomez and Raggi 1974; Darbyshire 1980). Cross-protection against challenge by homologous and heterologous virus strains can be measured by observation of the ciliary activity of tracheal explants from vaccinated chickens.

The local antibody response is quite independent of the kinetics of the serum antibody development (Holmes 1973; Leslie and Martin 1973; Watanabe et al. 1975; Chhabra and Goel 1980). Whereas antibodies were first detected 3 days p.i. in the trachea and the titers fall again several weeks later, serum antibodies maintain a persistently high titer. IgG, IgA, and IgM are all detectable in tracheal washings. However, it should be noted that the antibody patterns detected by neutralization tests and enzyme-linked immunoassay (Mockett and Darbyshire 1981) are not identical, indicating that antibodies with different specificity and avidity might be detected by the two techniques. Cell-mediated immunity is demonstrable by specific lymphoblast transformation, but the role in pathogenesis is not yet known (Timms et al. 1980). Maternal antibodies are transferred to chicks during development and may contribute to protection early after hatching (Darbyshire 1981).

Under experimental conditions respiratory symptoms are observed between 2 and 8 days p.i., accompanied by complete absence of ciliary activity. Maximum virus titers are obtained 3 days p.i. The morphology of epithelial cells changes and thickening of mucosa, edema, and lymphocytic infiltration are observed (Hawkes et al. personal communication). Using immunofluorescence small groups of fluorescent cells can still be demonstrated 6 weeks p.i., but the regenerated tracheal epithelium seems to be resistant to destruction by virus infection.

### 3.3.2 Turkey Coronavirus

In the 1950s a virus was suspected to induce a transmissible enteritis of turkeys in Minnesota. By electron microscopic studies, a coronavirus-like agent was identified (Ritchie et al. 1973; Panigrahy et al. 1974) and characterized by physiochemical and morphological criteria (Deshmukh and Pomeroy 1974; Naqui et al. 1975). By immunoelectron-microscopy no cross-reaction of this virus to other coronaviruses was found (Ritchie et al. 1973), but different isolates of TCV are probably antigenically identical (Pomeroy et al. 1975). No tissue culture system is available for propagation of the virus but the virus grows in embryonated eggs (Adams and Hofstad 1971).

The onset of the clinical disease caused by this virus is characterized by depression, loss of appetite, weight loss, and watery diarrhea. The mortality, especially in older poults, is very low. The lesions in experimental and field cases are very similar to the changes caused by mammalian enterotropic coronaviruses and consist in a marked shortening of the villi, loss of microvilli, epithelial desquamation, and hemorrhage in the jejunum, ileum, and cecum. The number of goblet cells decreases, and the appearance of epithelial cells changes from columnar to cuboidal form (*Adams et al. 1972; Desmukh et al. 1976; Gonder et al. 1976*). The lesions appear within 1 day p.i., and recovery and healing begins after about 5 days. No pathological changes are observed in other organs. The number of lymphoid cells in the lamina propria increases and the villus to crypt ratio remains depressed for 10 days. Despite an early regression of histopathological changes, viral antigen can be found by immunofluorescence up to 28 days p.i.

Turkeys that recover are immune throughout their lives (*Pomeroy et al. 1975*). This lifelong immunity is mediated by the secretory IgA antibody barrier (*Nagaraja and Pomeroy 1978, 1980, 1981*). Serum-neutralizing antibody titers are very low, but intestinal secretions and bile contain virus-specific IgA antibodies for at least 6 months p.i. By immunofluorescence, antibody-secreting cells can be localized in the intestines 4–5 months after recovery from disease. In addition to local immunity, peripheral lymphocytes are specifically stimulated by virus antigen. These lymphocytes probably migrate from the intestinal lamina propria to the peripheral blood. Circulating IgA and IgM antibodies appear only during the acute phase of the disease (*Carson et al. 1972*).

### 3.4 Feline Coronaviruses

#### 3.4.1 Feline Infectious Peritonitis Virus

Feline infectious peritonitis virus (FIPV) normally causes widespread inapparent infections of wild and domestic cats, but the infection can also lead to a fatal disease. The disease syndrome was first described by *Holzworth (1963)* and experimentally transmitted from field cases to other cats by *Wolfe and Griesemer (1966)*. A coronavirus was identified as the cause of this disease by both morphological and physicochemical criteria (*Ward 1970; Osterhaus et al. 1976; Pedersen 1976a; Horzinek et al. 1977*). Serologically this agent reveals an antigenic relationship to the TGEV of pigs (*Witte et al. 1977; Reynolds et al. 1977; Pedersen et al. 1978*) and to CCV (*Everman et al. 1981*). Recently a tissue culture system was found which supports the growth of FIPV, and several isolates from field cases are now available (*O'Reilly et al. 1979; Black et al. 1980; Everman et al. 1981; McKeirnan et al. 1981*). It is unknown, whether these isolates are serologically and biologically identical or represent different strains of FIPV.

In nature the virus infects cats and other feline species. Randomly collected sera from wild cats and catteries are often up to 90% positive, indicating a wide distribution of the virus (*Pedersen 1976b; Osterhaus et al. 1977; Loeffler et al. 1978*). However, the incidence of clinical disease is rather low, usually up to 10%. The virus probably causes a high rate of inapparent infections as coronavirus-like particles have been demonstrated in feces of normal cats. Furthermore, the virus replicates in organ cultures of both small intestines and trachea, causing only small ultrastructural changes of absorptive epithelial cells (*Hoshino and Scott 1978, 1980a, b*).

Under epizootic conditions, the incubation time of clinical disease ranges from

several weeks to 4 months (*Hardy and Hurvitz 1971; Robison et al. 1971*). Experimentally transmitted disease occurs after a much shorter incubation time, which may last only 2–3 days from oral inoculation (*Pedersen and Boyle 1980; Everman et al. 1981*).

The clinical onset of disease is rather unspecific and characterized by fever, loss of appetite, and general depression. In typical cases swelling of the abdomen is observed as a result of peritonitis, but this effusive form is not always clinically detectable. In the noneffusive (dry) form localized granulomatous lesions are found. Both the effusive and noneffusive forms are caused by the same virus inoculum (*Hayashi et al. 1980; Everman et al. 1981*). In addition, both neurological symptoms and pleuritis are observed.

After onset of disease, several pathophysiological changes indicate damage to the reticuloendothelial system, liver, and kidneys (*Gouffoux et al. 1975; Weiss et al. 1980*). A depression of several plasma factors and an increase of fibrin-fibrinogen degradation products is accompanied by anemia, neutrophilia, and leukopenia. The amount of gammaglobulins increases significantly and the urine contains elevated levels of proteins, bilirubin, and urobilinogen. The level of liver-specific enzymes is very high. In the effusive form, fibrin is deposited on abdominal organs. Granulomatous inflammatory reactions, vasculitis, and plaques of focal necrosis are scattered through the parenchyma of the liver, kidneys, lung, spleen, and lymph organs. Central nervous system and ocular lesions can also occur, depending on the route of inoculation (*Ward et al. 1974*). Virus can be isolated from peritoneal exudate, organ homogenates, and blood.

Several observations support the concept that FIP might be an immunopathologically mediated disease (*Horzinek et al. 1979*). High levels of antibodies are often detected in field cases, but do not prevent disease (*Pederson et al. 1976b; Horzinek et al. 1978*). Experimentally infected seronegative kittens survive significantly longer and develop a less fulminant disease than seropositive kittens (*Weiss et al. 1980; Pedersen and Boyle 1980*). Moreover, treatment of seronegative kittens with purified anti-FIP IgG results in an aggravation of the disease. In addition, lesions in the liver and serosa of seropositive kittens contain viral antigen, IgG bound to antigen, and complement. In these animals immune complexes can be demonstrated in renal glomeruli tissue (*Jacobse-Geels et al. 1980*).

These findings indicate that the immune response against FIPV infections does not have a protective but maybe a destructive effect. In this context it is of interest that the disease often occurs in association with other virus infections such as feline leukemia, feline panleukopenia (a parvovirus), and feline syncytial virus (*Cotter et al. 1973; Black 1980; McKeirnan et al. 1981*). In such cases an enhancement of the FIPV-induced disease process is observed. It is possible that a preexisting persistent viral infection either leads to a higher susceptibility to FIPV or supports the manifestation of a disease state.

### 3.5 Other Coronaviruses

#### 3.5.1 Bovine Coronavirus

Rotaviruses, parvoviruses and coronaviruses are the main causes of bovine viral diarrhea. *Mebus et al. (1973a, b)* described a coronavirus-like agent associated with diarrhea in young calves (neonatal calf diarrhea coronavirus), which was identified by morphological and physicochemical criteria (*Stair et al. 1972; Sharpee et al. 1976; Dea et al. 1980a, b*). This agent has been adapted to grow in tissue cultures and can easily be transmitted by the oral route.

Other BCV strains cannot be grown on tissue culture and must be maintained by passage in vivo (Doughri et al. 1976; Doughri and Storz 1977). Both these and the tissue-culture-adapted strains (Mebus et al. 1975) cause clinical signs of diarrhea within 24–30 h of inoculation. These symptoms last for 4–5 days and can be lethal. The most severe lesions develop in the small intestines, but the large intestines are also infected. The experimental observation that the addition of trypsin to culture media results in a significant enhancement of virus growth in vitro (Dea et al. 1980b; Storz et al. 1981) suggests that the initiation of infection might be promoted by the action of proteolytic enzymes in the intestinal tract. Virions derived from such trypsin-treated in vitro cultures show shorter surface projections than usual (Storz et al. 1981).

The destruction of the intestinal absorptive epithelium leads rapidly to pathophysiological changes followed by extensive loss of water, sodium, chloride, bicarbonate, and potassium. Metabolism of glucose and lactate becomes severely disturbed and hypoglycemia, lactic acidosis, and an elevated efflux of potassium to the hypovolemic plasma consequently lead to acute shock, heart failure, and death (summarized by Lewis and Phillips 1978; Phillips and Case 1980). Maternal antibodies (IgA and IgM) are transmitted via colostrum to calves and reduce the severeness of disease (Mebus et al. 1976).

More than 50% of bovine sera contain antibodies against the BCV strain LY-138 (Hajer and Storz 1978; Storz and Rott 1980). Furthermore, high percentages of human sera from different sources cross-react with BCV antigens in immunodiffusion, neutralization, and electron microscopic tests (Storz and Rott 1981). The common reactive antigen(s) responsible for neutralization is associated with the virion peplomers, but other studies indicate that additionally internal antigens may be responsible for cross-reactivity (Gerna et al. 1981). A single case of diarrhea caused in man by infection with a BCV has been observed (Storz and Rott 1981), and could indicate that the high degree of reactive antibodies in human sera may result from infection with bovine strains.

### 3.5.2 Canine Coronavirus

Canine coronaviruses usually induce a self-limiting mild gastroenteritis in dogs. CCV has been isolated during an epizootic outbreak of diarrheal disease in military dogs in 1971 (Binn et al. 1975) and during two outbreaks of a highly contagious vomiting and diarrheal disease in the USA (Appel et al. 1979). CCV often occurs in association with canine parvoviruses, which cause a similar but more severe enteric disease (Appel et al. 1979; Helfer-Baker et al. 1980). Serologically, CCV is more predominant among kennel dogs than among family dogs (62%–87% vs 22%) and the incidence of animals seropositive against coronavirus in combination with parvovirus is also much higher in kennel dogs than in family dogs (55.6% vs 7.4%). Epizootic fatal canine enteritis caused by both viruses can also occur among captive coyote populations (Everman et al. 1980).

Canine coronavirus cross-reacts strongly with the porcine TGEV, although it can be serologically differentiated (Reynolds et al. 1980; Garwes and Reynolds 1981). It is also serologically related to FIPV (Everman et al. 1981). CCV cannot infect piglets, but TGEV can be transmitted to dogs without causing clinical signs (Larson et al. 1979).

The oral inoculation of beagle pups leads within 1–7 days to enteritis and diarrhea (Keenan et al. 1976; Takeuchi et al. 1976; Nelson et al. 1979). The lesions, which consist of atrophy and fusion of intestinal villi, are most predominant in the ileum. Virus can be recovered from duodenum, jejunum, ileum, colon, and mesenteric lymph nodes, but no

further spread of virus is detectable. Within 1–2 weeks the diarrhea and histopathological changes disappear and antibodies are detectable. The disease has a more severe course in very young pups than in older pups.

### 3.5.3 Hemagglutinating Encephalomyelitis Virus

Hemagglutinating encephalomyelitis virus (HEV) selectively infects neuronal tissue of pigs and causes a vomiting and wasting disease. The disease was first described as an epizootic outbreak in Canadian swine herds leading to high morbidity in suckling pigs (*Roe and Alexander 1958*). Clinical symptoms consist of vomiting and depression which can lead to death after emaciation and starvation. Additionally, neurological signs of encephalomyelitis appear (*Werdin et al. 1976*). The mortality in young pigs is very high: older litters often survive but remain permanently stunted. Clinical outbreaks are now not so predominant but high percentages of sera contain antibodies, indicating a wide distribution of the virus. The virus exists as a subclinical infection in the presence of maternal antibodies. After the weaning period an active immunity develops (*Andries and Pensaert 1981*).

*Greig et al. (1962)* were the first to isolate HEV. *Mengeling and Cutlip (1976)* demonstrated that both the vomiting disease and encephalomyelitis are caused by the same virus. Pathogenetic studies reveal that after oronasal infection of newborn colostrum-deprived pigs the virus replicates in the respiratory tract, the tonsils, and small intestines and spreads via nerve tracts to the peripheral ganglia nearest to the sites of primary infection (*Andries and Pensaert 1980a, b; Andries et al. 1978*). Vomiting starts 4 days p.i. at the time when the virus is detected in neurons of peripheral ganglia. In the central nervous system the viral antigen is first detected in the sensory nuclei of the trigeminal and vagal nerve located in the medulla oblongata, and then spreads to the brain stem and occasionally to the cerebrum, cerebellum, and spinal cord. The infection of other organs or viremia does not play a significant role in the pathogenesis of the disease.

The local inoculation of the virus intragastrically, intraintestinally, intramuscularly, or into the cerebrospinal fluid always leads to the same clinical signs. However, the distribution of viral antigens is very different depending on the route of inoculation. Thus it seems probable that infection of neurons in different locations could lead to vomiting due to a disturbance of regulatory mechanisms. A further consequence of the infection of neuronal cells is paralysis of the ileum, which leads to emaciation and death by starvation.

### 3.5.4 Transmissible Gastroenteritis Virus

Transmissible gastroenteritis (TGE) is an acute disease affecting pigs of all ages. Especially in pigs under 2 weeks of age, the infection leads after a short incubation period to diarrhea and vomiting, resulting frequently in death within 3–6 days. Older pigs are less severely affected.

The targets for virus replication after oral transmission are absorptive cells of the small intestine (*Pensaert et al. 1970*). However, respiratory infection can also occur, and viruses can persist for prolonged times in lung tissue of older pigs (*Underthal et al. 1974, 1975; Watt 1978*). In the infected intestinal cells necrotic lesions develop and lead to pro-

gressive shortening of the villi. Replacement of the villous epithelial cells begins 18–72 h p.i. by migration of undifferentiated cells from the crypts. These crypt cells are resistant against infection. The epithelial cells of microvilli are important for the digestion of disaccharides and the absorption of monosaccharides and contribute to osmoregulation. Their destruction leads consequently to diarrhea, acidosis, and dehydration (Moon et al. 1978; Shepherd et al. 1979a, b).

A key role in the defence against TGE is played by the local immune response of secretory IgA and IgM production (Stone et al. 1977; Kodama et al. 1980). Recovery from infection might also be enhanced by a strong cell-mediated local immune response (Frederick and Bohl 1976; Shimizu and Shimizu 1979). Interferon (type 1) also appears early in the disease process and is probably secreted by local enterocytes. However, intestinal and serum interferon appear to have little protective effect, since up to 100% of newborn pigs die after infection (La Bonnardiere and Laude 1981).

The transfer of antibodies via colostrum and milk is of practical importance for protection of suckling pigs. Several attenuated virus strains with low virulence are now available for vaccination of pregnant sows (Hess et al. 1977; Saif and Bohl 1979). IgA-secreting lymphocytes are locally stimulated in the lamina propria and invade the mammary glands. The pathological changes after oral infection are strongly dependent on the virulence of the TGEV inoculated, since attenuated strains infect only short parts of the intestines and cause only little atrophy of microvilli (Hess et al. 1977). However, the advantage of restricted growth of attenuated virus is counterbalanced by only a weak stimulation of IgA-secreting cells.

Recently, a coronavirus designated CV-777 was isolated in epizootic diarrhea outbreaks (Pensaert and Debouck 1978). No antigenic relationships to other coronaviruses were detected (Pensaert et al. 1981). The disease course is slower than in TGE and accompanied by less cell destruction (Debouck and Pensaert 1980). CV-777 also replicates to a certain extent in the duodenum and colon and infects crypt cells without destroying their regenerative potential. Another porcine virus unrelated to TGEV was recently described by Horvath and Mocsári (1981).

### 3.5.5 Rat Coronavirus

Two different coronavirus strains have been isolated from rats. Parker's rat coronavirus (RCV) is pathogenic for the respiratory system of rats, whereas the sialodacryoadenitis virus (SDAV) has a pronounced tropism for salivary and lacrimal glands. These viruses replicate on primary rat kidney cells but not on cells susceptible to MHV infection.

Isolation of RCV was achieved by inoculation of lung tissue homogenates of rats into specific pathogen-free animals (Parker et al. 1970). Newborn rats infected intranasally with RCV develop respiratory disease and die within 6–12 days p.i. Rats older than 21 days remain clinically healthy. Histopathological lesions are typical for an interstitial pneumonitis. Virus replication is confined to the mucosal epithelium and lungs. Virus was only exceptionally recovered from salivary and submaxillary glands (Bhatt and Jacoby 1977).

Initially was SDAV detected by electron microscopy in the salivary glands of rats. Infectious virus was subsequently isolated by inoculation of organ homogenates into newborn mice (Jonas et al. 1969). The virus is pathogenic for newborn mice by intracerebral inoculation and causes neuronal degradation. Mouse passaged virus induces lesions

of the salivary and lacrimal glands in rats (*Bhatt et al. 1972*). After intranasal inoculation the virus spreads from the respiratory tract via cervical lymph nodes to submaxillary and parotid salivary glands (*Jacoby et al. 1975*). Within 2 days a rhinitis develops and necrotic lesions spread, especially in the ductal epithelium of the affected glands. The disease is self-limiting and no spread to other organs is detectable. Antibodies are demonstrable within 7 days. In addition to the infection of salivary glands, a keratoconjunctivitis and ophthalmic lesions can be associated with the disease (*Lai et al. 1976; Weisbroth and Peress 1977*). These lesions may be a secondary phenomenon due to bacterial invasion and impediment of the lacrimal glands.

## 4 Pathogenetic Aspects

The development of a disease process depends not only on the biological properties of the infectious agent but also on the host. Such factors as susceptibility, spread of virus through the body, type and severity of disease, and control and elimination of the infectious virus are all host-dependent. In this context, experiments carried out with MHVs have provided important information on the pathogenic mechanisms of coronavirus infections.

### 4.1 The Role of Resistance in The Development of Disease

#### 4.1.1 Acute Infections

##### 4.1.1.1 Murine Hepatitis Virus Type 2

The first evidence for an association of host genes with resistance to MHV infection was reported for MHV-2, which causes a fulminant hepatitis with high lethality in PRI mice but no clinical disease in adult C3H mice. *Bang and Warwick (1960)* observed that peritoneal macrophages derived from PRI mice and cultured *in vitro* are able to replicate MHV-2, whereas no virus growth was detected in cultures of C3H macrophages. Breeding experiments indicated that resistance is inherited by a single recessive gene. These observations suggested that the result of virus infection may depend on the genetically determined ability of cells from the macrophage lineage to replicate the virus. A difference in susceptibility of macrophages was also observed by *Taguchi et al. (1976)* who compared the mouse strains DDD and CDF 1. However, as the following experiments illustrate, a complex network of interactions with other cells of the immune system also influences and modifies the outcome of infection (*Bang 1981*).

*Shif and Bang (1970a, b)* demonstrated that macrophages of PRI and C3H mice absorb and take up the virus equally well although macrophage cultures derived from resistant C3H mice did not produce detectable amounts of infectious virus. The ability of macrophages from both strains to replicate equally well a variant virus which arose during high multiplicity of infection indicates, however, that this genetically determined resistance is not absolute and can be overcome by strain variation. *Weiser and Bang (1976)* bred a mouse strain (C3HSS) which contains the gene for MHV-2 susceptibility from PRI mice but is in all other respects congenic with the resistant C3H strain. *Cody et al. (cit. Bang 1981)* used this new strain to show that whilst MHV-2 can replicate under single-

cycle conditions in macrophage cultures from both resistant and susceptible strains, the virus produced in resistant macrophages is relatively much less infectious for the genetically incompatible system.

Additional experiments have also shown that the resistance of adult C3H mice to hepatitis induced by MHV-2 can be modulated by procedures affecting T-cell functions. Whilst normal mice develop transitory hepatic lesions which do not lead to clinical signs, thymectomized animals are no longer resistant and die with an acute hepatitis (*Sheets et al. 1978*). Macrophage cultures derived from thymectomized animals are, however, still relatively resistant. This indicates that in addition to macrophage resistance, thymus-dependent functions are involved in preventing the disease. Also treatment of C3H mice with hydrocortisone, a steroid which suppresses T-cell functions, abolishes the resistance of C3H mice to MHV-2 (*Gallily et al. 1964*). On the other hand, polyclonal stimulation of lymphocytes by inoculation of concanavalin A into normally susceptible PRI mice induces resistance, again suggesting the involvement of T cell-mediated factors (*Weiser and Bang 1977*). These *in vivo* observations were further supported by experiments with cultured macrophages *in vitro* (*Weiser and Bang 1976, 1977; Taylor et al. 1981*). These authors showed that macrophages from resistant mice can be modulated in their susceptibility by the addition of soluble mediators (lymphokines and interferon) which have been secreted by stimulated lymphocyte cultures.

#### 4.1.1.2 Murine Hepatitis Virus Type 3

In the MHV-3 system, the resistance or susceptibility of animals is correlated with the degree of virus growth in macrophage cultures (*Virelizier and Allison 1976; Macnaughton and Patterson 1980*; reviewed by *Virelizier 1981*). Whilst little or no virus replication occurs in macrophage cultures from A/J mice, a resistant strain (Table 6), the degree of virus replication in macrophage cultures from susceptible and semiresistant strains reflects the pathogenicity of MHV-3 for the particular host. Resistant and susceptible macrophage cultures absorb and incorporate virions to the same extent (*Krzystyniak and Dupuy,*

Table 6. Different diseases induced by MHV-3 in inbred strains of mice. (*Virelizier et al. 1975; Le Prévost et al. 1975a; Yamada et al. 1979*)

Type of disease	Mouse strain inoculated	Age at time of intraperitoneal infection
Lethal, fulminant hepatitis 5–8 days p.i., systemic infection	C57 BL/6, Balb/c, DBA2, and others	6–8 weeks
Lethal hepatitis 6–10 days p.i., selective destruction of T cells	C3H/He	4 weeks
Chronic vasculitis 2–12 months p.i.	C3H/He	5–8 months
Chronic chorioependymitis 2–12 months p.i.	A2G	6–8 weeks
Clearance of virus within 7 days, survival	A/J	Over 10 weeks
Inapparent hepatitis, clearance of virus within 7 days p.i.	DDD	4 weeks

1981). The restriction may affect later stages in virus replication. *Levy et al.* (1981) observed that infection of macrophages from susceptible mouse strains leads to a significant stimulation of the blood coagulation system. This may be an additional parameter which contributes to the development of disease. The genetically determined degree of susceptibility is not only restricted to peritoneal macrophages, since hepatocyte cultures obtained by perfusion of liver also reveal the same type of genetic restriction (*Arnheiter and Haller* 1981). *Levy-Leblond et al.* (1979) have shown that at least two recessive genes are responsible for resistance and that they are associated with the histocompatibility (H2) genes. This suggests that antigen recognition by T-lymphocytes plays a role in virus elimination. Further evidence that impairment of virus replication in macrophages and cooperation with cells of the T-cell lineage are both required for resistance is the observation that resistance can only be transferred if peritoneal cells and adherent spleen cells are inoculated together (*Levy-Leblond and Dupuy* 1977). Additionally, bone marrow cells enhance the protection transferred by spleen cells (*Tardieu et al.* 1980). The host-cell gene functions which regulate the susceptibility for MHV-3 are apparently not important for the replication of other viruses (*Arnheiter and Haller* 1981).

The interferon system represents another important line of defence against MHV-3 infection. Interferon is released by macrophages during the first cycles of virus replication and is induced in both resistant and susceptible mouse strains, with peak titers 1-2 days p.i. (*Virelizier et al.* 1976). Application of an antiserum against virus-induced (type 1) interferon amplifies the disease course in susceptible mice and abolishes resistance if inoculated into resistant strains shortly before virus infection (*Virelizier and Gresser* 1978). No enhancement of disease by anti-interferon globulin can be found in chronically diseased animals (Sect. 4.1.2).

As in the MHV-2 system, immunosuppressive treatments such as thymectomy or treatment with anti Thy-1 serum aggravate the disease – induced by MHV-3 and indicate that T cell-mediated immune mechanisms contribute to resistance (*Dupuy et al.* 1975). Specific antibodies are not of major importance, since transfer of serum from immunized, resistant mice to susceptible mice gives no protection (*LePrevost et al.* 1975). It should also be noted, however, that immunosuppressive treatments impair not only T-cell functions but also the production of virus-induced interferon (*Virelizier et al.* 1979). Interperitoneal inoculation of inactivated *Corynebacterium parvum* together with MHV-3 suppresses the development of disease (*Schindler et al.* 1981). This type of resistance may be due to a nonspecific immune stimulation and activation of macrophages.

This situation is still further complicated by the influence of age of the mouse at the time of MHV-3 infection. For example, in young C3H mice T cells and not macrophages are the primary target for MHV-3 replication (*Yamada et al.* 1979). Thus although C3H mice infected at 4 weeks of age are susceptible, whilst DDD mice are relatively resistant, (Table 6), peritoneal macrophages from both strains support virus growth to the same extent and serum interferon titers are very similar. However, in cultured spleen cells of C3H mice virus growth is associated only with Thy-1-antigen-positive cells.

#### 4.1.1.3 Murine Hepatitis Virus JHM

The third MHV strain that has been studied in some detail is MHV-JHM. *Stohlman and Frelinger* (1978) showed that resistance to JHM virus is a recessive genetic trait, not strongly associated with the H2 complex. The interaction of at least two host genes may be re-

quired. A similar result was reported by *Knobler et al.* (1981b). It seems most important that in infections with MHV-JHM the development of resistance correlates with the maturation of the macrophage cell population. This is indicated by the results of cell transfer experiments (*Stohlman et al.* 1980, *Stohlman and Frelinger* 1981). SJL/J mice at an age of 6 weeks are relatively susceptible compared to older mice of the same strain and resistance can be transferred from old to young mice by peritoneal exudate cells. Depletion of cells bearing markers for T or B cells does not influence the transfer of protection. Virus replication in macrophage cultures of both young and old mice is poor, and macrophages have the ability to suppress virus growth in another susceptible cell culture which is permissive for the virus. This type of suppression is not mediated by interferon. When comparing resistant and susceptible strains, *Knobler et al.* (1981b) found a correlation between virus replication in cultured macrophages and the outcome of disease in vivo. A similar age-dependent resistance associated with the maturation of macrophages was observed by *Taguchi et al.* (1977, 1979b, c, 1980) for MHV-S.

*Pickel et al.* (1981) also studied the development of resistance to MHV-JHM infection during host maturation and found that a mature immune system is not the only requirement for protection. Intraperitoneal infection of C3H mice with JHM virus up to 20 days of age results in an acute fatal disease, whilst mice older than 20 days rapidly acquire resistance. Suckling mice can be rendered resistant by transferring spleen cells from adult mice immunized against JHM virus. Nonimmune spleen cells from adult mice, however, cannot protect after transfer. The transferred normal spleen cells were able to mediate a normal immune response in the immature host.

#### 4.1.2 Chronic Infections

The infection of semiresistant strains (C3H/He and A2G) with MHV-3 results in a persistent infection associated with a chronic neurological disease (*Virelizier et al.* 1975; *Le-Prevost et al.* 1975b). The majority of animals survive the acute stage of infection and develop a progressive chronic disease characterized by incoordination and paresis of one or more limbs (Table 6). The pathological findings in A2G mice consist mainly of a chronic chorioependymitis resulting in hydromelia and hydrocephalus, whereas C3H mice reveal a diffuse vasculitis in kidney, liver, spleen, brain, and spinal cord. Perivascular infiltrations by polymorphonuclear lymphocytes and fibrinoid necrosis develop around veins and arteries. In the central nervous system destruction of myelin and axons can be found, but in contrast to neurotropic MHV strains virus antigen has never been demonstrated in neuronal cells. Antigens and immunoglobulin are, however, detectable in the walls of affected vessels. Inoculation of susceptible mouse strains with organ suspensions from chronically diseased animals induces a fatal acute hepatitis in the recipient. Therefore, persistency in semisusceptible mice is a consequence of the host response and not due to the biological properties of the virus.

The infection of host macrophages by MHV-3 results in modification of the immune response (*Virelizier et al.* 1976; *Lahmy and Virelizier* 1981). Application of antigen (sheep red blood cells) at the time of virus infection results in an immunostimulation against this antigen. However, an immunosuppression occurs if the antigen is inoculated later in infection. In persistent infection, a chronic immunosuppression is observed and may be associated with the continuous release of circulating (type 2) interferon. Furthermore,

prostaglandin(s) produced by stimulated macrophages contribute to immunosuppression. This modification might be one of the mechanisms in the pathogenesis of chronic disease in semisusceptible mice.

Athymic nude mice are another host in which a chronic MHV infection occurs. Several MHVs have been isolated from nude (nu/nu) mice (Table 5). The strain termed MHV-NuU is of low virulence and causes a persistent infection with progressive necrotizing hepatitis and perivascular infiltrations in the lung (*Furuta et al. 1979*). It is not pathogenic for heterozygote nu/+ mice with a Balb/c background. Interestingly, *Tamura et al. (1977, 1978, 1980)* found in infected athymic nu/nu mice an immune response normally not detectable in these animals. After inoculation with thymus-dependent antigen (sheep red blood cells), chronically infected nu/nu mice produce neutralizing antibodies (IgM and IgG) and also produce a secondary response (*Tamura and Fujiwara 1979*). This immunostimulation is thought to involve the differentiation of T-cell precursors. Humoral immunity alone however is not sufficient to prevent disease. Especially the functions of T cells are required for protection (*Kai et al. 1981*). Additionally, during the early phase of disease, the phagocytic activity and number of macrophages is enhanced. Impairment of macrophage functions by the toxic effects of silica inoculation aggravates the disease course and leads to a lethal acute hepatitis (*Tamura et al. 1979, 1980*).

## 4.2 Pathogenicity Associated with Viral Gene Sequences

In the preceding sections the importance of both viral and host factors in determining the outcome of coronaviral infection have been discussed. As a first step in attempting to define the viral gene sequences which might play a major role in the pathogenicity of a particular virus strain, *Lai* and co-workers have compared the genomes of several MHV strains and variants by oligonucleotide fingerprinting (*Lai and Stohlman 1981a,b; Lai et al. 1981*). Most interesting is their comparison of large- and small-plaque variants of MHV-JHM (*Stohlman et al. 1981*). The large-plaque variant (DL) is highly virulent for mice, whereas the small-plaque variant is less virulent and might induce a more extensive demyelination (*Fleming et al.*, personal communication). Each variant contains one unique oligonucleotide sequence which is missing in its counterpart. The unique oligonucleotide of the large-plaque variant is located on the genome about 14–15 kb from the 3' end, whilst the small-plaque variant oligonucleotide maps about 3–5 kb from the 3' end. The respective mRNAs for these oligonucleotides have been tentatively identified. Assuming that the same genes are associated with tropism in tissue culture and pathogenicity in animals, these studies, in conjunction with biochemical studies on viral replication (see *Siddell et al.* pp 131–165), could eventually indicate which protein(s) are important for the different biological properties of such mutant pairs.

A similar approach is based on the observation that MHV-3 is more hepatotropic in comparison to MHV-A59. The genomic RNAs of these two virus strains have been compared by T<sub>1</sub> oligonucleotide mapping and are very similar except for two oligonucleotide sequences. The mRNAs encoded by these sequences are known. Consequently, when the proteins encoded by these mRNAs are identified it may be possible to determine the proteins which are associated with the different pathogenicity of these virus strains.

## 5 Conclusions

It is evident that the framework of host age and genetic background, biological properties of the virus strain, and dose and route of inoculation are the major factors which determine the result of coronavirus infection.

The respiratory and intestinal tract may be the site of primary replication for all coronavirus infections under natural conditions, although the involvement of other target organs is important for the manifestation of disease in most cases. These target cells are hepatocytes and macrophages in the case of different MHV strains and FIPV, ependymal and endothelial cells for MHV-3 in semiresistant hosts, T cells in MHV-3 infection of young C3H mice, ductus cells in the salivary glands for SDAV infection, neurons in the case of infections with HEV and some MHV strains, and oligodendroglia cells for infection with MHV-JHM mutants.

For most viruses causing enteric diseases (BCV, CCV, some MHV strains, TGEV, and TCV) and respiratory diseases (IBV, HCV, and RCV) the pathophysiological events leading to clinical symptoms are almost certainly due to the acute cytotoxic infection of the target cells (epithelial cells of intestines or respiratory epithelium). These infections can be limited by the local immune response resulting in production of secretory antibodies. In enteric infections, maternal antibodies supplied by colostrum and milk are an additional important defence mechanism.

In contrast, many coronaviruses are maintained and spread in the population as inapparent and subclinical infections. Many murine strains have been isolated from clinically healthy animals, and chronic infection by IBV can result in prolonged virus shedding. TGEV can also be carried for a prolonged time. In the case of FIPV, although only a low percentage of animals develop disease there is good reason to believe that many more animals may be infected. In central nervous system infections with MHV-JHM a clinically silent acute encephalitis develops, which may later become a subacute to chronic demyelinating disease.

The sequence of events leading to chronic diseases is unknown. During the pathogenesis of chronic and acute diseases stages of viral persistency can be involved. The result depends on the expression of viral genes, the functional impairment of host cells and the interaction with the host immune response. At the present stage, no experimental data are available on the molecular mechanisms important in the development and maintenance of persistent infections. However, the use of permanent cultures of differentiated cells may be of great use in this respect. For example, neurotropic and non-neurotropic MHV viruses behave very differently in certain neuroblastoma lines (*Lucas et al. 1977*) and viral mutants exist which selectively replicate in oligodendroglia cells (*Knobler et al. 1981a*). Also, persistent infected neuroblastoma cell lines can harbor the virus without any indication of viral antigen expression and other lines can shed virus variants with altered pathogenicity (*Stohlman et al. 1979a, b; Holmes and Behnke 1981; Hirano et al. 1981b*). Such systems will be of value in investigating the physiological impairment of cell functions by virus persistence and as model systems to evaluate mechanisms of viral persistence.

For several murine systems the host genetic background is an essential parameter determining resistance and the outcome of disease. A valuable system for investigating the mechanism of genetic resistance is formed by inbred mice, which are congenic with the exception of the gene(s) responsible for different susceptibility. Many results indicate

that macrophages play a key role in this genetic restriction. The detailed mechanisms for this restriction are as yet difficult to define and cannot be generalized, and the effect of genes which influence the susceptibility may change during host maturation. Mutations within the virus population also have to be considered during prolonged virus-host relationships. Additionally, infection of macrophages and other cells of the immune system clearly modulates the host immune response and influences the outcome of the infection.

The first attempts to define viral genes which influence pathogenicity have been reported. If strain differences are defined in biochemical terms, it may be possible to describe the role of these gene products in pathogenesis. Further work on variants which differ in only few mutations and show clear differences in biological properties can help to elucidate the function of viral genes in pathogenesis.

Coronaviruses are pathogens of economic and clinical importance. Defined experimental systems have been established, especially for murine coronaviruses, which are valuable disease models representative for coronaviral and other diseases of man and animals. We may expect rapid progress to be made in the next few years.

*Acknowledgements.* We gratefully acknowledge the help of many colleagues who provided manuscripts and information. Data were also made available by the WHO Collaborating Centre for Collection and Evaluation of Data on Comparative Virology, University of Munich, Germany. We thank Helga Kriesinger for typing the manuscript. The authors were supported by the Deutsche Forschungsgemeinschaft.

## References

- Adams NR, Ball RA, Hofstad MS (1970) Intestinal lesions in transmissible enteritis of turkeys. *Avian Dis* 14:392-399
- Adams NR, Hofstad MS (1971) Isolation of transmissible enteritis agent of turkeys in avian embryos. *Avian Dis* 15:426-433
- Adams NR, Hofstad MS, Gough PM (1972) Physical and morphological characterization of transmissible enteritis virus of turkeys. *Avian Dis* 16:817-827
- Akashi H, Inaba Y, Miura Y, Sato K, Tokuhisa S, Asagi M, Hayashi Y (1981) Propagation of the Kakegawa strain of bovine coronavirus in suckling mice, rats and hamsters. *Arch Virol* 67:367-370
- Alexander DJ, Gough RE (1977) Isolation of infectious bronchitis virus from experimentally infected chickens. *Res Vet Sci* 23:344-347
- Alexander DJ, Gough RE, Pattison M (1978) A long-term study of the pathogenesis of infection of fowls with three strains of avian infectious bronchitis virus. *Res Vet Sci* 24:228-233
- Andries K, Pensaert MB (1980a) Virus isolation and immunofluorescence in different organs of pigs infected with hemagglutinating encephalomyelitis virus. *Am J Vet Res* 41:215-218
- Andries K, Pensaert MB (1980b) Immunofluorescence studies on the pathogenesis of hemagglutinating encephalomyelitis virus infection in pigs after oronasal inoculation. *Am J Vet Res* 41:1372-1378
- Andries K, Pensaert M (1981) Vomiting and wasting disease. In: ter Meulen V, Siddell S, Wege H (eds) *Biochemistry and biology of coronaviruses*. Plenum Press, New York, pp 399-408
- Andries K, Pensaert M, Callebaut P (1978) Pathogenicity of hemagglutinating encephalomyelitis (vomiting and wasting disease) virus of pigs, using different routes of inoculation. *Zentralbl Veterinaermed [B]* 25:461-468
- Apostolov K, Spasic P, Bojanic N (1975) Evidence of a viral aetiology of endemic (Balkan) nephropathy. *Lancet* II:1271-1273
- Appel MJG, Cooper BJ, Greisen H, Scott F, Carmichael LE (1979) Canine viral enteritis. I. Status report on corona- and parvo-like viral enteritides. *Cornell Vet* 69:123-133

- Arnheiter H, Haller O (1981) Inborn resistance of mice to mouse hepatitis virus type 3 (MHV3). In: ter Meulen V, Siddell S, Wege H (eds) *Biochemistry and biology of coronaviruses*. Plenum Press, New York, pp 409–417
- Bailey OT, Pappenheimer AM, Cheever FS, Daniels JB (1949) A murine virus (JHM) causing disseminated encephalomyelitis with extensive destruction of myelin. II. Pathology. *J Exp Med* 90:195–212
- Bang FB (1981) The use of a genetically incompatible combination of host and virus (MHV) for the study of mechanisms of host resistance. In: ter Meulen V, Siddell S, Wege H (eds) *Biochemistry and biology of coronaviruses*. Plenum Press, New York, pp 359–373
- Bang FB, Warwick A (1960) Mouse macrophages as host cells for the mouse hepatitis virus and the genetic base of their susceptibility. *Proc Natl Acad Sci USA* 46:1065–1075
- Bárdos V, Schwanzler V, Pesko J (1980) Identification of Tettngang virus (possible arbovirus) as mouse hepatitis virus. *Intervirology* 13:275–283
- Bass EP, Sharpee RL (1975) Coronavirus and gastroenteritis in foals. *Lancet* II:822
- Bearé AS, Reed SE (1977) The study of antiviral compounds in volunteers. In: Oxford JS (ed) *Chemoprophylaxis and virus infections of the respiratory tract*. CRC Press, Cleveland, pp 27–35
- Bhatt PN, Jacoby RC (1977) Experimental infection of adult axenic rats with Parker's rat coronavirus. *Arch Virol* 54:345–352
- Bhatt PN, Percy DH, Jones AM (1972) Characterization of the virus of sialodacryoadenitis of rats: a member of the coronavirus group. *J Infect Dis* 126:123–130
- Biggers DC, Kraft LM, Sprinz H (1964) Lethal intestinal virus infection of mice (LIVIM). *Am J Pathol* 45:413–422
- Binn LN, Lazar EC, Keenan KP, Huxson DL, Marchwicki RM, Strano AJ (1975) Recovery and characterisation of a coronavirus from military dogs with diarrhoea. *Proc. 78th Ann. Meeting U.S. Anim. Health Assoc. Roanoke, Va., Oct. 1974*, pp 366–459
- Black JW (1980) Recovery and in vitro cultivation of a coronavirus from laboratory induced cases of feline infectious peritonitis (FIP). *Vet Med Small Anim Clin* 75:811–814
- Bradburne AF, Bynoe ML, Tyrrell DAJ (1967) Effects of a "new" human respiratory virus in volunteers. *Br Med J* 3:767–769
- Broderson JR, Murphy FA, Hierholzer JC (1976) Lethal enteritis in infant mice caused by mouse hepatitis virus. *Lab Anim Sci* 26:824–827
- Burks JS, DeVald BL, Jankovsky LD, Gerdes JC (1980) Two coronaviruses isolated from central nervous system tissue of two multiple sclerosis patients. *Science* 209:933–934
- Carson CA, Naqi CA, Hall CF (1972) Serologic response of turkeys to an agent associated with infectious enteritis (bluecomb). *Appl Environ Microbiol* 23:903–907
- Carthew P (1977) Lethal intestinal virus of infant mice is mouse hepatitis virus. *Vet Rec* 101:465
- Carthew P, Sparrow S (1981) Murine coronaviruses: the histopathology of disease induced by intranasal inoculation. *Res Vet Sci* 30:270–273
- Caul EO, Clarke SKR (1975) Coronavirus propagated from patient with non bacterial gastroenteritis. *Lancet* II:953–954
- Caul EO, Egglestone SI (1977) Further studies on human enteric coronaviruses. *Arch Virol* 54:107–117
- Caul EO, Paver WK, Clarke SKR (1975) Coronavirus particles in faeces from patients with gastroenteritis. *Lancet* I:1192
- Cheever FS, Daniels JB, Pappenheimer AM, Bailey OT (1949) A murine virus (JHM) causing disseminated encephalomyelitis with extensive destruction of myelin. I. Isolation and biological properties of the virus. *J Exp Med* 90:181–194
- Chhabra PC, Goel MC (1980) Normal profile of immunoglobulins in sera and tracheal washings of chickens. *Res Vet Sci* 29:148–152
- Clewley JP, Morser J, Avery RJ, Lomniczi B (1981) Oligonucleotide fingerprinting of the RNA of different strains of infectious bronchitis virus. *Infect Immun* 32:1227–1233
- Collins MS, Alexander DJ (1980) The polypeptide composition of isolated surface projections of avian infectious bronchitis virus. *J Gen Virol* 48:213–217
- Cotter SM, Gilmore CE, Rollins C (1973) Multiple cases of feline leukemia and feline infectious peritonitis in a household. *J Am Vet Med Assoc* 162:1054–1058
- Darbyshire JH (1980) Assessment of cross-immunity in chickens to strains of avian infectious bronchitis virus using tracheal organ cultures. *Avian Pathol* 9:179–184

- Darbyshire JH (1981) Immunity to avian infectious bronchitis virus. In: Rose ME, Payne LN, Freeman BM (eds) *Avian Immunology*. Poultry Science Symposium Series 16, Edinburgh, Brit Poul Science Ltd, pp 205–226
- Darbyshire JH, Cook JKA, Peters RW (1975) Comparative growth kinetic studies on avian infectious bronchitis virus in different systems. *J Comp Pathol* 85:623–630
- Darbyshire JH, Rowell JG, Cook JKA, Peters RW (1979) Taxonomic studies on strains of avian infectious bronchitis virus using neutralisation tests in tracheal organ cultures. *Arch Virol* 61:227–238
- Dawson PS, Gough RE (1971) Antigenic variation in strains of avian infectious bronchitis virus. *Arch Ges Virusforsch.* 34:32–39
- Dea S, Roy RS, Begin ME (1980a) Physicochemical and biological properties of neonatal calf diarrhoea coronaviruses isolated in Quebec and comparison with the Nebraska calf coronavirus. *Am J Vet Res* 41:23–29
- Dea S, Roy RS, Begin ME (1980b) Bovine coronavirus isolation and cultivation in continuous cell lines. *Am J Vet Res* 41:30–38
- Debouck P, Pensaert M (1980) Experimental infection of pigs with a new porcine enteric coronavirus CV 777. *Am J Vet Res* 41:219–223
- Deshmukh DR, Pomeroy BS (1974) Physicochemical characterization of a bluecomb coronavirus of turkeys. *Am J Vet Res* 35:1549–1552
- Deshmukh DR, Sautter JH, Patel BL, Pomeroy BS (1976) Histopathology of fasting and bluecomb disease in turkey poults and embryos experimentally infected with bluecomb disease coronavirus. *Avian Dis* 20:631–640
- Dick GWA, Niven JFS, Gledhill AW (1956) A virus related to that causing hepatitis in mice (MHV). *Br J Exp Pathol* 37:90–97
- Doughri AM, Storz J (1977) Light and ultrastructural pathologic changes in intestinal coronavirus infection of newborn calves. *Zentralbl Veterinärmed [B]* 24:367–385
- Doughri AM, Storz J, Hajer I, Fernando HS (1976) Morphology and morphogenesis of a coronavirus infecting intestinal epithelial cells of newborn calves. *Exp Molec Pathol* 25:355–370
- Doyle LP, Hutchings LM (1946) A transmissible gastroenteritis in pigs. *J Am Vet Assoc* 108:257–259
- Dupuy JM, Levy-Leblond E, Le Prevost C (1975) Immunopathology of mouse hepatitis virus type 3 infection. II. Effects of immunosuppression in resistant mice. *J Immunol* 114:226–230
- Estola T (1967) Sensitivity of suckling mice to various strains of infectious bronchitis virus. *Acta Vet Scand* 8:86–87
- Evermann JF, Foreyt W, Maag-Miller L, Leathers W, McKeirnan AJ, Leamaster B (1980) Acute hemorrhagic enteritis associated with canine coronavirus and parvovirus infections in a captive coyote population. *J Am Vet Med Assoc* 177:784–786
- Evermann JF, Baumgartener L, Ott RL, Davis EV, McKeirnan AJ (1981) Characterization of a feline infectious peritonitis isolate. *Vet Pathol* 18:256–265
- Fleury HJA, Sheppard RD, Bornstein MB, Raine CS (1980) Further ultrastructural observations of virus morphogenesis and myelin pathology in JHM virus encephalomyelitis. *Neuropath Appl Neurobiol* 6:165–179
- Frederick GT, Bohl E (1976) Local and systemic cell-mediated immunity against transmissible gastroenteritis, an intestinal viral infection of swine. *J Immunol* 116:1000–1004
- Furuta T, Goto Y, Tamura T, Kai Ch, Ueda K (1979) Pulmonary vascular lesions in nude mice persistently infected with mouse hepatitis virus. *Jpn J Exp Med* 49:423–428
- Gallily A, Warwick A, Bang FB (1964) Effect of cortisone on genetic resistance to mouse hepatitis virus in vivo and in vitro. *Proc Natl Acad Sci USA* 51:1158–1164
- Garwes DJ, Reynolds DJ (1981) The polypeptide structure of canine coronavirus and its relationship to porcine transmissible gastroenteritis virus. *J Gen Virol* 52:153–157
- Garwes DJ, Lucas MH, Higgins DA, Pike BV, Cartwright SF (1978/1979) Antigenicity of structural components from porcine transmissible gastroenteritis virus. *Vet Microbiol* 3:179–190
- Gerdes JC, Jankovsky LD, DeVald BL, Klein I, Burks JS (1981a) Antigenic relationships of coronaviruses detectable by plaque neutralization, competitive enzyme linked immunoabsorbent assay and immunoprecipitation. In: ter Meulen V, Siddell S, Wege H (eds) *Biochemistry and biology of coronaviruses*. Plenum Press, New York, pp 29–41

- Gerdes JC, Klein I, DeVald B, Burks JS (1981b) Coronavirus isolates SK and SD from multiple sclerosis patients are serologically related to murine coronavirus A59 and JHM and human coronavirus OC43 but not to human coronavirus 229E. *J Virol* 38:231–238
- Gerna G, Cattano E, Cereda P, Revello MG (1978) Seroepidemiological study on human coronavirus OC43 infections in Italy. *Boll Inst Sieroter. Milan.* 57:535–542
- Gerna G, Cereda PM, Revello MG, Cattaneo E, Battaglia M, Gerna MT (1981) Antigenic and biological relationships between human coronavirus OC43 and neonatal calf diarrhoea coronavirus. *J Gen Virol* 54:91–102
- Gledhill AW, Andrewes CA (1951) A hepatitis virus of mice. *Br J Exp Pathol* 32:559–568
- Gomez L, Raggi LG (1974) Local immunity to avian infectious bronchitis in tracheal organ culture. *Avian Dis* 18:346–368
- Gonder E, Patel BL, Pomeroy BS (1976) Scanning electron, light and immunofluorescent microscopy of coronavirus enteritis of turkeys (bluecomb). *Am J Vet Res* 37:1435–1439
- Goto N, Hirano N, Aiuchi M, Hayashi T, Fujiwara K (1977) Nasoencephalopathy of mice infected intranasally with a mouse hepatitis virus, JHM strain. *Jpn J Exp Med* 47:59–70
- Goto N, Takahashi K, Huang KJ, Katami K, Fujiwara K (1979) Giant cell formation in the brain of suckling mice infected with mouse hepatitis virus, JHM strain. *Jpn J Exp Med* 49:169–177
- Gouffaux M, Pastoret PP, Henroteaux M, Massip A (1975) Feline infectious peritonitis proteins of plasma and ascitic fluid. *Vet Pathol* 12:335–348
- Gough RE, Alexander DJ (1979) Comparison of duration of immunity in chickens infected with a live infectious bronchitis vaccine by three different routes. *Res Vet Sci* 26:329–332
- Greig AS, Mitchell D, Corner AH (1962) A hemagglutinating virus producing encephalomyelitis in baby pigs. *Can J Comp Med* 26:49–56
- Haelterman EO (1962) Epidemiological studies of transmissible gastroenteritis of swine. Proc. 66th Annual Meeting US Livestock San. Assoc., pp 305–315
- Hajer I, Storz J (1978) Antigens of bovine coronavirus strain LY-138 and their diagnostic properties. *Am J Vet Res* 39:441–444
- Hardy WD, Hurvitz AI (1971) Feline infectious peritonitis: experimental studies. *J Am Vet Med Assoc* 158:994–1002
- Hasony HJ, Macnaughton MR (1981) Antigenicity of mouse hepatitis strain 3 subcomponents in C57 strain mice. *Arch Virol* 69:33–41
- Haspel VM, Lampert PW, Oldstone MBA (1978) Temperature-sensitive mutants of mouse hepatitis virus produce a high incidence of demyelination. *Proc Natl Acad Sci USA* 75:4033–4036
- Hayashi T, Utsumi F, Takahashi R, Fujiwara K (1980) Pathology of non-effusive type feline infectious peritonitis and experimental transmission. *Jpn J Vet Sci* 42:197–210
- Helfer-Baker C, Evermann JF, McKeirnan AJ, Morrison WB, Slack RL, Miller CW (1980) Serological studies on the incidence of canine enteritis viruses. *Canine Pract* 7:37–42
- Herndon RM, Griffin DE, McCormick U, Weiner LP (1975) Mouse hepatitis virus-induced recurrent demyelination. *Arch Neurol* 32:32–35
- Herndon RM, Price DL, Weiner LP (1977) Regeneration of oligodendroglia during recovery from demyelinating disease. *Science* 195:693–694
- Hess RG, Bachmann PA, Hänichen T (1977) Versuche zur Entwicklung einer Immunprophylaxe gegen die übertragbare Gastroenteritis (TGE) der Schweine. I. Pathogenität des Stammes B1 im Verlaufe von Serienpassagen. *Zentralbl Veterinärmed [B]* 24:753–763
- Hierholzer JC, Broderson JR, Murphy FA (1979) New strain of mouse hepatitis virus as the cause of lethal enteritis in infant mice. *Infect Immun* 24:508–522
- Hirai K, Hitchner SB, Calnek BW (1979) Characterization of a new coronavirus like agent isolated from parrots. *Avian Dis* 23:515–525
- Hirano N, Tamura T, Taguchi F, Ueda K, Fujiwara K (1975) Isolation of low virulent mouse hepatitis virus from nude mice with wasting syndrome and hepatitis. *Jpn J Exp Med* 45:492–496
- Hirano N, Miyajima H, Fujiwara K (1979) Isolation of low virulent mouse hepatitis virus from feces in infected mouse breeding colony. *Jpn J Vet Sci* 41:31–40
- Hirano N, Goto N, Ogawa T, Ono K, Murakami T, Fujiwara K (1980) Hydrocephalus in suckling rats infected intracerebrally with mouse hepatitis virus, MHV A59. *Microbiol Immunol* 24:825–834
- Hirano N, Murakami T, Taguchi F, Fujiwara K, Matumoto M (1981a) Comparison of mouse hepatitis strains for pathogenicity in weanling mice infected by various routes. *Arch Virol* 70:69–73

- Hirano N, Goto N, Makino S, Fujiwara K (1981b) Persistent infection with mouse hepatitis virus JHM strain in DBT cell culture. In: ter Meulen V, Siddell S, Wege H (eds) *Biochemistry and biology of coronaviruses*. Plenum Press, New York, pp 301-308
- Holmes HC (1973) Neutralizing antibody in nasal secretions of chickens following administration of avian infectious bronchitis virus. *Arch Ges Virusforsch* 43:235-241
- Holmes KV, Behnke JN (1981) Evolution of a coronavirus during persistent infection in vitro. In: ter Meulen V, Siddell S, Wege H (eds) *Biochemistry and biology of coronaviruses*. Plenum Press, New York, pp 287-300
- Holzworth J (1963) Some important disorders of cats. *Cornell Vet* 53:157-160
- Hopkins SR (1974) Serological comparisons of strains of infectious bronchitis virus using plaque-purified isolants. *Avian Dis* 18:231-239
- Horváth I, Mocsári E (1981) Ultrastructural changes in the small intestinal epithelium of suckling pigs affected with a transmissible gastroenteritis (TGE)-like disease. *Arch Virol* 68:103-113
- Horzinek MC, Osterhaus ADME, Ellens DJ (1977) Feline infectious peritonitis virus. *Zentralbl Veterinaarmed [B]* 24:398-405
- Horzinek MC, Osterhaus ADME (1978) Feline infectious peritonitis: a coronavirus disease of cats. *J Small Anim Pract* 19:561-568
- Horzinek MC, Osterhaus ADME (1979) The virology and pathogenesis of feline infectious peritonitis. *Arch Virol* 59:1-15
- Hoshino Y, Scott FW (1978) Replication of feline infectious peritonitis virus in organ cultures of feline tissue. *Cornell Vet* 68:411-417
- Hoshino Y, Scott FW (1980a) Immunofluorescent and electron microscopic studies of feline small intestinal organ cultures infected with feline infectious peritonitis virus. *Am J Vet Res* 41:672-675
- Hoshino Y, Scott FW (1980b) Coronavirus-like particles present in the feces of normal cats. *Arch Virol* 63:147-152
- Ishida T, Fujiwara K (1979) Pathology of diarrhea due to mouse hepatitis virus in the infant mouse. *Jpn J Exp Med* 49:33-41
- Ishida T, Taguchi G, Lee Y, Yamada A, Tamura T, Fujiwara K (1978) Isolation of mouse hepatitis virus from infant mice with fatal diarrhea. *Lab Anim Sci* 28:269-276
- Jacobse-Geels H, Daha MR, Horzinek M (1980) Isolation and characterization of feline C3 and evidence for the immune complex pathogenesis of feline infectious peritonitis. *J Immunol* 125:1606-1610
- Jacoby RO, Bhatt PN, Jonas AM (1975) Pathogenesis of sialodacryoadenitis in gnotobiotic rats. *Vet Pathol* 12:196-206
- Jonas AM, Craft J, Black CL, Bhatt PN, Hilding D (1969) Sialodacryoadenitis in rats. A light and electron microscope study. *Arch Pathol* 88:613-622
- Kai C, Tamura T, Fujiwara K (1981) Effect of immune heterozygous spleen cell transfer on resistance to mouse hepatitis virus infection in nude mice. *Microbiol Immunol* 25:1011-1018
- Kapikian AZ (1975) The coronaviruses. *Dev Biol Stand* 28:42-64
- Katami K, Taguchi F, Nakayama M, Goto N, Fujiwara K (1978) Vertical transmission of mouse hepatitis virus infection in mice. *Jpn J Exp Med* 48:481-490
- Kaye HS, Dowdle WR (1975) Seroepidemiologic survey of coronavirus (strain 229E) infections in a population of children. *Am J Epidemiol* 101:238-244
- Kaye HS, Yarbrough WB, Reed CV (1975) Calf diarrhoea coronavirus. *Lancet* II:509
- Kaye HS, Yarbrough WB, Reed CJ, Harrison AK (1977) Antigenic relationship between human coronavirus strain OC43 and hemagglutinating encephalomyelitis virus strain 67N of swine: antibody response in human and animal sera. *J Infect Dis* 135:201-209
- Keenan KP, Jervis HR, Marchwicki RH, Binn LN (1976) Intestinal infection of neonatal dogs with canine coronavirus 1-71: studies by virologic, histologic, histochemical and immunofluorescent techniques. *Am J Vet Res* 37:247-256
- Kersting G, Pette E (1956) Zur Pathohistologie und Pathogenese der experimentellen JHM-Virus-encephalomyelitis des Affen. *Dtsch Zeitschr f Nervenheilkd* 174:283-304
- Knobler RL, Dubois-Dalcq M, Haspel MV, Claysmith AP, Lampert PW, Oldstone MBA (1981a) Selective localization of wild type and mutant mouse hepatitis virus (JHM strain) antigens in CNS tissue by fluorescence, light and electron microscopy. *J Neuroimmunol* 1:81-92

- Knobler RL, Haspel MV, Oldstone MBA (1981b) Mouse hepatitis virus type-4 (JHM strain) induced fatal nervous system disease, part I (Genetic control and the murine neuron as the susceptible site of disease). *J Exp Med* 133:832-843
- Kodama Y, Ogata M, Shimizu Y (1980) Characterization of immunoglobulin A antibody in serum of swine inoculated with transmissible gastroenteritis virus. *Am J Vet Res* 41:740-745
- Kraaijeveld CA, Madge MH, Macnaughton MR (1980a) Enzyme-linked immunosorbent assay for coronaviruses HCV 229E and MHV 3. *J Gen Virol* 49:83-89
- Kraaijeveld CA, Reed SE, Macnaughton MR (1980b) Enzyme-linked immunosorbent assay for detection of antibody in volunteers experimentally infected with human coronavirus 229E group viruses. *J Clin Microbiol* 12:493-497
- Kraft LM (1962) An apparently new lethal virus disease of infant mice. *Science* 137:282-283
- Krystyniak K, Dupuy JM (1981) Early interaction between mouse hepatitis virus 3 and cells. *J Gen Virol* 57:53-61
- La Bonnardiére C, Laude H (1981) High interferon titer in newborn pig intestine during experimentally induced viral enteritis. *Infect Immun* 32:28-31
- Lahmy C, Virelizier JL (1981) Role of prostaglandins in the suppression of antibody production by mouse hepatitis virus infection. *Ann Immunol (Paris)* 132C:101-105
- Lai MMC, Stohman SA (1981a) Comparative analysis of RNA genome of mouse hepatitis viruses. *J Virol* 38:661-670
- Lai MMC, Stohman SA (1981b) Genome structure of mouse hepatitis virus: comparative analysis by oligonucleotide mapping. In: ter Meulen V, Siddell S, Wege H (eds) *Biochemistry and biology of coronaviruses*. Plenum Press, New York, pp 69-82
- Lai MMC, Brayton PR, Armen RC, Patton CD, Stohman SA (1981) Mouse hepatitis virus A59 messenger RNA structure and genetic localization of the sequence divergence from the hepatotropic strain MHV-3. *J Virol* 39:823-834
- Lai YL, Jacoby RO, Bhatt PN, Jonas AM (1976) Keratoconjunctivitis associated with sialodacryoadenitis in rats. *Invest Ophthalmol* 15:538-541
- Lampert PW, Sims JK, Kniazeff AJ (1973) Mechanism of demyelination in JHM virus encephalomyelitis. *Electron microscopic studies*. *Acta Neuropathol (Berlin)* 24:76-85
- Larson DJ, Morehouse LG, Solorzano RF, Kinden DA (1979) Transmissible gastroenteritis in dogs: Experimental intestinal infection with transmissible gastroenteritis virus. *Am J Vet Res* 40:477-486
- Larson HE, Reed SE, Tyrrell DAJ (1980) Isolation of rhinoviruses and coronaviruses from 38 colds in adults. *J Med Virol* 5:221-229
- Leinikki PO, Holmes KV, Shekarchi J, Iivanainen M, Madden D, Sever JL (1981) Coronavirus antibodies in patients with multiple sclerosis. In: ter Meulen V, Siddell S, Wege H (eds) *Biochemistry and biology of coronaviruses*. Plenum Press, New York, pp 323-326
- Le Prévost C, Levy-Leblond B, Virelizier JL, Dupuy JM (1975a) Immunopathology of mouse hepatitis virus type 3 infection. I. Role of humoral and cell-mediated immunity in resistance mechanism. *J Immunol* 117:221-225
- Le Prévost C, Virelizier JL, Dupuy JM (1975b) Immunopathology of mouse hepatitis virus type 3 infection. III. Clinical and virologic observation of a persistent viral infection. *J Immunol* 115:640-645
- Leslie GA, Martin LN (1973) Studies on the secretory immunologic system of fowl. III. Serum and secretory IgA of the chicken. *J Immunol* 110:1-9
- Levy GA, Leibowitz JL, Edgington TS (1981) Induction of monocyte procoagulant activity by murine hepatitis type 3 parallels disease susceptibility in mice. *J Exp Med* 154:1150-1163
- Levy-Leblond E, Dupuy JM (1977) Neonatal susceptibility to MHV-3 infection in mice. I. Transfer of resistance. *J Immunol* 118:1219-1222
- Levy-Leblond E, Oth D, Dupuy JM (1979) Genetic study of mouse sensitivity to MHV-3 infection: Influence of the H-2 complex. *J Immunol* 122:1359-1362
- Lewis LD, Phillips RW (1978) Pathophysiologic changes due to coronavirus induced diarrhea in the calf. *J Am Vet Med Assoc* 173:636-641
- Loeffler DG, Ott RL, Evermann JF, Alexander JE (1978) The incidence of naturally occurring antibodies against feline infectious peritonitis in selected cat populations. *Feline Pract* 8:43-45
- Lucas A, Flintoff W, Anderson R, Percy D, Coulter M, Dales S (1977) In vivo and in vitro models

- of demyelinating diseases: tropism of the JHM strain of murine hepatitis virus for cells of glial origin. *Cell* 12:553-560
- Macnaughton MR (1981) Structural and antigenic relationships between human, murine and avian coronaviruses. In: ter Meulen V, Siddell S, Wege H (eds) *Biochemistry and biology of coronaviruses*. Plenum Press, New York, pp 19-28
- Macnaughton MR, Davies HA (1981) Human enteric coronaviruses. *Arch Virol* 70:301-313
- Macnaughton MR, Hasony HJ, Madge H, Reed S (1981) Antibody to virus components in volunteers experimentally infected with human coronaviruses 229E group viruses. *Infect Immun* 31:845-849
- Macnaughton MR, Patterson S (1980) Mouse hepatitis virus strain 3 infection of C57, A/Sn and A/J strain mice and their macrophages. *Arch Virol* 66:71-75
- Málková D, Holubová J, Kolman JM, Lobkovic F, Pohlreichová L, Zikmundová L (1980) Isolation of Tettngang coronavirus from man. *Acta Virol* 24:363-366
- Manaker RA, Piczak CV, Miller AA, Stanton MF (1961) A hepatitis virus complicating studies with mouse leukemia. *J Natl Cancer Inst* 27:29-44
- McIntosh K (1974) Coronaviruses: a comparative review. *Curr Top Microbiol Immunol* 63:85-129
- McIntosh K, Becker WB, Chanock RM (1967) Growth in suckling mouse brain of "IBV-like" viruses from patients with upper respiratory tract disease. *Proc Natl Acad Sci USA* 58:2268-2273
- McIntosh K, Kapikian AZ, Hardison KA, Hartley JW, Chanock RM (1969) Antigenic relationships among coronaviruses of man and between human and animal coronaviruses. *J Immunol* 102:1109-1118
- McIntosh K, Kapikian AZ, Turner HC, Hartley JW, Parrot RH, Chanock RM (1970) Seroepidemiologic studies of coronavirus infection in adults and children. *Am J Epidemiol* 91:585-592
- McIntosh K, Ellis EF, Hoffman LS, Lybass TG, Eller JJ, Fulginiti VA (1973) The association of viral and bacterial respiratory infections with exacerbations of wheezing in young asthmatic children. *J Pediatr* 82:578-593
- McIntosh K, Chao RK, Krause HE, Wasil R, Mocega HE, Mufson MA (1974) Coronavirus infection in acute lower respiratory tract disease of infants. *J Infect Dis* 139:502-507
- McIntosh K, McQuillin J, Reed SE, Gardener PS (1978) Diagnosis of human coronavirus infection by immunofluorescence: method and application to respiratory disease in hospitalized children. *J Med Virol* 2:341-346
- McKeirnan AJ, Evermann JF, Hargis A, Miller LM, Ott RE (1981) Isolation of feline coronaviruses from two cats with diverse disease manifestations. *Feline Pract* 11:16-20
- Mebus CA, Stair EL, Rhodes MB, Twiehaus MJ (1973a) Pathology of neonatal calf diarrhoea induced by a coronavirus-like agent. *Vet Pathol* 10:45-64
- Mebus CA, Stair EL, Rhodes MB, Twiehaus MJ (1973b) Neonatal calf diarrhoea: propagation, attenuation, and characteristics of a coronavirus-like agent. *Am J Vet Res* 34:145-150
- Mebus CA, Newman LE, Stair EL (1975) Scanning electron, light and immunofluorescent microscopy of intestine of gnotobiotic calf infected with calf diarrhea coronavirus. *Am J Vet Res* 36:1719-1725
- Mebus CA, Torres-Medina A, Twiehaus MJ (1976) Immune response to orally administered calf reovirus-like agent and coronavirus vaccine. *Dev Biol Stand* 33:396-403
- Mengeling WL, Cutlip RC (1976) Pathogenicity of field isolates of hemagglutinating encephalomyelitis virus for neonatal pigs. *J Am Vet Med Assoc* 128:236-239
- Mockett A, Darbyshire JH (1981) Comparative studies with an enzyme-linked immunosorbent assay (ELISA) for antibodies to avian infectious bronchitis virus. *Avian Pathol* 10:1-10
- Monto AS (1974) Medical Reviews: Coronaviruses. *Yale J Biol Med* 47:234-251
- Moon HW (1978) Mechanisms in the pathogenesis of diarrhea: a review. *J Am Vet Med Assoc* 172:443-448
- Moore B, Lee P, Hewish M, Dixon B, Mukherjee T (1977) Coronaviruses in training centre for intellectually retarded. *Lancet* I:261
- Moscovici O, Chany C, Lebon P, Rousset S, Laporte (1980) Association d'infection à coronavirus avec l'entérocolite hémorragique du nouveau-né. *CR Acad Sc [D] (Paris)* 290:869-872
- Nagaraja KV, Pomeroy BS (1980) Cell-mediated immunity against turkey coronaviral enteritis (bluecomb). *Am J Vet Res* 41:915-917
- Nagaraja KV, Pomeroy BS (1978) Secretory antibodies against turkey coronaviral enteritis. *Am J Vet Res* 39:1463-1465

- Nagaraja KV, Pomeroy BS (1981) Immunofluorescent studies on localization of secretory immunoglobulins in the intestines of turkeys recovered from turkey coronavirus enteritis. *Am J Vet Res* 41:1283-1284
- Nagashima K, Wege H, ter Meulen V (1978a) Early and late CNS-effects of coronavirus infection in rats. *Adv Exp Med Biol* 100:395-409
- Nagashima K, Wege H, Meyermann R, ter Meulen V (1978b) Corona virus induced subacute demyelinating encephalomyelitis in rats. A morphological analysis. *Acta Neuropathol (Berlin)* 44:63-70
- Nagashima K, Wege H, Meyermann R, ter Meulen V (1979) Demyelinating encephalomyelitis induced by a long-term corona virus infection in rats. *Acta Neuropathol (Berlin)* 45:205-213
- Nagy E, Lomniczi B (1979) Polypeptide patterns of infectious bronchitis virus serotypes fall into two categories. *Arch Virol* 61:341-345
- Naqi SA, Panigrahy B, Hall CF (1975) Purification and concentration of viruses associated with transmissible (coronaviral) enteritis of turkeys (bluecomb). *Am J Vet Res* 36:548-552
- Nelson DT, Eustin SL, McAdarach JP, Stotz I (1979) Lesions of spontaneous canine viral enteritis. *Vet Pathol* 16:680-686
- Nelson JB (1952) Acute hepatitis associated with mouse leukemia. I. Pathological features and transmission of the disease. *J Exp Med* 96:293-303
- O'Reilly KJ, Fishman B, Hitchcock LM (1979) Feline infectious peritonitis. Isolation of a coronavirus. *Vet Rec* 104:348
- Osterhaus ADME, Horzinek MC, Ellens DJ (1976) Untersuchungen zur Ätiologie der feline infektiösen Peritonitis. *Berl Münch Tierärztl Wochenschr* 89:135-137
- Osterhaus ADME, Horzinek MC, Reynolds DJ (1977) Seroepidemiology of feline infectious peritonitis virus infections using transmissible gastroenteritis virus as antigen. *Zentralbl Veterinaermed [B]* 24:835-841
- Osterhaus ADME, Horzinek MC, Wirahadiredja RMS (1978a) Feline infectious peritonitis (FIP) virus. II. Propagation in suckling mouse brain. *Zentralbl Veterinaermed [B]* 25:301-307
- Osterhaus ADME, Horzinek MC, Wirahadiredja RMS, Kroo A (1978b) Feline infectious peritonitis (FIP) virus. IV. Propagation in suckling rat and hamster brain. *Zentralbl Veterinaermed [B]* 25:816-825
- Panigrahy B, Naqi SA, Hall CE (1973) Isolation and characterization of viruses associated with transmissible enteritis (bluecomb) of turkeys. *Avian Dis* 17:430-438
- Parker JC, Cross SS, Rowe WP (1970) Rat coronavirus (RCV): a prevalent, naturally occurring pneumotropic virus of rats. *Arch Ges Virusforsch* 31:293-302
- Pedersen NC (1976a) Morphologic and physical characteristics of feline infectious peritonitis virus and its growth in autochthonous peritoneal cell cultures. *Am J Vet Res* 37:567-572
- Pedersen NC (1976b) Serological studies of naturally occurring feline infectious peritonitis. *Am J Vet Res* 37:1449-1453
- Pedersen NC, Boyle BS (1980) Immunologic phenomena in the effusive form of feline infectious peritonitis. *Am J Vet Res* 41:868-876
- Pedersen NC, Ward J, Mengeling WL (1978) Antigenic relationship of the feline infectious peritonitis virus to coronaviruses of other species. *Arch Virol* 58:45-53
- Pensaert M, Callebaut P (1978) The coronaviruses: clinical and structural aspects with some practical implications. *Ann Méd Vét* 122:301-322
- Pensaert M, Haelterman EO, Burnstein T (1970) Transmissible gastroenteritis of swine: virus-intestinal cell interactions. *Arch Ges Virusforsch* 31:321-334
- Pensaert MB, Debouck P (1978) A new coronavirus-like particle associated with diarrhoea in swine. *Arch Virol* 58:243-247
- Pensaert MB, Debouck P, Reynolds DJ (1981) An immunoelectron microscopic and immunofluorescent study on the antigenic relationship between the coronavirus like agent, CV777, and several coronaviruses. *Arch Virol* 68:45-52
- Phillips RW, Case GL (1980) Altered metabolism, acute shock and therapeutic response in a calf with coronavirus induced diarrhea. *Am J Vet Res* 41:1039-1044
- Piazza M (1969) Experimental viral hepatitis. Thomas, Springfield Illinois
- Pickel K, Müller MA, ter Meulen V (1981) Analysis of age-dependent resistance to murine coronavirus JHM infection in mice. *Infect Immun* 34:375-387

- Pomeroy BS, Larsen CT, Deshmukh DR, Patel BL (1975) Immunity to transmissible (coronaviral) enteritis of turkeys (bluecomb). *Am J Vet Res* 36:553-555
- Powell HC, Lampert PW (1975) Oligodendrocytes and their myelin-plasma membrane connections in JHM mouse hepatitis virus encephalomyelitis. *Lab Invest* 33:440-445
- Purcell DA, McFerran JB (1972) The histopathology of infectious bronchitis in the domestic fowl. *Res Vet Sci* 13:116-122
- Reynolds DJ, Garwes DJ (1979) Virus isolation and serum antibody responses after infection of cats with transmissible gastroenteritis virus. *Arch Virol* 60:161-166
- Reynolds DJ, Garwes DJ, Gaskell CJ (1977) Detection of transmissible gastroenteritis virus neutralizing antibody in cats. *Arch Virol* 55:77-86
- Reynolds DJ, Garwes DJ, Lucey S (1980) Differentiation of canine coronavirus and porcine transmissible gastroenteritis virus by neutralization with canine, porcine and felina sera. *Vet Microbiol* 5:283-290
- Riski H, Hovi T (1980) Coronavirus infections of man associated with diseases other than the common cold. *J Med Virol* 6:259-265
- Ritchie AE, Deshmukh DR, Larsen CT, Pomeroy BS (1973) Electron microscopy of coronavirus-like particles characteristic of turkey bluecomb disease. *Avian Dis* 17:546-558
- Robb JA, Bond CW (1979) Coronaviridae. In: Fraenkel-Conrat H, Wagner RR (eds) *Comprehensive virology*. Plenum Press, New York 14:193-247
- Robb JA, Bond CW, Leibowitz JL (1979) Pathogenic murine Coronaviruses III. Biological and biochemical characterization of temperature sensitive mutants of JHMV. *Virology* 94:385-399
- Robison RL, Holzworth J, Gilmore CE (1971) Naturally occurring feline infectious peritonitis: signs and clinical diagnosis. *J Am Vet Med Assoc* 158:981-986
- Roe CK, Alexander TJL (1958) A disease of nursing pigs previously unreported in Ontario. *Can J Comp Med* 22:305-307
- Rosenberger JK, Gelb J (1978) Response to several avian respiratory viruses as affected by infectious bursal disease virus. *Avian Dis* 22:95-105
- Rowe WP, Hartley JW, Capps UJ (1963) Mouse hepatitis virus infection as a highly contagious, prevalent, enteric infection of mice. *Proc Soc Exp Biol Med* 112:161-165
- Sabesin SM (1972) Isolation of a latent murine hepatitis virus from cultured mouse liver cells. *Am J Gastroenterol* 58:259-274
- Saif LJ, Bohl EH (1979) Passive immunity in transmissible gastroenteritis of swine. *Am J Vet Res* 40:115-117
- Sato K, Maru M, Wata T (1976) Some characteristics of corona-like virus isolated from infant mice with diarrhea and inflammatory submaxillary gland of rats (in Japanese). *Virus* 26:97
- Sato K, Inaba Y, Kurogi H, Takahashi E, Ito Y, Goto Y, Omori T, Matumoto M (1977) Physicochemical properties of calf coronavirus. *Vet Microbiol* 2:73-81
- Schalk AE, Hawn MC (1931) An apparently new respiratory disease of baby chicks. *J Am Vet Med Assoc* 78:413-422
- Schindler L, Streissle H, Kirchner H (1981) Protection of mice against mouse hepatitis virus by *Corynebacterium parvum*. *Infect Immun* 32:1128-1131
- Schmidt OW, Kenny GE (1981) Immunogenicity and antigenicity of human coronavirus 229E and OC43. *Infect Immun* 32:1000-1006
- Schnagl RD, Holmes IH, Mackay-Scollay EM (1978) Coronavirus-like particles in aboriginals and non-aboriginals in Western Australia. *Med J Australia* 1:307-310
- Sebesteny A, Hill AC (1974) Hepatitis and brain lesions due to mouse hepatitis virus accompanied by wasting in nude mice. *Lab Anim* 8:317-326
- Sharpee RL, Mebus CA, Bass EP (1976) Characterization of a calf diarrheal coronavirus. *Am J Vet Res* 37:1031-1041
- Sheets P, Shah K V, Bang FB (1978) Mouse hepatitis virus (MHV) infection in thymectomized C<sub>3</sub>H mice. *Proc Soc Exp Biol Med* 159:34-38
- Shepherd RW, Butler DG, Cutz E, Gall DG, Hamilton JR (1979a) The mucosal lesion in viral enteritis: extent and dynamics of the epithelial response to virus invasion in transmissible gastroenteritis in piglets. *Gastroenterol* 75:770-777
- Shepherd RW, Gall DG, Butler DG, Hamilton JR (1979b) Determinants of diarrhea in viral enteritis. The role of ion transport and epithelial changes in the ileum in transmissible gastroenteritis in piglets. *Gastroenterol* 76:20-24

- Shif I, Bang FB (1970a) In vitro interaction of mouse hepatitis virus and macrophages from genetically resistant mice. I. Absorption of virus and growth curve. *J Exp Med* 131:843-850
- Shif I, Bang FB (1970b) In vitro interaction of mouse hepatitis virus and macrophages from genetically resistant mice. II. Biological characterization of a variant virus (MHV C<sub>3</sub>H) isolated from stocks of MHV (PRI). *J Exp Med* 131:851-862
- Shimizu M, Shimizu Y (1979) Demonstration of cytotoxic lymphocytes to virus infected target cells in pigs inoculated with transmissible gastroenteritis virus. *Am J Vet Res* 40:208-213
- Sorensen O, Percy D, Dales S (1980) In vivo and in vitro models of demyelinating diseases. III. JHM virus infection of rats. *Arch Neurol* 37:478-484
- Sorensen O, Coulter-Mackie M, Percy D, Dales S (1981) In vivo and in vitro models of demyelinating diseases. In: ter Meulen V, Siddell S, Wege H (eds) *Biochemistry and biology of coronaviruses*, Plenum Press, New York, pp 271-286
- Stair EL, Rhodes MB, White RG (1972) Neonatal calf diarrhea: purification and electron microscopy of a coronavirus-like agent. *Am J Vet Res* 33:1147-1156
- Stohlman SA, Frelinger JA (1978) Resistance to fatal central nervous system disease by mouse hepatitis virus strain JHM. I. Genetic analysis. *Immunogen* 6:277-281
- Stohlman SA, Weiner LP (1981) Chronic central nervous system demyelination in mice after JHMV virus infection. *Neurology* 31:38-44
- Stohlman SA, Sakaguchi A, Weiner LP (1979a) Rescue of a positive stranded RNA virus from antigen negative neuroblastoma cells. *Life Sci* 24:1029-1036
- Stohlman SA, Sakaguchi AY, Weiner LP (1979b) Characterization of the cold-sensitive murine hepatitis virus mutants rescued from latently infected cells by cell fusion. *Virology* 98:448-455
- Stohlman SA, Frelinger JA, Weiner LP (1980) Resistance to fatal central nervous system disease by mouse hepatitis virus, strain JHM. II. Adherent cell mediated protection. *J Immunol* 124:1733-1739
- Stohlman SA, Frelinger JA (1981) Macrophages and resistance to JHM virus. In: ter Meulen V, Siddell S, Wege H (eds) *Biochemistry and biology of coronaviruses*. Plenum Press, New York, pp 387-398
- Stone SS, Kemeny LS, Woods RD, Jensen MT (1977) Efficacy of isolated colostral IgA, IgG, IgM(A) to protect neonatal pigs against the coronavirus of transmissible gastroenteritis. *Am J Vet Res* 38:1285-1288
- Storz J, Rott R (1980) Über die Verbreitung der Coronavirusinfektion bei Rindern in ausgewählten Gebieten Deutschlands: Antikörpernachweis durch Mikroimmundiffusion und Neutralisation. *Dtsch Tierärztl Wochenschr* 87:252-254
- Storz J, Rott R (1981) Reactivity of antibodies in human serum with antigens of an enteropathogenic bovine coronavirus. *MMI* 169:169-178
- Storz J, Rott R, Kaluza G (1981) Enhancement of plaque formation and cell fusion of an enteropathogenic coronavirus by trypsin treatment. *Infect Immun* 31:1214-1222
- Sugiyama K, Amano Y (1980) Hemagglutination and structural polypeptides of a new coronavirus associated with diarrhea in infant mice. *Arch Virol* 66:95-105
- Taguchi F, Hirano N, Kiuchi Y, Fujiwara K (1976) Difference in response to mouse hepatitis virus among susceptible mouse strains. *Jpn J Microbiol* 20:293-302
- Taguchi F, Aiuchi M, Fujiwara K (1977) Age-dependent response of mice to a mouse hepatitis virus, MHV-S. *Jpn J Exp Med* 47:109-115
- Taguchi F, Yamada A, Fujiwara K (1979a) Asymptomatic infection of mouse hepatitis virus in the rat. *Arch Virol* 59:275-279
- Taguchi F, Yamada A, Fujiwara K (1979b) Factors involved in the age-dependent resistance of mice infected with low-virulence mouse hepatitis virus. *Arch Virol* 62:333-340
- Taguchi F, Goto Y, Aiuchi M, Hayashi T, Fujiwara K (1979c) Pathogenesis of mouse hepatitis virus infection. The role of nasal epithelial cells as a primary target of low-virulence virus, MHV-S. *Microbiol Immunol* 23:249-262
- Taguchi F, Yamada A, Fujiwara K (1980) Resistance to highly virulent mouse hepatitis virus acquired by mice after low-virulence infection: enhanced antiviral activity of macrophages. *Infect Immun* 29:42-49
- Takahashi K, Hirano N, Goto N, Fujiwara K (1980) Massive cerebral cortical necrosis in suckling rats inoculated intracerebrally with mouse hepatitis virus, A59 strain. *Jpn J Vet Sci* 42:311-321

- Takeuchi A, Binn LN, Jervis HR, Keenan KP, Hildebrandt PK, Valas RB, Bland FF (1976) Electron microscope study of experimental enteric infection in neonatal dogs with a canine coronavirus. *Lab Invest* 34:539-549
- Tamura T, Fujiwara K (1979) IgM and IgG response to sheep red blood cells in mouse hepatitis virus infected nude mice. *Microbiol Immunol* 23:177-183
- Tamura T, Ueda K, Hirano N, Fujiwara K (1976) Response of nude mice to a mouse hepatitis virus isolated from a wasting nude mouse. *Jpn J Exp Med* 46:19-30
- Tamura T, Taguchi F, Ueda K, Fujiwara K (1977) Persistent infection with mouse hepatitis virus of low virulence in nude mice. *Microbiol Immunol* 21:683-691
- Tamura T, Machii K, Ueda K, Fujiwara K (1978) Modification of immune response in nude mice infected with mouse hepatitis virus. *Microbiol Immunol* 22:557-564
- Tamura T, Kai Ch, Sakaguchi A, Ishida T, Fujiwara K (1979) The role of macrophages in the early resistance to mouse hepatitis virus infection in nude mice. *Microbiol Immunol* 23:965-974
- Tamura T, Sakaguchi A, Kai Ch, Fujiwara K (1980) Enhanced phagocytic activity of macrophages in mouse hepatitis virus infected nude mice. *Microbiol Immunol* 24:243-247
- Tardieu M, Hery C, Dupuy JM (1980) Neonatal susceptibility to MHV-3 infection in mice. II. Role of natural effector marrow cells in transfer of resistance. *J Immunol* 124:418-423
- Taylor CE, Weiser WY, Bang FB (1981) In vitro macrophage manifestation of cortisone induced decrease in resistance to mouse hepatitis virus. *J Exp Med* 153:732-737
- Timms LM, Bracewell CD, Alexander DJ (1980) Cell mediated and humoral immune response in chickens infected with avian infectious bronchitis. *Br Vet J* 136:349-356
- Traavik T, Mehl R, Kjeldsberg E (1977) "Runde" virus, a coronavirus-like agent associated with seabirds and ticks. *Arch Virol* 55:25-38
- Tyrrell DAJ, Bynoe ML (1965) Cultivation of a novel type of common cold virus in organ cultures. *Br Med J* 1:1467-1470
- Tyrrell DAJ, Almeida JD, Berry DM, Cunningham CH, Hamre D, Hofstad MS, Malluci L, McIntosh K (1968) Coronaviruses. *Nature* 220:650
- Tyrrell DAJ, Almeida JD, Cunningham CH, Dowdle WR, Hofstad MW, McIntosh K, Tajima M, Zastelskaya LY, Easterday BC, Kapikian A, Bingham RW (1975) Coronaviridae. *Intervirology* 5:76-82
- Tyrrell DAJ, Alexander JD, Almeida JD, Cunningham CH, Easterday BC, Garwes DJ, Hierholzer JC, Kapikian AZ, Macnaughton MR, McIntosh K (1978) Coronaviridae: second report. *Intervirology* 10:321-328
- Underdahl NR, Mebus EL, Stair EL, Rhodes MB, McGill LD, Twiehaus MJ (1974) Isolation of transmissible gastroenteritis virus from lungs of market-weight swine. *Am J Vet Res* 35:1209-1216
- Underdahl NR, Mebus CA, Torres-Medina A (1975) Recovery of transmissible gastroenteritis virus from chronically infected experimental pigs. *Am J Vet Res* 36:1473-1476
- Virelizier JL (1979) Effects of immunosuppressive agents on leucocyte interferon production in normal or thymus-deprived mice. *Transplantation* 27:353-355
- Virelizier JL (1981) Role of macrophages and interferon in natural resistance to mouse hepatitis virus infection. *Curr Top Microbiol Immunol* 92:51-64
- Virelizier JL, Allison AC (1976) Correlation of persistent mouse hepatitis virus (MHV-3) infection with its effect on mouse macrophage cultures. *Arch Virol* 50:279-285
- Virelizier JL, Gresser I (1978) Role of interferon in the pathogenesis of viral diseases of mice as demonstrated by the use of anti-interferon serum. V. Protective role in mouse hepatitis virus type 3 infection of susceptible and resistant strains of mice. *J Immunol* 120:1616-1619
- Virelizier JL, Dayan AD, Allison AC (1975) Neuropathological effects of persistent infection of mice by mouse hepatitis virus. *Infect Immun* 12:1127-1140
- Virelizier JL, Virelizier AM, Allison AC (1976) The role of circulating interferon in the modifications of immune responsiveness by mouse hepatitis virus (MHV-3). *J Immunol* 117:748-753
- Ward JM (1970) Morphogenesis of a virus in cats with experimental feline infectious peritonitis. *Virology* 41:191-194
- Ward JM, Gribble DH, Dungworth DL (1974) Feline infectious peritonitis: experimental evidence for its multiphasic nature. *Am J Vet Res* 35:1271-1275
- Ward JM, Collins MJ, Parker JC (1977) Naturally occurring mouse hepatitis virus infection in the nude mouse. *Lab Anim Sci* 27:372-376

- Watanabe H, Kobayashi K, Isayama Y (1975) Peculiar secretory IgA system identified in chickens. II. Identification and distribution of free secretory component and immunoglobulins of IgA, IgM and IgG in chicken external secretions. *J Immunol* 115:998-1001
- Watt RG (1978) Virological study of two commercial pig herds with respiratory disease. *Res Vet Sci* 24:147-153
- Wege H, Stephenson JR, Koga M, Wege Hanna, ter Meulen V (1981a) Genetic variation of neurotropic and non neurotropic murine coronaviruses. *J Gen Virol* 54:67-74
- Wege H, Koga M, Wege Hanna, ter Meulen V (1981b) JHM infections in rats as a model for acute and subacute demyelinating disease. In: ter Meulen V, Siddell S, Wege H (eds) *Biochemistry and biology of coronaviruses*. Plenum Press, New York, pp 327-349
- Weiner LP (1973) Pathogenesis of demyelination induced by a mouse hepatitis virus (JHM virus). *Arch Neurol* 28:293-303
- Weisbroth SH, Peress N (1977) Ophthalmic lesions and dacryoadenitis: a naturally occurring aspect of sialodacryoadenitis virus infection of the laboratory rat. *Lab Animal Sci* 27:466-473
- Weiser WY, Bang FB (1976) Macrophages genetically resistant to mouse hepatitis virus converted in vitro to susceptible macrophages. *J Exp Med* 143:690
- Weiser WY, Bang FB (1977) Blocking of in vitro and in vivo susceptibility to mouse hepatitis virus. *J Exp Med* 146:1467
- Weiser W, Vellisto I, Bang FB (1976) Congenic strains of mice susceptible and resistant to mouse hepatitis virus. *Proc Soc Exp Biol Med* 152:499
- Weiss RC, Dodds JW, Scott FW (1980) Disseminated intravascular coagulation in experimentally induced feline infectious peritonitis. *Am J Vet Res* 41:663-671
- Weiss SR, Leibowitz JL (1981) Comparison of the RNAs of murine and human coronaviruses. In: ter Meulen V, Siddell S, Wege H (eds) *Biochemistry and biology of coronaviruses*. Plenum Press, New York, pp 245-259
- Werdin RE, Sorensen DK, Stewart MS (1976) Porcine encephalomyelitis caused by hemagglutinating encephalomyelitis virus. *J Am Vet Med Assoc* 168:240-246
- Witte KH, Tuch K, Dubenkropp H, Walther C (1977) Untersuchungen über die Antigenverwandtschaft der Viren der feline infektiösen Peritonitis (FIP) und der transmissiblen Gastroenteritis (TGE) des Schweines. *Berl Münch Tierärztl Wochenschr* 80:396-401
- Wolfe LG, Griesemer RD (1966) Feline infectious peritonitis. *Pathol Vet* 3:255-270
- Woods RD, Cheville NF, Gallagher JE (1981) Lesions in the small intestine of newborn pigs inoculated with porcine, feline and canine coronaviruses. *Am J Vet Res* 42:1163-1169
- Yamada A, Taguchi F, Fujiwara K (1979) T-lymphocyte dependent difference in susceptibility between DDD and C<sub>3</sub>H mice to mouse hepatitis virus, MHV 3. *Jpn J Exp Med* 49:413-421
- Yaseen SA, Johnson-Lussenburg M (1981) Antigenic studies on coronavirus. I. Identification of the structural antigens of human coronavirus, strain 229E. *Can J Microbiol* 27:334-342