



ELSEVIER

# Implications of altered replication fidelity on the evolution and pathogenesis of coronaviruses

Everett C Smith<sup>1,3</sup> and Mark R Denison<sup>1,2,3</sup>

RNA virus evolution results from viral replication fidelity and mutational robustness in combination with selection. Recent studies have confirmed the impact of increased fidelity on RNA virus replication and pathogenesis; however, the impact of decreased fidelity is less defined. Coronaviruses have the largest RNA genomes, and encode an exoribonuclease activity that is required for high-fidelity replication. Genetically stable exoribonuclease mutants will allow direct testing of viral mutational tolerance to RNA mutagens and other selective pressures. Recent studies support the hypothesis that coronavirus replication fidelity may result from a multi-protein complex, suggesting multiple pathways to disrupt or alter virus fidelity and diversity, and attenuate pathogenesis.

## Addresses

<sup>1</sup> Department of Pediatrics, Vanderbilt University Medical Center, Nashville, TN 37232, USA

<sup>2</sup> Department of Microbiology and Immunology, Vanderbilt University Medical Center, Nashville, TN 37232, USA

<sup>3</sup> Elizabeth B. Lamb Center for Pediatric Research, Vanderbilt University Medical Center, Nashville, TN 37232, USA

Corresponding author: Denison, Mark R ([mark.denison@Vanderbilt.edu](mailto:mark.denison@Vanderbilt.edu))

Current Opinion in Virology 2012, 2:519–524

This review comes from a themed issue on **Virus evolution**

Edited by **Raul Andino** and **Marco Vignuzzi**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 1st August 2012

1879-6257/\$ – see front matter, Published by Elsevier B.V.

<http://dx.doi.org/10.1016/j.coviro.2012.07.005>

## Introduction

RNA viruses cause many existing and recently emerged human diseases, and contain diverse replication machinery capable of generating enormous numbers of viral progeny. Despite this diversity, the size of RNA virus genomes spans only an order of magnitude ( $10^3$  to  $10^4$ ), while DNA virus genomes vary in size over three orders of magnitude ( $10^4$  to  $10^7$ ) [1,2]. Such differences in genome size between RNA and DNA viruses have been theoretically coupled to the inherent low fidelity of the RNA-dependent RNA polymerases (RdRps) used in viral RNA synthesis ( $10^{-3}$  to  $10^{-4}$  mutations/nucleotide/round of replication; subsequently referred to as  $\mu$ ), which has long been proposed to result from the lack of proofreading within replicase complexes during viral RNA synthesis [3]. While this concept of constitutively low replication fidelity has been useful for predicting RNA virus evol-

ution, viral systems have recently begun to be developed to directly test the impact of altered replication fidelity on viral evolution, genome size, viral replication, and/or viral pathogenesis. Studies using poliovirus and chikungunya virus have demonstrated that even modest increases in RNA replication fidelity (2–4-fold) result in decreased viral fitness *in vitro* and pathogenicity *in vivo* [4,5,6<sup>\*</sup>]. Similarly, only very small decreases in replication fidelity (<4-fold) were tolerated during poliovirus replication [5,7]. Thus, work with these viruses demonstrates rather stringent limits on the variation in fidelity tolerated during viral replication, suggesting a finely tuned balance between genome stability and the diversity required for survival. Mutational robustness, or the capacity of a virus population to tolerate mutations, has also been shown to influence sensitivity to environmental conditions or mutagens. Coxsackie virus demonstrates increased sensitivity to RNA mutagens compared to poliovirus in a population size-dependent manner, suggesting that population tolerance for mutational diversity is likely a distinct feature of virus families [8<sup>\*</sup>].

These exciting developments, in combination with advances in next generation sequencing, will allow direct experimental testing of the roles of fidelity and diversity in RNA virus genome size and complexity, host range expansion and adaptation, tropism, sensitivity to RNA mutagens or environmental stressor such as temperature, virulence or attenuation, and antigenicity and immune escape. The impact of decreased replication fidelity on attenuation or pathogenesis *in vivo* is less well studied, however, recent studies with the coronaviruses (CoVs) suggest that they may be excellent models to examine these important questions.

## Coronavirus genome size: is bigger smarter?

Bioinformatic models suggest that RNA genome size is constrained to an upper threshold of approximately 30 kb for an RNA based organism; beyond which, both the stability and the faithful replication of the viral genome cannot be maintained [1,9]. Coronaviruses fail to abide by these theoretical rules, and encode the largest known RNA genomes (27–32 kb) at almost twice the length of the next-largest non-segmented RNA viruses [1]. Assuming a fixed genomic mutation rate, coronaviruses and other large nidoviruses (e.g. roniviruses, toroviruses) must possess mechanisms to limit the accumulation of deleterious mutations while concomitantly maintaining the genetic diversity required for adaptation. Without such mechanisms, larger RNA viruses would accumulate


excessive numbers of deleterious mutations leading to a dramatic loss of fitness [10]. The demonstrated capacity of coronaviruses for host switching, [11] and the emergence of SARS-CoV into the human population [12,13] demonstrate that coronaviruses have strong adaptive capacity all while encoding such large genomes. Finally, the lack of larger or more complex RNA genomes exceeding those of the nidoviruses suggests that they may represent the upper limit of replicating RNA molecules, continually negotiating genome size and/or complexity with replicative stability. How then do the coronaviruses tolerate such a large and complex genome, and how did it arise?

### Coronaviruses encode multiple RNA modifying enzymes, including a 3'-to-5' exoribonuclease

Coronaviruses encode the most complex array of viral replicase proteins of any positive-strand RNA virus family [14]. The coronavirus replicase polyproteins contain up to


16 nonstructural protein domains (nsp1–16), many of which have known or predicted functions in viral RNA synthesis or modification (Figure 1) [9,14,15], including: RNA primase [16], RNA-dependent RNA polymerase (RdRp) [17], helicase/ATPase [18,19], N-methyltransferase [20], endoribonuclease [21], and 2'-O-methyltransferase [22] activities. Additionally, the coronaviruses encode a 3'-to-5' exoribonuclease (ExoN) domain within nsp14 [23]. ExoN is encoded in all larger members of the *Nidovirales* order (i.e. *Coronaviridae* and *Roniviridae*), but is absent within the smaller *Arteriviridae* family members ('small nidoviruses') (Figure 2) [9]. Furthermore, ExoN activity has not been predicted or demonstrated to be present within any RNA viruses containing genomes smaller than the arteriviruses. Coronavirus nsp14-ExoN is a predicted member of the 'DEDD' superfamily that includes both RNA and DNA exonucleases from a diverse group of eukaryotic and prokaryotic organisms [9,24], and which derives its name from the four invariant acidic amino acids distributed across three

Figure 1



Coronavirus genome organization and identified viral nsp interactions. (a) Coronaviruses express 16 nonstructural proteins (nsp1–16) from open reading frames ORF1a and ORF1b, along with several structural and accessory proteins. Several nsp have been shown to have roles in RNA synthesis or modification: nsp12 (RdRp, green), nsp13 [helicase (Hel)], nsp14 [3'-to-5' exoribonuclease (ExoN) and N7-methyltransferase (N7-MT); blue], nsp15 [endoribonuclease (NEndoU)], and nsp16 [2'-O-methyltransferase (2'-O-MT), red]. (b) ExoN shares conserved motifs (I, II and III) with other identified 3'-to-5' exonucleases, and requires the amino acids D-E-D-D (white boxes) for activity. Nsp14 also contains a zinc finger (Zn<sup>2+</sup> F, grey box) and a functional N7-methyltransferase domain. (c) Multiple nsp have been demonstrated to form higher-order complexes, and/or functionally regulate one another *in vitro* and in recombinant viruses.

Figure 2



Relationship between ssRNA virus genome size and the presence or absence of ExoN. The sizes of full viral genomes identified using the NCBI Viral Genome Resource [48] (with the exception of Yellow head virus strain YHV1999, GenBank ID: FJ848675.1) are plotted, and the average genome size per virus family or subfamily is shown as a horizontal black bar. Members of the order *Nidovirales* are denoted with an N, while the proposed new *Nidovirales* candidate family, *Mesoniviridae* [31], is denoted with an asterisk.

conserved sequence motifs: motifs I (DE), II (D) and III (D). Nsp14-ExoN is distinguished from other cellular ExoN homologs by the presence of a highly conserved putative zinc-finger domain positioned between motifs I and II [9] that could potentially confer specificity for RNA over DNA, and thus be important for proper targeting and function. What then is the known or predicted relationship of nsp14-ExoN to CoV genome size, stability and replication fidelity?

### ExoN and the expansion of the RNA genome

The original prediction of the CoV ExoN domain within nsp14 suggested that it might serve a role as a possible proofreading exonuclease, an activity without precedent in RNA viruses [9]. Though proofreading and repair mechanisms were long thought to occur only during DNA replication, it is now clear that both DNA-dependent RNA polymerases (DdRps) and RNA-dependent RNA polymerases (RdRps) are capable of proofreading (reviewed in [25]). Cellular DdRps, such as human RNA pol II, have been shown to excise misincorporated nucleotides via 3'-to-5' exonuclease activity that is stimulated upon binding of specific cleavage-stimulatory factors [26,27]. While nuclease activity has been reported for the influenza virus RdRp [28], evidence supporting either an intrinsic 3'-to-5' exonuclease activity of viral RdRps, or the occurrence of proofreading-repair mechanisms during

replication of other RNA viruses has yet to be found [3]. The identification of the CoV ExoN within the large nidoviruses suggested that acquisition of ExoN allowed for genome expansion to double that of the arteriviruses [1,9]. Closteroviruses (RNA viruses outside of the order *Nidovirales*) have non-segmented genomes larger than the arteriviruses, but smaller than the coronaviruses, and do not encode any known exonucleases [1], leaving a theoretical genome size gap of >10 kb between genomes lacking and encoding ExoN (Figure 2). However, important support for the ExoN hypothesis has been provided by the recent discovery of two invertebrate nidoviruses (Nam Dinh virus, NDiV; and Cavally virus, CAVV) containing genomes of approximately 20 kb that encode an nsp14-ExoN homolog [29,30,31], further narrowing the boundaries of the genome size gap to <4 kb for genomes lacking and containing ExoN. Is ExoN-mediated genome expansion due to alterations in RNA genome replication fidelity?

### ExoN is required for replication fidelity and is a probable RNA-dependent RNA proofreading enzyme

The predicted enzymatic activity [9] of nsp14-ExoN was confirmed *in vitro* for bacterially expressed SARS-CoV nsp14, demonstrating that expression of nsp14 alone was sufficient for 3'-to-5' exoribonuclease activity [23]. Recombinant mutant MHV-A59 and SARS-CoV viruses containing motif 1 (DE to AA) substitutions (named S-ExoN and M-ExoN), which were shown to significantly diminished or abolished ExoN activity [23], are viable and have less than 1 log reduction in peak titers [32,33] compared to WT. Single cycle replication of M-ExoN and S-ExoN resulted in 15–20-fold increases in mutation accumulation and similar calculated increased mutation rates ( $\mu$ ) as compared to their respective WT counterparts. The estimated mutation rate of the ExoN mutants ( $\mu \sim 10^{-4}$  to  $10^{-5}$ ) more closely aligns with RNA viruses encoding smaller genomes, while the WT viruses containing intact ExoN activity appear to have a profound increase in replication fidelity ( $\mu \sim 10^{-6}$  to  $10^{-7}$ ). Both the *in vitro* exoribonuclease activity and the demonstrated requirement for ExoN for high fidelity replication are consistent with the hypothesis that nsp14-ExoN is involved in RNA-dependent RNA proofreading. Given that ExoN activity is distinct from RdRp activity, could ExoN be a component of a larger multi-protein polymerase proofreading complex?

### ExoN is likely a proofreading component of a larger multi-subunit error recognition and repair complex

Identification of ExoN activity distinct from the viral RdRp [23] suggests that nsp14-ExoN is a component of a larger multi-protein complex that includes nsp8, 10, 12 and 16 and possibly others. As other DEDD superfamily exonucleases are subunits of larger proteins

with polymerase activity, and serve to recognize and repair mismatched nucleotides, ExoN would be predicted to associate with other viral and/or cellular proteins. Several lines of evidence support this hypothesis (Figure 1). Nsp14 also contains N7-methyltransferase activity (N7-MTase) [20,34], demonstrating a requirement for ExoN to function in that context, and potentially interact with other virus-encoded proteins in the capping pathway, specifically nsp16, a 2'-O-methyltransferase (2'-O-MTase) [22]. While, nsp14 possesses independent ExoN activity, very recent work demonstrates that ExoN activity *in vitro* is enhanced up to 35-fold by the binding of nsp10 [35\*\*], a protein with no known independent enzymatic activity. Nsp10 is also required for activation of SARS-CoV nsp16 2'-O-MTase activity [34], and surprisingly, nsp10–16 and nsp10–14 interactions appear to bind at overlapping sites on the surface of nsp10 [35\*\*,36]. Given the distinct functions of nsp10–16 and nsp10–14 binding events [35\*\*], nsp14 could possibly form at least two higher-order complexes during viral replication, one involved in putative viral RNA proofreading and composed of nsps10, 12 and 14 (at minimum), and a second viral RNA capping complex composed of nsps10, 14 and 16. While there is currently no experimental evidence for cooperation of nsp14 with the nsp15 endonuclease, such an interaction has been proposed for viral RNA modification [9]. Finally, nsp8 (primase) interacts with nsp7 and is required for nsp12 RdRp activity *in vitro* [37,38], further supporting the possibility of a multi-protein complex. Such multi-protein complexes would allow for testing if multiple proteins regulate virus replication fidelity, and if natural or induced variability in fidelity exists under differing replication conditions. If so, could the modular nature of the coronavirus replicase represent a genetically encoded system for fidelity regulation?

### Is replication fidelity of RNA viruses fixed or responsive to selective pressure?

Mutation rates of RNA viruses have been the subject of extensive research [39\*\*], with the general paradigm being that the inherent low fidelity of viral RdRps is a major contributor to mutation rates. While such biochemical restraints on RdRp function help in understanding replication fidelity determinants, the existence of high fidelity RdRp variants [6,40,41], as well as the high fidelity replication of coronaviruses, demonstrates that increased RdRp fidelity is both feasible and attainable for RNA viruses. Consistent with the need to balance adaptability with genome maintenance, viral replication in the presence of RNA mutagens demonstrates that RNA viruses can only accommodate limited changes in mutation rates without incurring significant fitness costs [8\*,42,43]. Such sensitivity not only suggests that RNA viruses replicate close to a maximum error threshold [44], but also suggests that high mutation rates are a product of selection. Additionally, selection for further reductions or

enhancements in replication fidelity has been described for RNA viruses [6\*,40,41,45–47], suggesting that replication fidelity may represent an evolutionarily defined range rather than a fixed value, to maintain optimal population fitness. Given the likely modular nature of a coronavirus proofreading complex and the tolerance for fidelity variation (up to 20-fold at least), they may represent a unique model by which to measure viral mutation rates under variable selective pressures.

### Conclusions

The multiple viral systems becoming available for study of increased and decreased fidelity create important new platforms and research applications, particularly the opportunity to study the impact of altered fidelity as a universal approach for attenuation of entire taxonomic groups of RNA viruses. Increased replication fidelity has already been shown to be attenuating for several RNA viruses. In contrast, the concept of decreased fidelity as an attenuation strategy for live viruses has been subject to concerns, when in fact there is no experimental evidence that decreasing virus replication fidelity accelerates emergence of virulence, or expansion of host range. If fidelity is a long-term evolved phenotype, then fidelity alterations could tip the balance toward continual emergence of attenuating mutations that would trump the emergence of virulence while possibly maintaining and expanding the repertoire of immune response. Stable coronavirus mutants that tolerate profoundly decreased replication fidelity may represent an excellent model to study the implications of decreased replication fidelity on attenuation and vaccine design in human viruses with strong animal models.

### References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
  - of outstanding interest
1. Gorbalenya AE, Enjuanes L, Ziebuhr J, Snijder EJ: **Nidovirales: evolving the largest RNA virus genome.** *Virus Research* 2006, **117**:17-37.
  2. Holmes EC: **Evolution in health and medicine Sackler colloquium: the comparative genomics of viral emergence.** *Proceedings of the National Academy of Sciences of the United States of America* 2010, **107**(Suppl. 1):1742-1746.
  3. Steinhauer DA, Domingo E, Holland JJ: **Lack of evidence for proofreading mechanisms associated with an RNA virus polymerase.** *Gene* 1992, **122**:281-288.
  4. Vignuzzi M, Stone JK, Arnold JJ, Cameron CE, Andino R: **Quasispecies diversity determines pathogenesis through cooperative interactions in a viral population.** *Nature* 2006, **439**:344-348.
  5. Pfeiffer JK, Kirkegaard K: **Increased fidelity reduces poliovirus fitness and virulence under selective pressure in mice.** *PLoS Pathogenesis* 2005, **1**:e11.
  6. Coffey LL, Beeharry Y, Borderia AV, Blanc H, Vignuzzi M: **Arbovirus high fidelity variant loses fitness in mosquitoes and mice.** *Proceedings of the National Academy of Sciences of the United States of America* 2011, **108**:16038-16043.



This work describes the first high-fidelity arbovirus variant, which was due to a single point mutation (C483Y) within the viral RdRp. Infection experiments in mosquitoes revealed that increased replication fidelity decreased infection and viral dissemination rates. Additional infection experiments in newborn mice demonstrated truncated viremia and lower organ titers, providing evidence that reduced genetic diversity during arbovirus infections negatively impacts viral fitness.

7. Crotty S, Cameron CE, Andino R: **RNA virus error catastrophe: direct molecular test by using ribavirin.** *Proceedings of the National Academy of Sciences of the United States of America* 2001, **98**:6895-6900.
8. Graci JD, Gnading NF, Galarraga JE, Castro C, Vignuzzi M, Cameron CE: **Mutational robustness of an RNA virus influences sensitivity to lethal mutagenesis.** *Journal of Virology* 2012, **86**:2869-2873.  
Experiments using the known RNA mutagen ribavirin demonstrated that coxsackievirus B3 (CVB3) has a lower mutational robustness relative to poliovirus (PV). Such sensitivity of CVB3 to ribavirin was dependent upon population size, providing strong evidence that RNA viruses abide by the tenants of lethal mutagenesis theory. This work has powerful implications for the testing and development of potential mutagenic antivirals.
9. Snijder EJ, Bredenbeek PJ, Dobbe JC, Thiel V, Ziebuhr J, Poon LL, Guan Y, Rozanov M, Spaan WJ, Gorbalenya AE: **Unique and conserved features of genome and proteome of SARS-coronavirus, an early split-off from the coronavirus group 2 lineage.** *Journal of Molecular Biology* 2003, **331**:991-1004.
10. Holmes EC: **Error thresholds and the constraints to RNA virus evolution.** *Trends in Microbiology* 2003, **11**:543-546.
11. Baric RS, Yount B, Hensley L, Peel SA, Chen W: **Episodic evolution mediates interspecies transfer of a murine coronavirus.** *Journal of Virology* 1997, **71**:1946-1955.
12. Lau SK, Woo PC, Li KS, Huang Y, Tsoi HW, Wong BH, Wong SS, Leung SY, Chan KH, Yuen KY: **Severe acute respiratory syndrome coronavirus-like virus in Chinese horseshoe bats.** *Proceedings of the National Academy of Sciences of the United States of America* 2005, **102**:14040-14045.
13. Li W, Shi Z, Yu M, Ren W, Smith C, Epstein JH, Wang H, Crameri G, Hu Z, Zhang H *et al.*: **Bats are natural reservoirs of SARS-like coronaviruses.** *Science* 2005, **310**:676-679.
14. Perlman S, Netland J: **Coronaviruses post-SARS: update on replication and pathogenesis.** *Nature Reviews Microbiology* 2009, **7**:439-450.
15. Ziebuhr J, Snijder EJ, Gorbalenya AE: **Virus-encoded proteinases and proteolytic processing in the Nidovirales.** *Journal of General Virology* 2000, **81**:853-879.
16. Imbert I, Guillemot JC, Bourhis JM, Bussetta C, Coutard B, Egloff MP, Ferron F, Gorbalenya AE, Canard B: **A second, non-canonical RNA-dependent RNA polymerase in SARS coronavirus.** *EMBO Journal* 2006, **25**:4933-4942.
17. te Velthuis AJW, Arnold JJ, Cameron CE, van den Worm SHE, Snijder EJ: **The RNA polymerase activity of SARS-coronavirus nsp12 is primer dependent.** *Nucleic Acids Research* 2011, **39**:9458.
18. Seybert A, Hegyi A, Siddell SG, Ziebuhr J: **The human coronavirus 229E superfamily 1 helicase has RNA and DNA duplex-unwinding activities with 5'-to-3' polarity.** *RNA* 2000, **6**:1056-1068.
19. Ivanov KA, Ziebuhr J: **Human coronavirus 229E nonstructural protein 13: characterization of duplex-unwinding, nucleoside triphosphatase, and RNA 5'-triphosphatase activities.** *Journal of Virology* 2004, **78**:7833-7838.
20. Chen Y, Cai H, Pan J, Xiang N, Tien P, Ahola T, Guo DY: **Functional screen reveals SARS coronavirus nonstructural protein nsp14 as a novel cap N7 methyltransferase.** *Proceedings of the National Academy of Sciences of the United States of America* 2009, **106**:3484-3489.
21. Ivanov KA, Hertzog T, Rozanov M, Bayer S, Thiel V, Gorbalenya AE, Ziebuhr J: **Major genetic marker of nidoviruses encodes a replicative endoribonuclease.** *Proceedings of the National Academy of Sciences of the United States of America* 2004, **101**:12694-12699.
22. Decroly E, Imbert I, Coutard B, Bouvet M, Selisko B, Alvarez K, Gorbalenya AE, Snijder EJ, Canard B: **Coronavirus nonstructural protein 16 is a cap-0 binding enzyme possessing (nucleoside-2'-O)-methyltransferase activity.** *Journal of Virology* 2008, **82**:8071-8084.
23. Minskaia E, Hertzog T, Gorbalenya AE, Campanacci V, Cambillau C, Canard B, Ziebuhr J: **Discovery of an RNA virus 3'-5' exoribonuclease that is critically involved in coronavirus RNA synthesis.** *Proceedings of the National Academy of Sciences of the United States of America* 2006, **103**:5108-5113.
24. Zuo Y, Deutscher MP: **Exoribonuclease superfamilies: structural analysis and phylogenetic distribution.** *Nucleic Acids Research* 2001, **29**:1017-1026.
25. Sydow JF, Cramer P: **RNA polymerase fidelity and transcriptional proofreading.** *Current Opinion in Structural Biology* 2009, **19**:732-739.
26. Poole AM, Logan DT: **Modern mRNA proofreading and repair: clues that the last universal common ancestor possessed an RNA genome?** *Molecular Biology and Evolution* 2005, **22**:1444-1455.
27. Thomas MJ, Platas AA, Hawley DK: **Transcriptional fidelity and proofreading by RNA polymerase II.** *Cell* 1998, **93**:627-637.
28. Ishihama A, Mizumoto K, Kawakami K, Kato A, Honda A: **Proofreading function associated with the RNA-dependent RNA polymerase from influenza virus.** *Journal of Biological Chemistry* 1986, **261**:10417-10421.
29. Nga PT, Parquet Mdel C, Lauber C, Parida M, Nabeshima T, Yu F, Thuy NT, Inoue S, Ito T, Okamoto K *et al.*: **Discovery of the first insect nidovirus, a missing evolutionary link in the emergence of the largest RNA virus genomes.** *PLoS Pathogenesis* 2011, **7**:e1002215.  
The emergence of virally encode exonuclease activity among RNA viruses is thought to have driven RNA virus genome expansion. This work describes the discovery of a new insect nidovirus, Nam Dinh virus (NiDV), which helps to evolutionary link the 'small' and 'large' nidoviruses. NiDV was shown to encode a homolog of nsp14-ExoN, making it the 'smallest' nidovirus (~20 kb) encoding ExoN activity.
30. Zirkel F, Kurth A, Quan PL, Briesse T, Ellerbrok H, Pauli G, Leendertz FH, Lipkin WI, Ziebuhr J, Drosten C *et al.*: **An insect nidovirus emerging from a primary tropical rainforest.** *mBio* 2011, **2**:e00077-00011.
31. Lauber C, Ziebuhr J, Junglen S, Drosten C, Zirkel F, Nga PT, Morita K, Snijder EJ, Gorbalenya AE: **Mesoniviridae: a proposed new family in the order Nidovirales formed by a single species of mosquito-borne viruses.** *Archives of Virology* 2012 <http://dx.doi.org/10.1007/s00705-012-1295-x>:1-6.
32. Eckerle LD, Becker MM, Halpin RA, Li K, Venter E, Lu X, Scherbakova S, Graham RL, Baric RS, Stockwell TB *et al.*: **Infectivity of SARS-CoV Nsp14-exonuclease mutant virus replication is revealed by complete genome sequencing.** *PLoS Pathogenesis* 2010, **6**:e1000896.  
Using full genome deep sequencing, the authors showed that recombinant SARS-CoV virus lacking ExoN exhibits a profound decrease in replication fidelity. The mutant viruses had stable growth defects and demonstrated a 21-fold increase in mutation frequency during replication in cell culture. This work provides strong experimental evidence for putative proofreading during RNA virus replication.
33. Eckerle LD, Lu X, Sperry SM, Choi L, Denison MR: **High fidelity of murine hepatitis virus replication is decreased in nsp14 exoribonuclease mutants.** *Journal of Virology* 2007, **81**:12135-12144.
34. Bouvet M, Debarnot C, Imbert I, Selisko B, Snijder EJ, Canard B, Decroly E: **In vitro reconstitution of SARS-coronavirus mRNA cap methylation.** *PLoS Pathogenesis* 2010, **6**:e1000863.
35. Bouvet M, Imbert I, Subissi L, Gluais L, Canard B, Decroly E: **RNA 3'-end mismatch excision by the severe acute respiratory syndrome coronavirus nonstructural protein nsp10/nsp14 exoribonuclease complex.** *Proceedings of the National Academy of Sciences of the United States of America* 2012, **109**:9372-9377.  
Nsp14-ExoN has previously been shown to have 3'-to-5' exoribonuclease activity, and this work provides evidence that nsp14-ExoN functions in complex with other viral nsps. Using recombinant protein, the authors

demonstrated that incubation of nsp14 with nsp10 greatly enhances nsp14-ExoN activity, and that this complex was capable of hydrolyzing single mismatched nucleotides at the 3' end of synthetic RNA molecules.

36. Decroly E, Debarnot C, Ferron F, Bouvet M, Coutard B, Imbert I, Gluais L, Papageorgiou N, Sharff A, Bricogne G *et al.*: **Crystal structure and functional analysis of the SARS-coronavirus RNA cap 2'-O-methyltransferase nsp10/nsp16 complex.** *PLoS Pathogenesis* 2011, **7**:e1002059.
37. te Velthuis AJ, van den Worm SH, Snijder EJ: **The SARS-coronavirus nsp7 + nsp8 complex is a unique multimeric RNA polymerase capable of both de novo initiation and primer extension.** *Nucleic Acids Research* 2012, **40**:1737-1747.
38. Xiao Y, Ma Q, Restle T, Shang W, Svergun DI, Ponnusamy R, Sczakiel G, Hilgenfeld R: **Nonstructural proteins 7 and 8 of feline coronavirus form a 2:1 heterotrimer that exhibits primer-independent RNA polymerase activity.** *Journal of Virology* 2012, **86**:4444-4454.
39. Sanjuan R, Nebot MR, Chirico N, Mansky LM, Belshaw R: **Viral mutation rates.** *Journal of Virology* 2010, **84**:9733-9748.  
Viral mutation rates have been the subject of intense study, and this work provides the most current and extensive analysis of viral mutation rates across RNA, DNA and retrovirus genomes. Additionally, the authors thoroughly discuss the inherent experimental problems associated with measuring mutation rates, including both the sources of, and how to account for, selection bias.
40. Pfeiffer JK, Kirkegaard K: **A single mutation in poliovirus RNA-dependent RNA polymerase confers resistance to mutagenic nucleotide analogs via increased fidelity.** *Proceedings of the National Academy of Sciences of the United States of America* 2003, **100**:7289-7294.
41. Levi LI, Gnadig NF, Beaucourt S, McPherson MJ, Baron B, Arnold JJ, Vignuzzi M: **Fidelity variants of RNA dependent RNA polymerases uncover an indirect, mutagenic activity of amiloride compounds.** *PLoS Pathogenesis* 2010, **6**:e1001163.
42. Anderson JP, Daifuku R, Loeb LA: **Viral error catastrophe by mutagenic nucleosides.** *Annual Review of Microbiology* 2004, **58**:183-205.
43. Dapp MJ, Clouser CL, Patterson S, Mansky LM: **5-Azacytidine can induce lethal mutagenesis in human immunodeficiency virus type 1.** *Journal of Virology* 2009, **83**:11950-11958.
44. Manrubia SC, Domingo E, Lazaro E: **Pathways to extinction: beyond the error threshold.** *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences* 2010, **365**:1943-1952.
45. Arias A, Arnold JJ, Sierra M, Smidansky ED, Domingo E, Cameron CE: **Determinants of RNA-dependent RNA polymerase (in)fidelity revealed by kinetic analysis of the polymerase encoded by a foot-and-mouth disease virus mutant with reduced sensitivity to ribavirin.** *Journal of Virology* 2008, **82**:12346-12355.
46. Sierra M, Airaksinen A, Gonzalez-Lopez C, Agudo R, Arias A, Domingo E: **Foot-and-mouth disease virus mutant with decreased sensitivity to ribavirin: implications for error catastrophe.** *Journal of Virology* 2007, **81**:2012-2024.
47. Mansky LM, Bernard LC: **3'-Azido-3'-deoxythymidine (AZT) and AZT-resistant reverse transcriptase can increase the in vivo mutation rate of human immunodeficiency virus type 1.** *Journal of Virology* 2000, **74**:9532-9539.
48. Bao Y, Federhen S, Leipe D, Pham V, Resenchuk S, Rozanov M, Tatusov R, Tatusova T: **National center for biotechnology information viral genomes project.** *Journal of Virology* 2004, **78**:7291-7298.
49. Donaldson EF, Graham RL, Sims AC, Denison MR, Baric RS: **Analysis of murine hepatitis virus strain A59 temperature-sensitive mutant TS-LA6 suggests that nsp10 plays a critical role in polyprotein processing.** *Journal of Virology* 2007, **81**:7086-7098.