

# Roadmap to developing a recombinant coronavirus S protein receptor-binding domain vaccine for severe acute respiratory syndrome

*Expert Rev. Vaccines* 11(12), 1405–1413 (2012)

Shibo Jiang<sup>1,2</sup>,  
Maria Elena  
Bottazzi<sup>3,4,5,6,7,8</sup>,  
Lanying Du<sup>1</sup>,  
Sara Lustigman<sup>1</sup>,  
Chien-Te Kent Tseng<sup>9</sup>,  
Elena Curti<sup>3,4,5,6,7,8</sup>,  
Kathryn Jones<sup>3,4,5,6,7,8</sup>,  
Bin Zhan<sup>3,4,5,6,7,8</sup> and  
Peter J Hotez<sup>\*3,4,5,6,7,8</sup>

<sup>1</sup>Lindsley F Kimball Research Institute,  
New York Blood Center, New York,  
NY, USA

<sup>2</sup>Key Laboratory of Medical Molecular  
Virology of Ministries of Education  
and Health, Shanghai Medical College  
and Institute of Medical Microbiology,  
Fudan University, Shanghai, China

<sup>3</sup>Sabin Vaccine Institute, Washington,  
DC, USA

<sup>4</sup>Texas Children's Hospital Center for  
Vaccine Development, Houston, TX,  
USA

<sup>5</sup>Section of Pediatric Tropical Medicine,  
Baylor College of Medicine, Houston,  
TX, USA

<sup>6</sup>Departments of Pediatrics, Baylor  
College of Medicine, Houston, Texas,  
USA

<sup>7</sup>Molecular Virology & Microbiology,  
Baylor College of Medicine, Houston,  
TX, USA

<sup>8</sup>National School of Tropical Medicine,  
Baylor College of Medicine, Houston,  
TX, USA

<sup>9</sup>Department of Microbiology &  
Immunology, University of Texas  
Medical Branch, Galveston, TX, USA

\*Author for correspondence:

Tel.: +1 713 798 1199

[hotez@bcm.edu](mailto:hotez@bcm.edu)

<sup>†</sup>Authors contributed equally to the  
manuscript.

A subunit vaccine, RBD-S, is under development to prevent severe acute respiratory syndrome (SARS) caused by SARS coronavirus (SARS-CoV), which is classified by the US NIH as a category C pathogen. This vaccine is comprised of a recombinant receptor-binding domain (RBD) of the SARS-CoV spike (S) protein and formulated on alum, together with a synthetic glucopyranosyl lipid A. The vaccine would induce neutralizing antibodies without causing Th2-type immunopathology. Vaccine development is being led by the nonprofit product development partnership; Sabin Vaccine Institute and Texas Children's Hospital Center for Vaccine Development in collaboration with two academic partners (the New York Blood Center and University of Texas Medical Branch); an industrial partner (Immune Design Corporation); and Walter Reed Army Institute of Research. A roadmap for the product development of the RBD-S SARS vaccine is outlined with a goal to manufacture the vaccine for clinical testing within the next 5 years.

**KEYWORDS:** receptor-binding domain • recombinant vaccine • SARS • severe acute respiratory syndrome • spike (S) protein

Severe acute respiratory syndrome (SARS) emerged as a new human infection in South China in 2002, and ultimately caused over 8000 infections and almost 800 deaths as it spread to five continents along airline routes [1,2]. To date, most SARS patients have been adults with very few cases in children less than 15 years of age [2]. While the overall mortality rate from SARS during the 2002–2003 pandemic was approximately 10%, the mortality among the elderly exceeded 50% [1,2]. Subsequent outbreaks were reported later in 2003 with the latest outbreak in 2004 in China occurring as a result of accidental exposure to a laboratory virus isolate [1]. Early on, a new coronavirus, SARS coronavirus (SARS-CoV), was identified as the etiologic agent responsible for human SARS; the virus was also determined to be of zoonotic origin with bats, Chinese palm civets and other mammals serving as natural reservoir hosts [2]. SARS-CoV is an enveloped virus with a positive and ssRNA genome encoding a replicase and several

structural proteins including an important spike (S) protein, which binds to host receptors and can elicit neutralizing antibodies linked to protective immunity [1]. SARS outbreaks remain as a serious global concern because of the continued threat of zoonotic transmission to humans or the accidental or intentional release of laboratory isolates [1]. The SARS-CoV is classified by the National Institute of Allergy and Infectious Diseases of the US NIH (NIH-NIAID) as a category C pathogen, which also includes yellow fever virus, influenza virus and other emerging pathogens that could be bioengineered for mass dissemination based on their availability, ease of production and dissemination, and potential for high morbidity and mortality [101].

The prospect of renewed transmission of SARS-CoV to human populations suggests an urgent need to prepare vaccines for this emerging agent [2]. Vaccines to protect high-risk groups, including the elderly, healthcare workers and laboratory personnel are particularly critical [2].

Several recent reviews have been published on the development of SARS vaccines, including those targeting the SARS-CoV S protein [1–3]. Here, the authors provide a rationale for developing a vaccine based on eliciting neutralizing antibodies to the receptor-binding domain (RBD) of the S protein (RBD-S) [4] and building on this concept the authors provide a roadmap for the development and manufacture of a prototype RBD-S SARS vaccine. Our roadmap includes using the peer-reviewed literature to invite the scientific community to follow the progress of the development of a new vaccine through the activities of a nonprofit product development partnership (PDP).

### Proposed product

The proposed RBD-S SARS vaccine is a recombinant protein-based vaccine formulated with alum (either Alhydrogel® or aluminum phosphate), together with a suitable synthetic Toll-like receptor 4 (TLR4) agonist known as glucopyranosyl lipid A (GLA).

### Antigen

The recombinant protein is comprised of a 193 amino acid (21 kDa) polypeptide corresponding to the minimal RBD (amino acid residues: 318–510) of the SARS-CoV S protein [4]. The full-length SARS-CoV S protein belongs to the class I of viral fusion proteins, which also includes the glycoprotein gp160 of HIV and hemagglutinin of influenza virus [5,6]. The S protein fragment spanning the 193-mer polypeptide is the minimal RBD responsible for virus binding to the host-cell receptor, angiotensin-converting enzyme 2 [4,7].

RBD-S, which contains multiple conformation-dependent epitopes, induces strong neutralizing antibody responses, CD8<sup>+</sup> T-cell responses and long-term protective immunity in animal models, as well as potent cross-neutralizing antibodies against pseudoviruses expressing S proteins of 2002–2003 and 2003–2004 human SARS-CoV strains, and an isolate from a civet zoonotic host [8–11]. Moreover, more than 50% of the neutralizing activity in the sera of convalescent SARS patients is mediated by RBD-S-specific antibodies [9,12], while RBD-S can absorb and remove the majority of neutralizing antibodies in the antisera of rabbits, mice and monkeys immunized with inactivated SARS-CoV or modified vaccinia virus expressing S protein [9,13]. These studies provide the rationale for selecting RBD-S as an important target for vaccine development. The recombinant 21 kDa RBD-S will be expressed in yeast (*Pichia pastoris*), bacteria (*Escherichia coli*), or baculovirus (pending studies demonstrating optimal protection, as described below). More recently, a second RBD-S 219-mer construct corresponding to residues 318–536 was expressed in Chinese hamster ovary-K1 cells and shown to exhibit correct folding and conformation followed by equivalent or enhanced protection [14]; this construct will also be evaluated as a back-up vaccination strategy.

### Adjuvant

GLA is a 1.7 kDa synthetic TLR4 agonist functionally equivalent to monophosphoryl lipid A, a nontoxic derivative of the lipopolysaccharide [15–18]. In a study in mice with a commercial killed influenza vaccine, chemically synthesized GLA was effective in

enhancing IgG1 and IgG2a titers, both as an oil–water emulsion (squalene oil and surfactant) and as an aqueous formulation [15,16]. Subsequent studies revealed that GLA also enhances T-cell responses to influenza vaccine among older adults [18], which would be a desirable feature for a SARS vaccine.

### Indication & target product profile of the RBD-S SARS vaccine

The vaccine is initially indicated for adults and individuals over the age of 15 who are considered at greatest risk of mortality from SARS [2]. The vaccine would be administered as an injectable product to protect against lethal SARS-CoV infection, either as a single dose or two doses spaced closely together (within 2–4 weeks), so that vulnerable populations could be rapidly immunized in an outbreak setting (TABLE 1). The proposed target product profile also accounts for the stability of the RBD-S SARS vaccine (up to 4 years at 2–8°C) to allow for emergency stockpiling and to protect at low-cost at-risk populations in underdeveloped and developing countries (particularly southeast Asia where the risk of reemergence is greatest), as well as developed countries.

### Comparative advantage of an RBD-S SARS vaccine

Several different technologies have been proposed for developing human SARS-CoV vaccines, including inactivated virus vaccines, live–attenuated vaccines, adenovirus or poxvirus vectored vaccines, and DNA vaccines [1]. Among the concerns regarding these vaccines is the possibility of vaccine enhancement of disease, which has been noted with feline CoVs causing feline infectious peritonitis [1,19,20]. Similarly, laboratory animals administered alum-adjuvanted whole-virus SARS-CoV vaccine and subsequently challenged with infectious virus experienced enhanced disease and immunopathology comprised of eosinophilic infiltration and Th2-mediated inflammatory alveolar damage [21], similar to immunopathologic lung reactions found in infants and animals challenged with respiratory syncytial virus (RSV) [22,23]. A double-inactivated SARS-CoV (formalin and ultraviolet radiation) was also shown to elicit eosinophilic and immunoenhancing pathology, as well as poor protection, especially in aged animals, suggesting the possibility that aged humans (a desired target population for the vaccine) may be particularly vulnerable to vaccine-induced effects [24]. Further studies showed that a Venezuelan equine encephalitis vector containing the SARS nucleocapsid (NP) gene also elicited eosinophilic immunopathology in BALB/c mice after challenge without any noticeable protection [25]. Thus, it was hypothesized that the NP antigen of SARS-CoV may be responsible for this property of inactivated vaccines [25,26]. It was also reported that the formalin-inactivated RSV vaccine was associated with lack of antibody affinity maturation followed by poor TLR stimulation, suggesting that NP may not account entirely for immune enhancement [27].

Both Venezuelan equine encephalitis vector containing S-expressing plasmid and alum-adjuvanted recombinant protein (expressed in baculovirus) were shown to elicit protection in BALB/c mice when challenged with live SARS-CoV [28]. However, it remains unclear whether such S protein constructs

**Table 1. Proposed target product profile of the recombinant receptor-binding domain spike protein-based severe acute respiratory syndrome coronavirus vaccine.**

Item	Desired target
Indication	A preventative vaccine to protect against lethal infection caused by the SARS-CoV
Target population	Adults and children >15 years of age
Route of administration	Intramuscular injection
Product presentation	Single-dose vials. 1.0 ml volume of delivery
Dosage schedule	Maximum of two immunizations regardless of age, with the second injection given 2–4 weeks after the first immunization
Warnings and precautions/pregnancy and lactation	Mild-to-moderate local injection site reactions, such as erythema, edema and pain, the character, frequency and severity of which is similar to licensed recombinant protein vaccines. Less than 0.01% risk of urticaria and other systemic allergic reactions. Incidence of SAEs no more than licensed comparator vaccines
Expected efficacy	80% efficacy at preventing SARS-associated deaths
Co-administration	All doses may be co-administered with antiviral drugs and/or other vaccines used in public health emergencies
Shelf-life	4 years
Storage	Refrigeration between 2 and 8°C. Cannot be frozen. Can be out of refrigeration (at temperatures up to 25°C) for up to 72 h
Product registration	Licensure by the US FDA
Target price	Less than US\$10 per dose for use in low- and middle-income countries

SAE: Serious adverse event; SARS: Severe acute respiratory syndrome; SARS-CoV: Severe acute respiratory syndrome coronavirus.

also elicit eosinophilic pathology. One report, for example, has indicated that S protein administered either as a baculovirus-expressed and purified recombinant protein or as a virus-like particle, causes Th2-type pathology with eosinophils, albeit reduced when compared with whole virus or NP protein constructs [28]. Similarly, a SARS vaccine candidate based on recombinant, properly folded, native full-length S protein trimers (expressed in mammalian cells) was also shown to elicit some immune enhancement [29].

Thus, while the recombinant S protein elicits protection, its use as a human vaccine might be limited by host immunopathology.

In order to re-engineer the S protein as a safe and effective vaccine, early studies indicate that the RBD component is highly protective in laboratory animals, while significantly reducing the risk of antibody enhancement of disease [5,6,8–12]. Additional data confirm that sera from SARS convalescent patients contain robust neutralizing antibodies to RBD-S [9,12].

To summarize, our vaccine technology has the potential advantages over existing vaccine technologies that have previously been reported:

- The RBD-S subunit vaccine is expected to be safer than other SARS vaccines described above because the latter may elicit eosinophilic immunopathology or antibody-mediated enhancement of disease, while our vaccine is not expected to cause harmful immune responses.
- The RBD-S subunit vaccine may induce more potent cross-neutralizing antibody responses than other SARS vaccines reported.

### Scientific & technical value of the project

The goal and deliverable is to produce a formulated recombinant RBD-S vaccine under current good manufacturing practices (cGMP), in preparation for clinical testing to assess safety, immunogenicity and efficacy against human SARS-CoV infection.

The key technological objectives involve developing a process for vaccine manufacture utilizing a bacterial, yeast or baculovirus expression system, as well as an inexpensive purification method that maximizes the production of RBD-S antigen component of the vaccine at a minimum cost. These activities would take place through a consortium led by Sabin Vaccine Institute and Texas Children's Hospital Center for Vaccine Development ('Sabin-TCH'), an established PDP with more than a decade-long track record of transitioning recombinant protein vaccines through process development, cGMP and clinical testing [30]. PDPs are non-profit organizations that use industry practices and often partner with for-profit organizations, including biotechnology and multinational pharmaceutical companies, as well as academic and other partners [30]. To develop the RBD-S SARS vaccine, Sabin-TCH will work with the Immune Design Corporation (Seattle, WA, USA) for adjuvant access and formulation, and with the Lindsley F Kimball Research Institute of the New York Blood Center (New York, NY, USA) and the University of Texas Medical Branch (Galveston, TX, USA), in the areas of assay development and confirmatory preclinical efficacy testing.


The Walter Reed Army Institute of Research (WRAIR) will take the lead on the final cGMP manufacture (FIGURE 1). Batch production records will be utilized to ensure reproducibility

and documentation under a quality umbrella. Before transferring technology to the WRAIR Pilot Bioproduction Facility for cGMP manufacture, the final process will be repeated a minimum of three times at the 10-l scale to assure that the vaccine manufacture, from fermentation through formulation, reproducibly results in a product of sufficient yield and quality. In addition, antigen/adjuvant formulation studies will be established both biochemically and by immunological assessment (potency). By performing a minimum of three full pilot runs with quality control for both process and release, the authors will maximize the probability of successful technology transfer for the cGMP manufacture and with optimal yield and purity. Accelerated and long-term stability studies will be conducted on the final formulations, with chemical stability emphasizing provocative tests at high pH to induce deamidation and with selective oxidizing agents to produce oxidation. In addition, stability will be assessed at different time points

after conducting temperature incursion experiments (repeated freeze–thaw cycles and storage at different temperatures). Following completion of process development of the vaccine, the production technology will be transferred to WRAIR for cGMP manufacture. The final product will be subjected to a rigorous battery of assays for lot and product release at Sabin-TCH, WRAIR and other contractors (TABLE 2). Following GLP toxicology testing, a regulatory filing for clinical testing will be submitted to the US FDA.

The major innovations for this product include:

- Producing the RBD-S polypeptide formulation, rather than the complete S protein, with the use of GLA, a novel TLR4 agonist, at point of injection in order to limit vaccine-induced Th2-type immunopathology enhancement.
- The long-term stability of the vaccine will be maximized by evaluating the biochemical and biophysical characteristics [31,32].



**Figure 1. Partnership for the development of a recombinant receptor-binding domain spike protein-based severe acute respiratory syndrome coronavirus vaccine.** The roles of the major partner institutions are described in the text. cGMP: Current good manufacturing practice.

**Table 2. Assays proposed for product characterization and release.**

Assay (application)	Description
SE-HPLC (bulk protein) in-process, release, stability	A SE-HPLC assay is used to analyze recombinant protein samples (crude fermentation supernatant to release of purified bulk). This assay has been developed for other antigens and adapted to RBD-S protein to quantitate target protein from other contaminants during in-process analysis. This assay also serves as a purity assay for RBD-S protein
RP-HPLC (bulk protein) release, stability	Column and condition screening of a variety of RP-HPLC columns (including silica- and polymer-based) has been initiated to provide an assay that can better resolve full-length RBD-S protein from its product-derived cleaved forms (i.e., <1 kDa difference)
Multi-angle light scattering (bulk protein), information only	A multi-angle light scattering detector utilized both online (with SE-HPLC) and offline (batch) has provided valuable data regarding the aggregation and native state of candidate antigens in solution. This assay could be used to detect polydispersity of RBD-S protein and assist in development of a stable formulation
SDS-PAGE (bulk protein and formulation) in-process, release, stability	SDS-PAGE assays under both nonreduced and reducing conditions allow determination of molecular weight, aggregation and purity of samples. A calibrated scanning densitometer allows a relatively quantitative aspect to the assay, involving the determination of limit of detection and linear range of the assay to assess purity and relative quantity. This assay is already adapted and implemented routinely for RBD-S protein analysis
N-terminal sequencing (bulk protein), information only	N-terminal sequencing has been completed on major, minor and contaminant bands to identify whether contaminants are product derived and cleavage sites. SDS-PAGE is completed, then transferred to a PVDF membrane and sent to a contracted service, audited by Sabin's Quality Assurance team and subject to a quality agreement
Mass spectrometry (bulk protein), information only	ESMS is carried out on a Micromass Q-ToF mass spectrometer on samples to monitor mass and any post-translational modifications. This assay will be used for RBD-S protein
OPA (formulation) release, stability	The OPA assay measures protein content by fluorescent detection of primary amines using OPA, a highly sensitive amine-reacting reagent. This technique is especially applicable to vaccine formulations with Alhydrogel® (Brenntag Biosector, Frederikssund, Denmark) where protein content cannot be measured by absorbance at 280 nm and measurement of the percentage of Alhydrogel-bound protein and Alhydrogel-unbound protein is desired. This assay will be adapted to RBD-S protein and its formulation buffers and has been widely used to determine Alhydrogel protein binding efficiency thus far
HCP content in-process, release	Residual HCP impurities in in-process fermentation and purification products and final purified protein samples are evaluated using a slot blot system with a scanning calibrated densitometer with a picogram level of sensitivity

Potency, ELISA (formulation) release and stability to be determined.

ESMS: Electrospray ionization mass spectrometry; HCP: Host cell protein; OPA: O-phthalaldehyde; PVDF: Polyvinylidene fluoride; RBD: Receptor-binding domain; RP-HPLC: Reversed-phase HPLC; S: Spike; SE-HPLC: Size-exclusion HPLC.

Such activities are essential for a vaccine that might require emergency stockpiling. Biophysical assessment is initiated evaluating the recombinant protein using multiple biophysical techniques, including circular dichroism, intrinsic and extrinsic fluorescence, light scattering, and differential scanning calorimetry [31,32]. Collectively, these studies provide specific information concerning the physical state of the protein as a

function of temperature and pH. Biophysical data are then used to design high-throughput screening assays for the identification of potential stabilizers, typically GRAS (generally regarded as safe)-designated excipients and stabilizers that would be acceptable to the FDA and other regulatory agencies [32].

- Preclinical testing in an innovative human angiotensin-converting enzyme 2 transgenic (Tg) mouse model [33,34]. Following



intranasal infection of such Tg mice with SARS-CoV, weight loss and other clinical manifestations become apparent before reaching 100% mortality within 8 days post-infection. In contrast, infected transgene-negative (Tg) mice survive without showing any clinical illness [33]. The severity of the disease developed in these Tg mice makes them valuable for preclinical testing of SARS-CoV vaccines [33,34].

### Summary of immediate project goals & key technological objectives

- Development of a process for high-yield, low-cost expression and purification of the recombinant RBD-S protein antigen, followed by formulation on alum in the presence of GLA;
- Development and qualification of product-specific assays, followed by process and formulation optimization, which incorporates biophysical profiling;
- Completion of three process development and release of pilot runs with quality control;
- Technology transfer for cGMP manufacture (both drug substance and product);
- Formal release of the drug product based on qualified assays and an established formal stability program, which includes vaccine potency;
- A pre-IND meeting followed by completion of a GLP toxicology study;
- IND submission and approval for clinical testing.

The first major deliverable will be an RBD-S protein-based SARS vaccine approved for Phase I clinical testing by the FDA.

### Product & clinical development challenges

The major product challenges includes: selection of antigen expression systems and demonstrating feasibility of scale-up expression with process development at acceptable yields, purity, and stability; preclinical optimization for testing immunological responses and efficacy in laboratory animals; technology transfer for cGMP manufacture; completion of a GLP toxicology study and a clinical development plan that meets requirements of the FDA for 'the animal rule'; and immune correlates of protection [102].

### Selection of antigen expression systems & feasibility of scale-up expression

Three protein expression systems, bacterial (*E. coli*), yeast (*P. pastoris*) and baculovirus (and possibly mammalian cells), will be evaluated in parallel for the feasibility of scalable expression of the recombinant RBD-S protein. Preliminary studies have shown that *E. coli*-expressed RBD-S protein was able to elicit high levels of neutralizing antibodies and protection in an animal model [6]. However, the *E. coli*-expressed protein was not evaluated and formally characterized for its scalability; hence, a more formal evaluation is proposed.

Alternatively, Sabin-TCH has demonstrated that *P. pastoris* is a very robust and sometimes simpler, system for expression. Through Sabin's Human Hookworm Vaccine Initiative, at least two hookworm vaccine antigens, *Na*-ASP-2 and *Na*-GST-1, have already been shown to be suitable for scalability, producing high resolution crystals for X-ray diffraction [35–38], as well as antigens suitable for clinical development [39,40]. For RBD-S, the optimal expression systems will be selected for the subsequent studies based on the following criteria: yield, scalability, purity and stability profile of the RBD-S protein expressed; recognition of the RBD-S by conformational monoclonal antibodies and by human convalescent sera; ability of the RBD-S to inhibit viral infection *in vitro*; and ability of the RBD-S to induce neutralizing antibodies and other protective immune responses in vaccinated mice against subsequent lethal challenge with homologous and heterogeneous strains of SARS-CoV.

### Preclinical optimization & immunological testing

Using the RBD-S protein from the selected expression system and adjuvant formulation, the authors will optimize the immunization regimens to induce the highest levels of immune responses, particularly neutralizing antibodies and protection against subsequent challenge. Immunological and protection studies using wild-type and Tg mice will require optimization in terms of antigen dose and formulation; number and interval of immunizations and route of administration; evaluation of efficacy of the optimized regimen; extended studies to optimize long-term immunological responses and length of protection; and no vaccine-induced Th2-derived immunopathology.

### Process development, characterization, formulation & stability evaluation

Following selection of an expression system, a research cell bank will be generated, and expression of RBD-S in a 10-l bioreactor (fermentation) will be optimized. To improve yield, purity, stability and activity of the soluble protein, the optimal growth and induction conditions for the fermentation processes will be determined first using flasks. Once the best parameters are selected, they will be evaluated at the 10-l scale, with the aim of obtaining a reproducible high-yield production (minimum of 100 mg/l target). The major components of this activity will include: production of the RBD-S protein at the 10-l fermentation scale and purification using ion-exchange, hydrophobic interaction or other column chromatographies; documentation of three successive process development runs (fermentation and protein purification); development of product-specific assays to monitor the antigen identity, purity and integrity during fermentation and purification, and for clinical lot release; development of buffer and adjuvant formulations, and product stability studies.

### Technology transfer, cGMP manufacture & lot release, GLP toxicology & IND preparation

The process development and formulation technology for the selected RBD-S vaccine will be transferred to WRAIR. Developed and qualified assays will be used for lot release and

stability/potency testing. The protein-specific assay will be conducted at Sabin-TCH (TABLE 2), while other regulated assays will be contracted to industrial partners.

These assays can provide important feedback about the purity, stability and integrity of the molecule essential to guide process development and formulation. Additionally, new assays and technology are typically identified and implemented, where appropriate, to ensure specific characteristics of the molecule are maintained. Several of these assays, either currently used or being developed for RBD-S protein, have been qualified by Sabin-TCH for other recombinant antigens for Phase I clinical trials. Following a pre-IND meeting with the FDA, a GLP toxicology testing will be performed. Sabin Vaccine Institute will serve as the sponsor and prepare the IND for the vaccine. The initial design of a global access roadmap for vaccine uptake and distribution will also be drafted.

### Clinical development & pivotal animal efficacy studies

The safety of RBD-S vaccine will be evaluated in a Phase I clinical trial. Under a new subpart of the US Code of Federal Regulations, it is acceptable to approve new drugs through pivotal animal efficacy studies based on the 'Animal Rule' [41] when human efficacy studies are not ethical or feasible because they would deliberately expose healthy human volunteers to a lethal agent [41]. While safety and immunogenicity studies with the RBD-S vaccine are feasible, human challenge studies with SARS-CoV would not be considered safe or ethical.

Therefore, pivotal animal efficacy studies will be conducted using at least one animal model exhibiting host pathogenic and immune responses that are matched as closely as possible to those

of humans [41]. Besides the nonhuman primate model, the diseases and lethality revealed by ACE2 transgenic mice infected by clinical isolates of SARS-CoV (e.g., Urbani) and those elicited by wild-type Balb/c mice in response to a mouse-adapted strain of SARS-CoV, designated MA-15) [33,34,42] make them among the best models for evaluating *in vivo* efficacy of SARS vaccines.

Determination of the virus neutralizing antibody titers among vaccine recipients in the pivotal animal efficacy study is an important bridging of animal protection data to humans. To serve as an immune correlate of protection, antibody titers would have to equal or exceed the neutralizing antibody titers and amounts found in the sera of convalescent patients, with desired quantities of specific neutralizing antibodies to be determined in consultation with expert clinical virologists. In addition, levels of antibody affinity and avidity may need to be established using surface plasmon resonance and other technologies [43].

### Expert commentary

Following the initial discovery of the SARS-CoV as the etiologic agent of human SARS in 2003, an international effort has been underway to develop and test prototype vaccines. These intensive studies determined that when delivered as an injectable vaccine, inactivated SARS-CoV can elicit protective neutralizing antibodies. However, such vaccines also caused a Th2-derived immunoenhancing pathology bearing resemblance to the immunopathology that derailed efforts to produce an inactivated RSV vaccine more than four decades ago. Subsequent efforts determined that protective neutralizing antibodies were directed primarily against S protein responsible for receptor-binding, but even vaccines comprised of the full-length S protein can elicit immunopathology, albeit in reduced

### Key issues

- SARS emerged as a new and highly lethal human infection in South China in 2002. SARS-CoV is considered as a potential bioweapons threat.
- Vaccines are particularly critical for protecting high-risk groups, including the elderly, healthcare workers and laboratory personnel.
- The proposed RBD-S SARS vaccine is a recombinant protein-based vaccine formulated with alum and a glucopyranosyl lipid A Toll-like receptor 4 agonist novel adjuvant.
- The recombinant protein is comprised of a 193 amino acid (21 kDa) polypeptide corresponding to the minimal RBD of the SARS-CoV S protein.
- The vaccine is initially indicated for adults and individuals over the age of 15 who are considered at greatest risk of mortality from SARS. The vaccine would be administered as an injectable product to protect against lethal SARS-CoV infection, and as either a single dose or two doses spaced closely together (within 2–4 weeks), so that vulnerable populations could be rapidly immunized in an outbreak setting. The high stability of the RBD-S SARS vaccine is such that it can be stockpiled for up to 4 years at 2–8°C for emergency use.
- Unlike other prototype vaccines, the RBD-S SARS vaccine is designed to reduce the likelihood of eosinophilic (Th2) immunopathology.
- The immediate goals include the development of a process for high-yield, low-cost expression and purification of the RBD-S protein antigen, followed by formulation on alum and with the use of glucopyranosyl lipid A at point of injection; development and qualification of product-specific assays, followed by process and formulation optimization, which incorporates biochemical and biophysical assessments; completion of three process development pilot runs with quality control testing; technology transfer for the cGMP manufacturing processes of both drug substance and drug product; formal release of the drug product and establishment of a formal stability program, which includes vaccine potency; a pre-investigational new drug meeting followed by completion of a good laboratory practice toxicology study and an investigational new drug submission followed by approval for clinical testing. Having met these objectives, we expect that the first major deliverable, an RBD-S SARS vaccine will be approved for Phase I clinical testing by the US FDA.
- Vaccine development will be led by a nonprofit product development partnerships in collaboration with key academic, military and industrial partners.

amounts. Therefore, efforts have, instead, focused on a subunit vaccine comprised of only the 193-mer RBD-S, the essential component responsible for receptor binding. In laboratory animals, recombinant RBD-S subunit vaccines elicited protection comparable with the S protein-based vaccines, but with minimal immune enhancement of immunopathology. A prototype recombinant RBD-S SARS vaccine formulated on alum, together with GLA at point of injection, is under development by the activities of a nonprofit PDP in collaboration with key academic, industrial and military partners.

### Five-year view

Within 5 years, it is anticipated that a prototype recombinant RBD-S SARS vaccine formulated on alum, and with GLA at point of injection, will have completed cGMP manufacture at WRAIR. Following lot release and GLP toxicology testing, the

vaccine will be ready for an IND submission and Phase I clinical testing. A full clinical development plan leading to product licensure will need outside consultation to confirm the quantity and quality, as well as affinity and avidity, of virus neutralizing antibodies required for protection.

### Financial & competing interests disclosure

Research reported in this publication is supported by the National Institute of Allergy and Infectious Diseases of the NIH under Award Number R01AI098775. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

No writing assistance was utilized in the production of this manuscript.

### References

Papers of special note have been highlighted as:

• of interest

•• of considerable interest

- 1 Du L, He Y, Zhou Y, Liu S, Zheng BJ, Jiang S. The spike protein of SARS-CoV – a target for vaccine and therapeutic development. *Nat. Rev. Microbiol.* 7(3), 226–236 (2009).
- Excellent review of the role of the spike protein in vaccine development.
- 2 Roper RL, Rehm KE. SARS vaccines: where are we? *Expert Rev. Vaccines* 8(7), 887–898 (2009).
- 3 Jaume M, Yip MS, Kam YW *et al.* SARS CoV subunit vaccine: antibody-mediated neutralisation and enhancement. *Hong Kong Med. J.* 18(Suppl. 2), 31–36 (2012).
- 4 Wong SK, Li W, Moore MJ, Choe H, Farzan M. A 193-amino acid fragment of the SARS coronavirus S protein efficiently binds angiotensin-converting enzyme 2. *J. Biol. Chem.* 279(5), 3197–3201 (2004).
- First description of the receptor-binding domain and its target receptor.
- 5 Du L, Zhao G, Li L *et al.* Antigenicity and immunogenicity of SARS-CoV S protein receptor-binding domain stably expressed in CHO cells. *Biochem. Biophys. Res. Commun.* 384(4), 486–490 (2009).
- 6 Du L, Zhao G, Chan CC *et al.* Recombinant receptor-binding domain of SARS-CoV spike protein expressed in mammalian, insect and *E. coli* cells elicits potent neutralizing antibody and protective immunity. *Virology* 393(1), 144–150 (2009).
- Report of different constructs eliciting neutralizing antibody to the severe acute respiratory syndrome coronavirus (SARS-CoV).
- 7 Li W, Moore MJ, Vasilieva N *et al.* Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus. *Nature* 426(6965), 450–454 (2003).
- 8 He Y, Lu H, Siddiqui P, Zhou Y, Jiang S. Receptor-binding domain of severe acute respiratory syndrome coronavirus spike protein contains multiple conformation-dependent epitopes that induce highly potent neutralizing antibodies. *J. Immunol.* 174(8), 4908–4915 (2005).
- 9 He Y, Zhu Q, Liu S *et al.* Identification of a critical neutralization determinant of severe acute respiratory syndrome (SARS)-associated coronavirus: importance for designing SARS vaccines. *Virology* 334(1), 74–82 (2005).
- 10 He Y, Li J, Li W, Lustigman S, Farzan M, Jiang S. Cross-neutralization of human and palm civet severe acute respiratory syndrome coronaviruses by antibodies targeting the receptor-binding domain of spike protein. *J. Immunol.* 176(10), 6085–6092 (2006).
- 11 Du L, Zhao G, He Y *et al.* Receptor-binding domain of SARS-CoV spike protein induces long-term protective immunity in an animal model. *Vaccine* 25(15), 2832–2838 (2007).
- Evidence for protective efficacy of the receptor-binding domain subunit of the spike protein inducing protection.
- 12 Cao Z, Liu L, Du L *et al.* Potent and persistent antibody responses against the receptor-binding domain of SARS-CoV spike protein in recovered patients. *Viol. J.* 7, 299 (2010).
- 13 Chen Z, Zhang L, Qin C *et al.* Recombinant modified vaccinia virus Ankara expressing the spike glycoprotein of severe acute respiratory syndrome coronavirus induces protective neutralizing antibodies primarily targeting the receptor binding region. *J. Virol.* 79(5), 2678–2688 (2005).
- 14 Du L, Zhao G, Chan CC *et al.* A 219-mer CHO-expressing receptor-binding domain of SARS-CoV S protein induces potent immune responses and protective immunity. *Viral Immunol.* 23(2), 211–219 (2010).
- 15 Fox CB, Friede M, Reed SG, Ireton GC. Synthetic and natural TLR4 agonists as safe and effective vaccine adjuvants. *Subcell. Biochem.* 53, 303–321 (2010).
- 16 Baldwin SL, Shaverdian N, Goto Y *et al.* Enhanced humoral and Type 1 cellular immune responses with Fluzone adjuvanted with a synthetic TLR4 agonist formulated in an emulsion. *Vaccine* 27(43), 5956–5963 (2009).
- 17 Reed SG, Bertholet S, Coler RN, Friede M. New horizons in adjuvants for vaccine development. *Trends Immunol.* 30(1), 23–32 (2009).
- 18 Behzad H, Huckriede AL, Haynes L *et al.* GLA-SE, a synthetic Toll-like receptor 4 agonist, enhances T-cell responses to influenza vaccine in older adults. *J. Infect. Dis.* 205(3), 466–473 (2012).
- Paper on the potential of glucopyranosyl lipid A as a new synthetic adjuvant.
- 19 Olsen CW. A review of feline infectious peritonitis virus: molecular biology, immunopathogenesis, clinical aspects, and vaccination. *Vet. Microbiol.* 36(1–2), 1–37 (1993).
- 20 Weiss RC, Scott FW. Antibody-mediated enhancement of disease in feline infectious peritonitis: comparisons with dengue hemorrhagic fever. *Comp. Immunol.*



- Microbiol. Infect. Dis.* 4(2), 175–189 (1981).
- 21 Perlman S, Dandekar AA. Immunopathogenesis of coronavirus infections: implications for SARS. *Nat. Rev. Immunol.* 5(12), 917–927 (2005).
  - 22 Castilow EM, Olson MR, Varga SM. Understanding respiratory syncytial virus (RSV) vaccine-enhanced disease. *Immunol. Res.* 39(1-3), 225–239 (2007).
- **Good overview of the problem of immune enhancement as it relates to respiratory syncytial virus.**
- 23 Collins PL, Graham BS. Viral and host factors in human respiratory syncytial virus pathogenesis. *J. Virol.* 82(5), 2040–2055 (2008).
  - 24 Bolles M, Deming D, Long K *et al.* A double-inactivated severe acute respiratory syndrome coronavirus vaccine provides incomplete protection in mice and induces increased eosinophilic proinflammatory pulmonary response upon challenge. *J. Virol.* 85(23), 12201–12215 (2011).
  - 25 Deming D, Sheahan T, Heise M *et al.* Vaccine efficacy in senescent mice challenged with recombinant severe acute respiratory syndrome-coronavirus bearing epidemic and zoonotic spike variants. *PLoS Med.* 3(12), e525 (2006).
  - 26 Yasui F, Kai C, Kitabatake M *et al.* Prior immunization with severe acute respiratory syndrome (SARS)-associated coronavirus (SARS-CoV) nucleocapsid protein causes severe pneumonia in mice infected with SARS-CoV. *J. Immunol.* 181(9), 6337–6348 (2008).
  - 27 Delgado MF, Coviello S, Monsalvo AC *et al.* Lack of antibody affinity maturation due to poor Toll-like receptor stimulation leads to enhanced respiratory syncytial virus disease. *Nat. Med.* 15(1), 34–41 (2009).
  - 28 Tseng CT, Sbrana E, Iwata-Yoshikawa N *et al.* Immunization with SARS coronavirus vaccines leads to pulmonary immunopathology on challenge with the SARS virus. *PLoS ONE* 7(4), e35421 (2012).
- **Description of the immunopathology that can potentially result from SARS-CoV vaccines.**
- 29 Jaume M, Yip MS, Kam YW *et al.* SARS CoV subunit vaccine: antibody-mediated neutralisation and enhancement. *Hong Kong Med. J.* 18(Suppl. 2), 31–36 (2012).
  - 30 Hotez P. A handful of ‘antipoverty’ vaccines exist for neglected diseases, but the world’s poorest billion people need more. *Health Aff. (Millwood)* 30(6), 1080–1087 (2011).
- **Overview of the role of product development partnerships in producing vaccines and other biotechnologies.**
- 31 Maddux NR, Joshi SB, Volkin DB, Ralston JP, Middaugh CR. Multidimensional methods for the formulation of biopharmaceuticals and vaccines. *J. Pharm. Sci.* doi:10.1002/jps.22618 (2011) (Epub ahead of print).
- **Overview of the use of biochemical and biophysical assessments in vaccine development.**
- 32 Plieskatt JL, Rezende WC, Olsen CM *et al.* Advances in vaccines against neglected tropical diseases: enhancing physical stability of a recombinant hookworm vaccine through biophysical and formulation studies. *Hum. Vaccin. Immunother.* 8(6), 765–776 (2012).
  - 33 Tseng CT, Huang C, Newman P *et al.* Severe acute respiratory syndrome coronavirus infection of mice transgenic for the human Angiotensin-converting enzyme 2 virus receptor. *J. Virol.* 81(3), 1162–1173 (2007).
- **First description of a unique laboratory animal model for SARS-CoV.**
- 34 McCray PB Jr, Pewe L, Wohlford-Lenane C *et al.* Lethal infection of K18-hACE2 mice infected with severe acute respiratory syndrome coronavirus. *J. Virol.* 81, 813–821 (2007).
  - 35 Goud GN, Bottazzi ME, Zhan B *et al.* Expression of the *Necator americanus* hookworm larval antigen Na-ASP-2 in *Pichia pastoris* and purification of the recombinant protein for use in human clinical trials. *Vaccine* 23(39), 4754–4764 (2005).
  - 36 Asojo OA, Goud G, Dhar K *et al.* X-ray structure of Na-ASP-2, a pathogenesis-related-1 protein from the nematode parasite, *Necator americanus*, and a vaccine antigen for human hookworm infection. *J. Mol. Biol.* 346(3), 801–814 (2005).
- 37 Asojo OA, Homma K, Sedlacek M *et al.* X-ray structures of Na-GST-1 and Na-GST-2 two glutathione *S*-transferase from the human hookworm *Necator americanus*. *BMC Struct. Biol.* 7, 42 (2007).
  - 38 Goud GN, Deumic V, Gupta R *et al.* Expression, purification, and molecular analysis of the *Necator americanus* glutathione *S*-transferase 1 (Na-GST-1): a production process developed for a lead candidate recombinant hookworm vaccine antigen. *Protein Expr. Purif.* 83(2), 145–151 (2012).
  - 39 Bethony JM, Simon G, Diemert DJ *et al.* Randomized, placebo-controlled, double-blind trial of the Na-ASP-2 hookworm vaccine in unexposed adults. *Vaccine* 26(19), 2408–2417 (2008).
  - 40 Hotez PJ, Bethony JM, Diemert DJ, Pearson M, Loukas A. Developing vaccines to combat hookworm infection and intestinal schistosomiasis. *Nat. Rev. Microbiol.* 8(11), 814–826 (2010).
  - 41 Burns DL. Licensure of vaccines using the Animal Rule. *Curr. Opin. Virol.* 2(3), 353–356 (2012).
  - 42 Roberts A, Deming D, Paddock CD *et al.* A mouse-adapted SARS-coronavirus causes disease and mortality in Balb/c mice. *PLoS Pathogens* 3, e5 (2007).
  - 43 Hearty S, Conroy PJ, Ayyar BV, Byrne B, O’Kennedy R. Surface plasmon resonance for vaccine design and efficacy studies: recent applications and future trends. *Expert Rev. Vaccines* 9(6), 645–664 (2010).

## Websites

- 101 Biodefence category A, B, C pathogens. [www.niaid.nih.gov/topics/biodefenserelated/biodefense/pages/cata.aspx](http://www.niaid.nih.gov/topics/biodefenserelated/biodefense/pages/cata.aspx) (Accessed 13 May 2012)
- 102 CFR – Code of Federal Regulations Title 21. [www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcfr/CFRSearch.cfm?CFRPart=314&showFR=1&subpartNo=21.5.0.1.1.4.9](http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcfr/CFRSearch.cfm?CFRPart=314&showFR=1&subpartNo=21.5.0.1.1.4.9) (Accessed 27 May 2012)