

Coronavirus infection, ER stress and Apoptosis

TO SING FUNG and Ding Xiang Liu

Journal Name: Frontiers in Microbiology

ISSN: 1664-302X

Article type: Review Article

Received on: 31 Mar 2014

Accepted on: 29 May 2014

Provisional PDF published on: 29 May 2014

www.frontiersin.org: www.frontiersin.org

Citation: Fung T and Liu D(2014) Coronavirus infection, ER stress and Apoptosis. *Front. Microbiol.* 5:296. doi:10.3389/fmicb.2014.00296

[/Journal/Abstract.aspx?s=1161&name=virology&ART_DOI=10.3389/fmicb.2014.00296](#): [/Journal/Abstract.aspx?s=1161&name=virology&ART_DOI=10.3389/fmicb.2014.00296](#)

(If clicking on the link doesn't work, try copying and pasting it into your browser.)

Copyright statement: © 2014 Fung and Liu. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](#). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

This Provisional PDF corresponds to the article as it appeared upon acceptance, after rigorous peer-review. Fully formatted PDF and full text (HTML) versions will be made available soon.

Coronavirus infection, ER stress and Apoptosis

To Sing Fung and Ding Xiang Liu*

School of Biological Sciences, Nanyang Technological University, 60 Nanyang Drive,
Singapore 637551

Running title: coronavirus and ER stress response

*Corresponding author: Phone +65 63162862; Fax +65 67936828; Email dxliu@ntu.edu.sg

Abstract

The replication of coronavirus, a family of important animal and human pathogens, is closely associated with the cellular membrane compartments, especially the endoplasmic reticulum (ER). Coronavirus infection of cultured cells was previously shown to cause ER stress and induce the unfolded protein response (UPR), a process that aims to restore the ER homeostasis by global translation shutdown and increasing the ER folding capacity. However under prolonged ER stress, UPR can also induce apoptotic cell death. Accumulating evidence from recent studies has shown that induction of ER stress and UPR may constitute a major aspect of coronavirus-host interaction. Activation of the three branches of UPR modulates a wide variety of signaling pathways, such as mitogen-activated protein (MAP) kinases activation, autophagy, apoptosis and innate immune response. ER stress and UPR activation may therefore contribute significantly to the viral replication and pathogenesis during coronavirus infection. In this review, we summarize current knowledge on coronavirus-induced ER stress and UPR activation, with emphasis on their cross-talking to apoptotic signaling.

Introduction

Coronaviruses are a family of enveloped viruses with positive sense, non-segmented, single stranded RNA genomes. Many coronaviruses are important veterinary pathogens. For example, avian infectious bronchitis virus (IBV) reduces the performance of both meat-type and egg-laying chickens and causes severe economic loss to the poultry industry worldwide [1]. Certain coronaviruses, such as HCoV-229E and HCoV-OC43, infect humans and account for a significant percentage of adult common colds [2,3]. Moreover, in 2003, a highly pathogenic human coronavirus (SARS-CoV) was identified as the causative agent of severe acute respiratory syndrome (SARS) with high mortality rate and led to global panic [4]. Afterwards, it was found that the SARS-CoV was originated from bat and likely jumped to humans via some intermediate host (palm civets) [5,6]. Recently, a live SARS-like coronavirus was isolated from fecal samples of Chinese horseshoe bats, which could use the SARS-CoV cellular receptor - human angiotensin converting enzyme II (ACE2) for cell entry [7]. This indicates an intermediate host may not be necessary and direct human infection by some bat coronaviruses is possible. Moreover, a novel human coronavirus – the Middle East respiratory syndrome coronavirus (MERS-CoV), emerged in Saudi Arabia in September 2012 [8]. Although the risk of sustained human-to-human transmission is considered low, infection

of MERS-CoV causes ~50% mortality in patients with comorbidities [9]. Initial studies had pointed to bats as the source of MERS-CoV [10], however, accumulating evidence strongly suggested the dromedary camels to be the natural reservoirs and animal source of MERS-CoV [11,12]. Thus, coronaviruses can cross the species barrier to become lethal human pathogens, and studies on coronaviruses are both economically and medically important.

Taxonomically, the family *Coronaviridae* is classified into two subfamilies, the *coronavirinae* and the *torovirinae*. The *coronavirinae* is further classified into three genera, namely the Alphacoronavirus, Betacoronavirus and Gammacoronavirus [13]. The classification was originally based on antigenic relationships and later confirmed by sequence comparisons of entire viral genomes [14]. Almost all Alphacoronaviruses and Betacoronaviruses have mammalian hosts, including humans. In contrast, Gammacoronaviruses have mainly been isolated from avian hosts.

Morphologically, coronaviruses are spherical or pleomorphic in shape with a mean diameter of 80-120 nm. They are characterized by the large (20 nm) “club-like” projections on the surface, which are the heavily glycosylated trimeric spike (S) proteins [13]. Two additional structural proteins are found on the envelope. The abundant membrane (M) proteins give the virion its shape, whereas the small envelope (E) proteins play an essential role during assembly [15,16]. Inside the envelope, the helical nucleocapsid is formed by binding of the nucleocapsid (N) proteins on the genomic RNA in a beads-on-a-string fashion. The genome, ranging from 27,000 to 32,000 nucleotides in size, is the largest RNA genomes known to date.

Coronavirus infection starts with receptor binding via the S protein (Figure 1). The S proteins of most coronaviruses are cleaved by host protease into two functional subunits: an N-terminal receptor binding domain (S1) and a C-terminal domain (S2) responsible for membrane fusion [17,18,19]. The interaction between the cell surface receptor and the S1 subunit is the major determinant of the tropism of coronaviruses [20]. Upon receptor binding of S1, a conformational change is triggered in the S2 subunit, exposing its hidden fusion peptide for insertion into the cellular membrane. This is followed by the packing of the two heptad repeats in the three monomers into a six-helix bundle fusion core. This close juxtaposition of the viral and cellular membrane enables fusion of the lipid bilayers, and the viral nucleocapsid is thus delivered into the cytoplasm [13].

After uncoating, the genomic RNA first acts as an mRNA for translation of the replicase polyprotein. The replicase gene consists of two open reading frames (ORF1a and ORF1b). Translation of ORF1a produces the polyprotein 1a (pp1a). Meanwhile, a ribosomal

frameshifting occurs at the junction of ORF1a and ORF1b, allowing translation to continue onto ORF1b, producing a larger polyprotein 1ab (pp1ab) [21]. Autoproteolytic cleavage of pp1a produces 11 non-structural proteins (nsp1-nsp11), while cleavage of pp1ab produces 15 non-structural proteins (nsp1-nsp10, nsp12-nsp16). The functions of these nsps are partially understood. Particularly, the autoproteolytic cleavage relies on nsp3 (a papain-like proteinase) and nsp5 (the main proteinase), whereas the RNA-dependent RNA polymerase (RdRp) is contained within nsp12 [22,23].

Using the genomic RNA as a template, the replicase then synthesizes the negative sense genomic RNAs, which are used as templates for synthesizing progeny positive sense RNA genomes. On the other hand, through discontinuous transcription of the genome, the replicase synthesizes a nested set of subgenomic RNAs (sgRNAs) [24]. Replication and transcription of the coronavirus genome involve the formation of the replication/transcription complexes (RTCs), which are anchored to the intracellular membranes via the multi-spanning transmembrane proteins nsp3, nsp4 and nsp6 [25]. Also, inside the infected cells, coronaviruses induce modification of the intracellular membrane network and formation of the double membrane vesicles (DMVs) [26]. Several studies have shown that the DMVs are closely associated with the coronavirus RTCs and the *de novo* synthesized viral RNAs [27,28]

The sgRNAs are translated into structural proteins and accessory proteins. Transmembrane structural proteins (S, M and E) are synthesized, inserted and folded in the endoplasmic reticulum (ER) and transported to the ER-Golgi intermediate compartment (ERGIC). The N proteins are translated in the cytoplasm and encapsidate the nascent progeny genomic RNA to form the nucleocapsids. Virion assembly occurs in the ERGIC and is likely to be orchestrated by the M protein through protein-protein interactions [13].

The virions budded into the ERGIC are exported through secretory pathway in smooth-wall vesicles, which ultimately fuse with the plasma membrane and release the mature virus particles [29]. For some coronaviruses, a portion of the S protein escapes from viral assembly and is secreted to the plasma membrane. These S proteins cause fusion of the infected cell with neighboring uninfected cells, resulting in the formation of a large multinucleated cell known as a syncytium, which enables the virus to spread without being released into the extracellular space [13].

In eukaryotic cells, ER is the major site for synthesis and folding of secreted and transmembrane proteins. The amount of protein entering the ER can vary substantially under different physiological states and environmental conditions. When protein synthesis surpasses the folding capacity, unfolded proteins accumulate in the ER and lead to ER stress. ER stress

can also be activated by excessive lipids or pro-inflammatory cytokines [30,31]. To maintain homeostasis, cells have evolved signaling pathways that are collectively known as the unfolded protein response (UPR) [32]. The UPR signaling starts with the unfolded proteins activating the three ER stress transducers: PKR-like ER protein kinase (PERK), activating transcriptional factor-6 (ATF6) or inositol-requiring protein-1 (IRE1) (Figure 2). Once activated, these sensors transmit the signal across the ER membrane to the cytosol and the nucleus, and the cell responds by lowering the protein synthesis and increasing the ER folding capacity. If homeostasis cannot be re-established, apoptosis is induced for the benefit of the entire organism [33].

In this review, current studies on the involvement of the UPR in coronavirus infection and pathogenesis will be summarized. The role of UPR activation in host response, in particular the induction of apoptosis, will also be reviewed.

Coronavirus infection and ER stress

Global proteomic and microarray analyses have shown that the expression of several genes related to the ER stress, such as glucose-regulated protein 94 (GRP94) and glucose-regulated protein 78 (GRP78, also known as immunoglobulin heavy chain-binding protein, or BiP), is up-regulated in cells infected with SARS-CoV or in cells over-expressing the SARS-CoV S2 subunit [34,35]. Using a luciferase reporter system, Chan et al found that both GRP94 and GRP78 were induced in SARS-CoV infected FRhK4 cells [36]. Consistently, the mRNA level of homocysteine-inducible, endoplasmic reticulum stress-inducible, ubiquitin-like domain member 1 (HERPUD1), an ER stress marker, was up-regulated in L cells infected with mouse hepatitis virus (MHV) or SARS-CoV [37]. Data from this group have shown a similar induction of ER stress in IBV infected Vero, H1299 and Huh-7 cells (unpublished observations). Although no parallel studies have been performed on Alphacoronaviruses, it is likely that all three genera of coronaviruses may induce ER stress in the infected cells.

Current evidence suggests the following three main mechanisms.

Formation of double membrane vesicles (DMVs). It is well-known that the replication of many plus-stranded RNA viruses induces modification of cellular membranes [38]. Among them, coronaviruses have been shown to induce the formation of DMVs in infected cells [39]. Based on immunocytochemistry electron microscopy data, the DMVs co-localize with coronavirus major replicase proteins and are presumably the sites where coronavirus RTCs are located [27,28]. Indeed, DMVs are induced in HEK293T cells co-expressing the SARS-CoV nsp3, nsp4 and nsp6, which are all multispinning transmembrane non-structural proteins

[40]. There have been different perspectives regarding the origin of the coronavirus-induced DMVs. The late endosomes, autophagosomes and the early secretory pathway have all been implicated as the membrane source of DMVs [41,42,43]. Also, co-localization has been observed between SARS-CoV non-structural proteins and protein disulfide isomerase (PDI), an ER marker [44]. Using high resolution electron tomography, Knoops et al have shown that infection of SARS-CoV reorganizes the ER into a reticulovesicular network, which consists of convoluted membranes and interconnected DMVs [26]. Recently, Reggiori et al have proposed a model in which coronaviruses hijack the EDEMosomes to derive ER membrane for DMVs formation [45]. The EDEMosomes are COPII-independent vesicles that export from the ER, which are normally used to fine-tune the level of ER degradation enhancer, mannosidase alpha-like 1 (EDEM1), a regulator of ER-associated degradation (ERAD) [46]. It has been demonstrated that MHV infection causes accumulation of EDEM1 and osteosarcoma amplified 9 (OS-9, another EDEMosome cargo), and that both EDEM1 and OS-9 co-localize with the RTCs of MHV [45]. These results thus add mechanical evidence to support the ER-origin of the coronavirus-induced DMVs.

Glycosylation of coronaviral structural proteins. Except for the N protein, all coronavirus structural proteins are transmembrane proteins synthesized in the ER. The M protein, which is the most abundant component of the virus particle, is known to undergo either O-linked (for most betacoronaviruses) or N-linked (for all alpha- and gammacoronaviruses) glycosylation in the ER [47,48,49]. The glycosylation of M protein is proposed to play a certain function in alpha interferon induction and *in vivo* tissue tropism [50,51,52]. The pre-glycosylated S monomers are around 128-160 kDa, whereas sizes can reach 150-200 kDa post-glycosylation (exclusively N-linked), indicating that the S protein is highly glycosylated [13]. At least for transmissible gastroenteritis coronavirus (TGEV), glycosylation is presumed to facilitate monomer folding and trimerization [53]. Moreover, the glycans on SARS-CoV S proteins have been shown to bind C-type lectins DC-SIGN (dendritic cell-specific intercellular adhesion molecule-3-grabbing non-integrin) and L-SIGN (liver lymph node-specific intercellular adhesion molecule-3-grabbing non-integrin), which can serve as alternative receptors for SARS-CoV independent of the major receptor ACE2 [54]. The folding, maturation and assembly of the gigantic S trimeric glycoprotein rely heavily on the protein chaperons inside the ER, such as calnexin. In fact, the N-terminal part of the S2 domain of SARS-CoV S protein has been found to interact with calnexin, and knock-down of calnexin decreases the infectivity of pseudotyped lentivirus carrying the SARS-CoV S protein [55]. Also, treatment of α -glucosidase inhibitors, which inhibit the interactions of calnexin with its

substrates, dose dependently inhibits the incorporation of S into pseudovirus and suppresses SARS-CoV replication in cell cultures [55]. During coronavirus replication, massive amount of structural proteins is synthesized to assembly progeny virions. The production, folding and modification of these proteins undoubtedly increase the workload of the ER.

Depletion of ER lipid during the budding of virions. Budding of coronaviruses occurs in the ERGIC, which is a structural and functional continuance of the ER. Thus the release of mature virions by exocytosis in effect depletes the lipid component of the ER. Taken together, coronavirus infection results in: 1. massive morphological rearrangement of the ER; 2. significant increase ER burden for protein synthesis, folding and modification; and 3. extensive depletion of ER lipid component. These factors together may contribute to the coronavirus-induced ER stress.

In the following sections, the activation of the three individual branches of the UPR by coronavirus infection will be discussed in detail.

The PERK branch of UPR

PERK-eIF2 α -ATF4 Signaling pathway. The PERK branch of the UPR is believed to be activated first in response to ER stress [56]. Activation of PERK begins with the dissociation from ER chaperon BiP, followed by oligomerization and auto-phosphorylation. Activated PERK then phosphorylates the α -subunit of eukaryotic initiation factor 2 (eIF2 α). Phosphorylated eIF2 α forms a stable complex with and inhibits the turnover of eIF2B, a guanine nucleotide exchange factor that recycles inactive eIF2-GDP to active eIF2-GTP. This results in a general shutdown of cellular protein synthesis and reduces the protein flux into the ER [32]. Besides PERK, three other kinases are known to phosphorylate eIF2 α , namely the protein kinase RNA-activated (PKR), heme-regulated inhibitor kinase (HRI) and general control non-derepressible-2 (GCN2) [32]. PKR is induced by interferon (IFN) and activated by the binding of double-stranded RNA (dsRNA) after virus infection [57]. HRI is activated in red blood cells and hepatocytes by low levels of heme [58]. GCN2 senses amino acid deficiency and is activated via binding to uncharged transfer RNAs [59]. Due to common outcome (eIF2 α phosphorylation and translation suppression), activation of these kinases is collectively known as the integrated stress response (ISR) [32].

Interestingly, the mRNAs of certain genes contain small open reading frames in their 5' UTR and bypass the eIF2 α -dependent translation block. One of these is the activating transcription factor 4 (ATF4), which is preferentially translated under ISR. ATF4 in turn trans-activates genes involved in amino acid metabolism, redox reactions and stress response.

One of ATF4's target genes is the growth arrest and DNA damage-inducible protein 153 (GADD153, also known as C/EBP homologous protein, or CHOP). GADD153 induces the growth arrest and DNA damage-inducible protein 34 (GADD34), which recruits protein phosphatase 1 (PP1) to dephosphorylate eIF2 α and release the translation block. To this end, if ER stress is resolved, normal protein synthesis can be resumed. However, if ER stress persists, GADD153 can induce apoptosis by suppressing the anti-apoptotic protein **B-cell lymphoma 2 (Bcl-2)** and inducing the pro-apoptotic proteins such as Bcl-2-interacting mediator of cell death (Bim) [60]. **GADD153 also activates endoplasmic reticulum oxidoreductin-1 α (ERO1 α)**, which encodes an ER oxidase. The increase protein influx to a hyper-oxidizing ER aggravates ER stress and induces apoptosis [61] (Figure 3).

Involvement of the PERK pathway during viral infections. Translation attenuation has been widely observed as a defensive mechanism of the host cells against viral infection. By reducing the translation of viral proteins, virus replication is hampered and the spread of infection is limited, giving enough time for the immune system to initiate effective antiviral responses. Among the four eIF2 α kinases, PKR, due to its interferon-inducible nature and specific recognition of viral dsRNAs, plays an especially important role in inducing translation attenuation in virus infected cells [62]. It is therefore not surprising, that viruses have evolved various mechanisms to counteract PKR. For example, the non-structural 5A (NS5A) protein of hepatitis C virus directly interact with the catalytic site of PKR, whereas the NS1 protein in the influenza A virus binds to dsRNAs and thus blocks PKR activation [63,64].

During virus infection, massive production of viral proteins can overload the folding capacities of ER and lead to activation of another eIF2 α kinase – PERK. Activation of PERK has been observed in cells infected with various DNA and RNA viruses, such as vesicular stomatitis virus, bovine viral diarrhea virus and herpes simplex virus 1 (HSV1), to name just a few [65,66,67]. However, similar to PKR, viruses have adopted counter measures to inhibit PERK mediated translation attenuation. For example, the E2 protein of hepatitis C virus (HCV) and the glycoprotein gB of HSV1 binds to PERK and inhibits its kinase activity to rescue translation [68,69].

Activation of PERK pathway during coronaviruses infection and its involvement in coronavirus-induced apoptosis. There have been diverging results on the activation of PKR and/or PERK during coronavirus infection. In an early study, it has been found that there is minimal transcriptional activation of PKR and another interferon stimulated gene, 2'5'-oligoadenylate synthetase (OAS) in cells infected with MHV-1 [70]. In a separate study,

phosphorylation of PKR and eIF2 α was also not observed in MHV A59-infected cells [71]. However, Bechill et al have detected significant eIF2 α phosphorylation and up-regulation of ATF4 in cells infected with MHV A59, although no induction of GADD153 and GADD34 was observed [72]. It has been suggested that due to the lack of GADD34-mediated eIF2 α dephosphorylation, MHV infection induces sustained translation repression of most cellular proteins [72]. However, the translation of MHV mRNAs seems to be resistant to eIF2 α phosphorylation, and the detailed mechanisms for such evasion are yet to be investigated. As for SARS-CoV, PKR, PERK and eIF2 α phosphorylation are readily detectable in virus infected cells [73]. However, knock-down of PKR using specific morpholino oligomers did not affect SARS-CoV-induced eIF2 α phosphorylation but significantly inhibited SARS-CoV-induced apoptosis [73]. It is possible that eIF2 α is phosphorylated by PERK in SARS-CoV-infected cells, but similar loss-of-function experiments have not been performed, although over-expression of SARS-CoV accessory protein 3a has been shown to activate the PERK pathway [74].

The discrepancy regarding the activation of PKR/PERK during coronavirus infection may be a result from the different cell culture systems and virus strains used. The interpretation is further complicated by the interferon-inducible nature of PKR. It is generally believed that coronaviruses are poor type I interferon inducers *in vitro* [75,76,77], although the interferon response may be essential for antiviral activities *in vivo* [78]. Moreover, it is known that coronaviruses employ multiple mechanisms to antagonize the interferon response. For example, the nsp16 has been shown to utilize the 2'-O-methyltransferase activity to modify coronavirus mRNAs, so as to evade from the cytosolic RNA sensor melanoma differentiation-associated protein 5 (MDA5) and type I IFN induction [79,80]. Furthermore, the activities of several interferon induced genes (ISGs) have also been shown to be modulated by coronaviruses during infection. For instance, Zhao et al have demonstrated that the MHV accessory protein ns2 cleaves 2',5'-oligoadenylate, the product of an ISG called OAS [81]. This results in the suppression of the cellular endoribonuclease RNase L activity and facilitates virus replication *in vitro* and *in vivo* [81,82]. Thus, similar uncharacterized mechanisms may be used by MHV and other coronaviruses to block the activation and/or downstream signaling of PKR. In this regard, the activation of PERK via ER stress seems to be an alternative pathway to activate eIF2 α , although coronaviruses may counteract by directly targeting eIF2 α , as described below.

Studies done by this group have shown that, phosphorylation of PKR, PERK and eIF2 α were detectable at early stage of IBV infection (0-8 hpi) but diminished quickly

afterwards [83,84]. The rapid de-phosphorylation of eIF2 α is likely due to the accumulation of GADD34, which is a component of the PP1 complex and a downstream target gene induced by GADD153 [84]. Despite of the rapid de-phosphorylation of eIF2 α , significant induction of GADD153 was observed at late stage of infection (16-24h) at both mRNA and protein levels [83]. The up-regulation of GADD153 was likely mediated by both PKR and PERK, since knock-down of either PKR or PERK by siRNA reduces IBV-induced GADD153 [83]. The up-regulation of GADD153 promotes apoptosis in IBV-infected cells, possibly via inducing the pro-apoptotic protein tribbles-related protein 3 (TRIB3) and suppressing the pro-survival kinase extracellular signal-related kinase (ERK) [83]. Based on the findings so far obtained, it is safe to conclude that the PERK/PKR-eIF2 α -ATF4-GADD153 pathway is activated by some, but not all, coronaviruses. In the infected cells, this pathway is activated at an early stage but quickly modulated by feedback de-phosphorylation. The PERK/PKR-eIF2 α -ATF4-GADD153 most likely plays a pro-apoptotic function during coronavirus infection.

Integrated stress response pathways and innate immunity. Several recent studies have demonstrated the critical roles of cellular stress response pathways in modulating the innate immune activation [85]. One of the key regulators that bridge stress and innate immunity is GADD34, a negative regulator of eIF2 α activation. It has been shown that when stimulated with polyriboinosinic:polyribocytidylic acid (polyI:C), the integrated stress response pathways were activated in dendritic cells (DCs), leading to up-regulation of ATF4 and GADD34 [86]. Interestingly, GADD34 expression did not significantly affect protein synthesis in DCs, but was shown to be crucial for the production of interferon β (IFN- β) and pro-inflammatory cytokines interleukin-6 (IL-6) [86]. In contrast, GADD34 has also been shown to specify PP1 to dephosphorylate the TGF- β -activated kinase 1 (TAK1), thus negatively regulating the Toll-like receptor (TLR) signaling and pro-inflammatory cytokines [IL-6 and TNF- α (Tumor necrosis factor alpha)] production in macrophages [87]. The functional disparities of GADD34 in DCs and macrophages indicate that the integrated stress response may be regulated by some other signaling pathways, resulting in cell-type specific outcomes in the innate immune activation. Since GADD34 induction was readily observed in cells infected with IBV [84], it will be intriguing to ask whether GADD34 also contributes to IBV-induced pro-inflammatory cytokines production, and to determine potential cross-talks between the PERK pathway and innate immune activation during IBV infection.

The massive production of pro-inflammatory cytokines (cytokine storm) has been associated with the immunopathogenesis and high mortality rate of SARS-CoV [88]. The

transcription factor nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B) is a master regulator of pro-inflammatory response and innate immunity [89]. It has been well established that NF- κ B is required for the induction of pro-inflammatory cytokines (such as IL-6 and IL-8) and the early expression of IFN- β during RNA virus infection [90,91,92,93,94]. Interestingly, induction of TNF- α , IL-6 and IL-8 has been detected in cells over-expressing the spike protein of SARS-CoV via the NF- κ B pathway [95,96]. Thus it is intriguing to consider the involvement of ER stress in activating the NF- κ B pathway during coronavirus infection. In its inactive form, NF- κ B is sequestered in the cytoplasm by inhibitor of NF- κ B alpha ($\text{I}\kappa\text{B}\alpha$), which masks the nuclear localization signal of NF- κ B [97]. The basal level of $\text{I}\kappa\text{B}\alpha$ is maintained by constitutive synthesis and degradation of the protein [98]. Under various stress conditions, phosphorylation of eIF2 α leads to global translation repression and a net decrease in $\text{I}\kappa\text{B}\alpha$ protein level [99]. This then results in the activation of NF- κ B and induction of pro-inflammatory response (Figure 3). Nonetheless, further studies are needed to characterize the actual contributions of ER stress in NF- κ B mediated cytokine induction during coronavirus infection.

Previous study done by this group has shown that infection of IBV induced the production of IL-6 and IL-8, which was dependent on the phosphorylation of MAP kinase p38 [100]. Interestingly, a protein phosphatase called dual-specificity phosphatase 1 (DUSP1) was also up-regulated in IBV-infected cells and dephosphorylated p38 to modulate pro-inflammatory cytokine production [100]. Previous studies have shown that the mRNA and protein levels of DUSP1 are modulated by ER stress [101,102]. ER stress-induced DUSP1 up-regulation is likely to be mediated by ATF3 in the PERK pathway, since knock-down of ATF3 significantly reduced DUSP1 induction in cells under ER stress [103]. Thus, it is possible that IBV infection activates the PERK branch of UPR to induce DUSP1 expression, which in turn dephosphorylates p38 to modulate IBV-induced pro-inflammatory cytokine production (Figure 3).

Besides p38, DUSP1 has also been shown to dephosphorylate c-Jun N-terminal kinase (JNK) and ERK [104,105]. It has been long proposed that ERK phosphorylation promotes cell survival, whereas prolonged JNK and p38 phosphorylation is linked to the induction of apoptosis [106]. Thus, the induction of DUSP1 by ER stress in coronavirus-infected cells may also contribute to virus-induced apoptosis via modulation of the MAP kinase pathways.

The IRE1 branch of UPR

IRE1-XBP1 Signaling pathway. The IRE1-XBP1 branch of the UPR is evolutionarily conserved from yeast to humans. In response to unfolded proteins, IRE1 undergoes oligomerization [107]. This results in trans-autophosphorylation of the kinase domain and the activation of IRE1's RNase domain. So far, the only known substrate for IRE1 RNase activity is the mRNA of the X box binding protein 1 (XBP1) gene [108,109]. IRE1 cuts the XBP1 mRNA twice, removing a 26-nucleotide intron to form a frameshifted transcript, the spliced XBP1 (XBP1s). Whereas the unspliced XBP1 mRNA (XBP1u) encodes an inhibitor of the UPR, XBP1s encodes a potent transcriptional activator, which translocates to the nucleus and enhances the expression of many UPR genes, including those encoding molecular chaperones and proteins contributing to ER-associated degradation [110,111] (Figure 4).

Apart from the XBP1 pathway, activated IRE1 has been shown to recruit TNF receptor-associated factor 2 (TRAF2) and induce apoptosis by activating the JNK [112]. This IRE1-JNK pathway is independent of IRE1's RNase activity, but requires IRE1's kinase domain and involves TRAF2-dependent activation of caspase-12 [113]. Moreover, one recent study has demonstrated that the IRE1-JNK pathway is required for autophagy activation after pharmacological induction of ER stress. It was found that the kinase domain but not the RNase activity of IRE1 was required, and treatment of a JNK inhibitor (SP600125) abolished autophagosome formation after ER stress [114]. Therefore the IRE1 branch of UPR is closely associated with the JNK pathway and involved in JNK-mediated apoptosis and autophagy signaling.

Activation of the IRE1 pathway during coronaviruses infection. The involvement of IRE1-XBP1 pathway during coronavirus infection has been investigated by several studies, using MHV as a model. Either MHV infection or overexpression of the MHV S protein (but not other structural proteins) induces XBP1 mRNA splicing [37,72]. However, although XBP1 mRNA is efficiently spliced, the protein product of spliced XBP1 cannot be detected in either the whole cell lysate or the nuclear fraction. Moreover, UPR downstream genes known to be activated by XBP1s, such as **endoplasmic reticulum DNA J domain-containing protein 4 (ERdj4)**, **EDEM1** and **protein kinase inhibitor of 58 kDa (p58^{IPK})**, are not significantly induced after infection [72]. Using a luciferase reporter system, it is shown that MHV infection does not inhibit transactivation of **unfolded protein response element (UPRE)** and **ER stress response element (ERSE)** promoter by XBP1s. Because MHV infection is associated with persistent eIF2 α phosphorylation and host translational repression, it is likely that failure to translate the XBP1s protein may be the main reason why activation of the IRE1

branch does not occur even though XBP1 mRNA splicing is observed. On the other hand, although SARS-CoV belongs to the same genera of Betacoronavirus as MHV, neither infection with SARS-CoV nor overexpression of SARS-CoV S protein induces XBP1 mRNA splicing [37,115]. It is possible that other viral proteins of SARS-CoV (such as the E protein mentioned below), function as an antagonist of IRE1-XBP1 activation.

Result from this group has also shown that the IRE1-XBP1 pathway is activated in cells infected with IBV. In IBV-infected Vero cells, significant splicing of XBP1 mRNA was detected starting from 12-16 hours post infection till the late stage of infection. The mRNA levels of XBP1 effector genes (EDE1, ERdj4 and p58^{IPK}) were up-regulated in IBV-infected Vero cells. The activation of IRE1-XBP1 pathway was also detectable, though at a lower level, in other cell lines such as H1299 and Huh-7 cells. Treatment of IRE1 inhibitor effectively blocked IBV-induced XBP1 mRNA splicing and effector genes up-regulation in a dosage-dependent manner. Consistently, knockdown of IRE1 inhibited IBV-induced XBP1 mRNA splicing, whereas overexpression of wild type IRE1 (but not its kinase dead or RNase domain deleted mutants) enhanced IBV-induced XBP1 mRNA splicing. These results suggest that the IRE1-XBP1 pathway is indeed activated in cells infected with IBV. Interestingly, an earlier onset and more significant apoptosis induction in IRE1-knockdown IBV-infected cells was observed, which is associated with hyper-phosphorylation of pro-apoptotic kinase JNK and hypo-phosphorylation of pro-survival kinase **RAC-alpha serine/threonine-protein kinase (Akt)**. Taken together, IRE1 may modulate IBV-induced apoptosis and serve as a survival factor during coronavirus infection.

Interestingly, a recent report by DeDiego et al demonstrates that the coronavirus E protein may modulate the IRE1-XBP1 pathway. Using a recombinant SARS-CoV that lacks the E protein (rSARS-CoV-ΔE), it is found that both XBP1 splicing and induction of UPR genes significantly increase in the absence of E protein. Moreover, E protein also suppresses ER stress induced by RSV and drugs (thapsigargin and tunicamycin) [115]. Whether the UPR modulating activity is related to the viroporin property of E protein remains to be investigated, but this study explains, at least in part, why SARS-CoV lacking the E protein is attenuated in animal models [116,117].

IRE1-dependent decay during virus infection. Notably, one recent study has demonstrated an alternative function of IRE1. It was found that at the late stage of ER stress, IRE1 mediates non-specific cleavage of membrane associated mRNA species. This was dubbed IRE1-dependent decay (RIDD) and was proposed to resolve ER stress by reducing the amount of transcripts influx [118]. It is intriguing to think of RIDD as a host anti-viral mechanism.

During prolonged ER stress induced by infection, non-specific RNase activity of IRE1 may decay the membrane associated viral mRNA. In fact, it has been recently suggested that RIDD is activated during Japanese encephalitis virus (JEV) infection in Neuro2a cells [119]. Interestingly, RIDD specifically degraded known target mRNA transcripts but not JEV RNAs. Also, treatment with IRE1 RNase activity inhibitor suppressed viral replication, indicating that JEV benefits from RIDD activation [119].

IRE1 pathway and innate immunity. Similar to the integrated stress response, the IRE1 pathway has also been implicated in the innate immune response [85]. Martinon et al have shown that in murine macrophages, the IRE1-XBP1 pathway is specifically activated by TLR4 and TLR2 [120]. Interestingly, the ER stress and TLR activation synergistically activate IRE1 and induce the production of pro-inflammatory cytokines such as IL-1 β and IL-6 [120]. Consistently, Hu et al have demonstrated that the IRE1-XBP1 pathway is also involved in IFN- β and pro-inflammatory cytokines production in murine DCs induced by polyI:C [121]. Significantly, it has been shown that overexpression of the spliced form of XBP1 enhanced IFN- β production in DCs and significantly suppressed vesicular stomatitis virus infection [121]. Preliminary results from this group have also found that the activation of IRE1-XBP1 pathway is required for IL-8 induction in cells infected with IBV (unpublished data). **On the other hand, the kinase but not the RNase activity of IRE1 has been associated with ER stress induced NF- κ B activation [122]. Under ER stress, IRE1 has been shown to phosphorylate TRAF2, which activates the I κ B kinase (IKK) and contributes to its basal activity (Figure 4). IKK in turns phosphorylates I κ B α and promotes its proteasomal degradation, releasing NF- κ B to activate downstream genes [122].** Taken together, these findings suggest that IRE1 may act synergistically with players in innate immunity and serve as a supplementary sensor and/or signaling factors during coronavirus infection.

The ATF6 branch of UPR

The ER stress sensor ATF6 has an N-terminal cytoplasmic domain, a single transmembrane segment and an ER luminal domain that sense the presence of unfolded/misfolded proteins. Under ER stress, ATF6 is translocated from the ER to the Golgi apparatus and cleaved by protease S1P and S2P [123]. The cleavage releases the cytosolic basic leucine zipper (bZIP) domain, which translocates into the nucleus and activates genes harboring the ERSE or ERSE II [124]. The identified targets genes of ATF6 include ER chaperones (such as GRP78, GRP94), protein disulfide isomerase and the UPR transcription factors GADD153 and XBP1 [56]. Previously, it was proposed the ATF6 pathway is mainly

pro-survival, as it enhances the ER protein folding capacity to counteract ER stress [56]. However, recent studies have demonstrated that, under certain circumstances, ATF6-mediated signals may also contribute to ER-stress induced apoptosis, possibly via activation of CHOP and/or suppression of myeloid cell leukemia sequence 1 (Mcl-1) [125,126,127].

The infection of cells by several viruses has been shown to activate the ATF6 pathway, including the Tick-borne encephalitic virus, African swine fever virus (ASFV), West Nile virus (WNV) and HCV [128,129,130,131]. In the case of ASFV, ATF6 activation has been shown to modulate ASFV-induced apoptosis and facilitate viral replication [129]. For WNV, it has been shown that ATF6 activation promotes efficient WNV replication by suppressing signal transducer and activator of transcription 1 (STAT1) phosphorylation and late-phase interferon signaling [132]. The NS4B protein of HCV has been shown to activate ATF6 signaling in cultured cells [133]. Induction of chronic ER stress and adaptation of infected hepatocyte to UPR have been considered important for HCV persistent infection and pathogenesis *in vivo* [131,134].

Compared with the PERK and IRE1 pathway, the induction of ATF6 pathway during coronaviruses infection has not been deeply investigated. In MHV-infected cells, significant cleavage of ATF6 could be detected starting from 7 hours post infection [72]. However, the levels of both full length and cleaved ATF6 protein diminished at later time points during infection. Moreover, activation of ATF6 target genes was not observed at the mRNA level, as determined by luciferase reporter constructs under the control of ERSE promoters [72]. It is also unlikely that MHV infection suppresses downstream signaling of the ATF6 pathway, because the reporter induction by overexpressed ATF6 was not inhibited by MHV infection. The authors thus conclude that global translation shutdown via eIF2 α phosphorylation prevent accumulation of ATF6 and activation of ATF6 target genes [72]. The involvement of ATF6 pathway during infection of other coronaviruses has not been well characterized.

Although the spike proteins of coronaviruses have been considered the major contributor in ER stress induction, overexpression of SARS-CoV spike protein fails to activate ATF6 reporter constructs [36]. On the other hand, the accessory protein 8ab of SARS-CoV has been identified to induce ATF6 activation [135]. The 8ab protein was found in SARS-CoV isolated from animals and early human isolates. In SARS-CoV isolated from humans during the peak of the epidemic, there is a 29-nt deletion in the middle of ORF8, resulting in the splitting of ORF8 into two smaller ORFs, namely ORF8a and ORF8b, which encode two truncated polypeptides 8a and 8b [136]. ATF6 cleavage and nuclear translocation was observed in cells transfected with SARS-CoV 8ab [135]. Physical interaction between

8ab and the luminal domain of ATF6 was also demonstrated by co-immunoprecipitation. However, similar experiments have not been performed for the 8a and 8b proteins. Also, further studies using recombinant SARS-CoV lacking 8a, 8b or 8ab would be required.

Conclusion

Coronaviruses constitute human and animal pathogens that are medically and economically important. Much remains unknown regarding the host virus interactions during infection. Recent studies have demonstrated that coronaviruses infection induces ER stress in infected cells and activates the UPR. Activation of the PERK pathway (possibly in synergy with PKR and/or other integrated stress response kinases) leads to phosphorylation of eIF2 α and a global translation shutdown. At late stage of infection, up-regulation of transcription factor GADD153 likely contributes to coronaviruses induced apoptosis. Activation of the IRE1 pathway induces XBP1 mRNA splicing and expression of downstream UPR genes. Interestingly, IRE1 but not XBP1 is also shown to modulate the JNK and Akt kinase activities, thus protecting infected cells from virus induced apoptosis. The ATF6 pathway is also activated in coronaviruses infected cells, resulting in the up-regulation of chaperon proteins to counteract ER stress.

However, many questions remain to be addressed. First, although the coronaviruses spike proteins are demonstrated to induce ER stress and UPR, detailed mechanisms regarding molecular interactions between the spike proteins and PERK/IRE1/ATF6 have not been determined. Secondly, it should be noted that the phenotypes observed in cells overexpressing viral proteins may not essentially reflect their physiological functions in the setting of a real infection. Further experiments using recombinant viruses with deletion of or modification in the target viral proteins should be performed to validate these findings [115]. Last but not the least, the three branches of UPR should not be considered functionally independent, but rather as an integrated regulatory network [32]. For example, besides being spliced by IRE1, XBP1 is also transcriptionally activated by PERK and ATF6 [109,137]. Also, it is difficult to separate the translation shutdown effect mediated by PERK and the induction of UPR genes by PERK and the other two ER stress sensors, as in the studies with MHV [72].

Nonetheless, there are scientific and clinical significance for studies on ER stress and UPR induction during infection with coronaviruses and other viruses. As an evolutionarily conserved and well characterized stress response pathway, it serves as a perfect model to study host-virus interactions and pathogenesis. Moreover, besides apoptosis, UPR has been recently demonstrated to crosstalk with other major cellular signaling pathways, including

MAP kinases pathways, autophagy and innate immune responses [86,113,114,120,121]. Thus, further investigations on coronaviruses induced UPR may also help identifying new targets for antiviral agents and developing more effective vaccines against coronaviruses.

1. Cavanagh D (2007) Coronavirus avian infectious bronchitis virus. *Veterinary research* 38: 281-297.
2. Hamre D, Procknow JJ (1966) A new virus isolated from the human respiratory tract. *Experimental Biology and Medicine* 121: 190-193.
3. Kaye HS, Ong SB, Dowdle WR (1972) Detection of coronavirus 229E antibody by indirect hemagglutination. *Applied microbiology* 24: 703-707.
4. Ksiazek TG, Erdman D, Goldsmith CS, Zaki SR, Peret T, et al. (2003) A novel coronavirus associated with severe acute respiratory syndrome. *The New England journal of medicine* 348: 1953-1966.
5. Li W, Shi Z, Yu M, Ren W, Smith C, et al. (2005) Bats are natural reservoirs of SARS-like coronaviruses. *Science* 310: 676-679.
6. Wang L, Eaton B (2007) Bats, civets and the emergence of SARS. *Wildlife and Emerging Zoonotic Diseases: the Biology, Circumstances and Consequences of Cross-Species Transmission*: 325-344.
7. Ge X-Y, Li J-L, Yang X-L, Chmura AA, Zhu G, et al. (2013) Isolation and characterization of a bat SARS-like coronavirus that uses the ACE2 receptor. *Nature*.
8. de Groot RJ, Baker SC, Baric RS, Brown CS, Drosten C, et al. (2013) Middle East respiratory syndrome coronavirus (MERS-CoV): announcement of the Coronavirus Study Group. *Journal of virology* 87: 7790-7792.
9. Graham RL, Donaldson EF, Baric RS (2013) A decade after SARS: strategies for controlling emerging coronaviruses. *Nature Reviews Microbiology* 11: 836-848.
10. Annan A, Baldwin HJ, Corman VM, Klose SM, Owusu M, et al. (2013) Human betacoronavirus 2c EMC/2012-related viruses in bats, Ghana and Europe. *Emerging infectious diseases* 19: 456.
11. Hemida M, Perera R, Wang P, Alhammadi M, Siu L, et al. (2013) Middle East Respiratory Syndrome (MERS) coronavirus seroprevalence in domestic livestock in Saudi Arabia, 2010 to 2013. *Euro surveillance: bulletin Européen sur les maladies transmissibles= European communicable disease bulletin* 18.
12. Alagaili AN, Briese T, Mishra N, Kapoor V, Sameroff SC, et al. (2014) Middle East respiratory syndrome coronavirus infection in dromedary camels in Saudi Arabia. *MBio* 5: e00884-00814.
13. Masters PS (2006) The molecular biology of coronaviruses. *Advances in virus research* 66: 193-292.
14. Gorbalenya AE, Snijder EJ, Spaan WJ (2004) Severe acute respiratory syndrome coronavirus phylogeny: toward consensus. *Journal of virology* 78: 7863-7866.
15. Liu D, Inglis S (1991) Association of the infectious bronchitis virus 3c protein with the virion envelope. *Virology* 185: 911-917.

16. Sturman LS, Holmes K, Behnke J (1980) Isolation of coronavirus envelope glycoproteins and interaction with the viral nucleocapsid. *Journal of virology* 33: 449-462.
17. Qiu Z, Hingley ST, Simmons G, Yu C, Sarma JD, et al. (2006) Endosomal proteolysis by cathepsins is necessary for murine coronavirus mouse hepatitis virus type 2 spike-mediated entry. *Journal of virology* 80: 5768-5776.
18. Huang I-C, Bosch BJ, Li F, Li W, Lee KH, et al. (2006) SARS coronavirus, but not human coronavirus NL63, utilizes cathepsin L to infect ACE2-expressing cells. *Journal of Biological Chemistry* 281: 3198-3203.
19. Yamada Y, Liu XB, Fang SG, Tay FPL, Liu DX (2009) Acquisition of Cell–Cell Fusion Activity by Amino Acid Substitutions in Spike Protein Determines the Infectivity of a Coronavirus in Cultured Cells. *PloS one* 4: e6130.
20. Kuo L, Godeke GJ, Raamsman MJ, Masters PS, Rottier PJ (2000) Retargeting of coronavirus by substitution of the spike glycoprotein ectodomain: crossing the host cell species barrier. *Journal of virology* 74: 1393-1406.
21. Brierley I, Bournsnel ME, Binns MM, Bilimoria B, Blok VC, et al. (1987) An efficient ribosomal frame-shifting signal in the polymerase-encoding region of the coronavirus IBV. *The EMBO journal* 6: 3779-3785.
22. Lu Y, Lu X, Denison MR (1995) Identification and characterization of a serine-like proteinase of the murine coronavirus MHV-A59. *Journal of virology* 69: 3554-3559.
23. Baker S, Yokomori K, Dong S, Carlisle R, Gorbalenya A, et al. (1993) Identification of the catalytic sites of a papain-like cysteine proteinase of murine coronavirus. *Journal of virology* 67: 6056-6063.
24. Sawicki SG, Sawicki DL, Siddell SG (2007) A contemporary view of coronavirus transcription. *Journal of virology* 81: 20-29.
25. Oostra M, Te Lintelo E, Deijns M, Verheije M, Rottier P, et al. (2007) Localization and membrane topology of coronavirus nonstructural protein 4: involvement of the early secretory pathway in replication. *Journal of virology* 81: 12323-12336.
26. Knoops K, Kikkert M, Van Den Worm SHE, Zevenhoven-Dobbe JC, Van Der Meer Y, et al. (2008) SARS-coronavirus replication is supported by a reticulovesicular network of modified endoplasmic reticulum. *PLoS biology* 6: e226.
27. Gosert R, Kanjanahaluethai A, Egger D, Bienz K, Baker SC (2002) RNA replication of mouse hepatitis virus takes place at double-membrane vesicles. *Journal of virology* 76: 3697-3708.
28. Snijder EJ, van der Meer Y, Zevenhoven-Dobbe J, Onderwater JJ, van der Meulen J, et al. (2006) Ultrastructure and origin of membrane vesicles associated with the severe acute respiratory syndrome coronavirus replication complex. *Journal of virology* 80: 5927-5940.

29. Krijnse-Locker J, Ericsson M, Rottier P, Griffiths G (1994) Characterization of the budding compartment of mouse hepatitis virus: evidence that transport from the RER to the Golgi complex requires only one vesicular transport step. *The Journal of cell biology* 124: 55-70.
30. Pineau L, Colas J, Dupont S, Beney L, Fleurat - Lessard P, et al. (2009) Lipid - Induced ER Stress: Synergistic Effects of Sterols and Saturated Fatty Acids. *Traffic* 10: 673-690.
31. Kharroubi I, Ladrière L, Cardozo AK, Dogusan Z, Cnop M, et al. (2004) Free fatty acids and cytokines induce pancreatic β -cell apoptosis by different mechanisms: role of nuclear factor- κ B and endoplasmic reticulum stress. *Endocrinology* 145: 5087-5096.
32. Ron D, Walter P (2007) Signal integration in the endoplasmic reticulum unfolded protein response. *Nature reviews Molecular cell biology* 8: 519-529.
33. Tabas I, Ron D (2011) Integrating the mechanisms of apoptosis induced by endoplasmic reticulum stress. *Nature cell biology* 13: 184-190.
34. Jiang XS, Tang LY, Dai J, Zhou H, Li SJ, et al. (2005) Quantitative analysis of severe acute respiratory syndrome (SARS)-associated coronavirus-infected cells using proteomic approaches. *Molecular & Cellular Proteomics* 4: 902-913.
35. Yeung YS, Yip CW, Hon CC, Chow KYC, Ma I, et al. (2008) Transcriptional profiling of Vero E6 cells over-expressing SARS-CoV S2 subunit: Insights on viral regulation of apoptosis and proliferation. *Virology* 371: 32-43.
36. Chan CP, Siu KL, Chin KT, Yuen KY, Zheng B, et al. (2006) Modulation of the unfolded protein response by the severe acute respiratory syndrome coronavirus spike protein. *Journal of virology* 80: 9279-9287.
37. Versteeg GA, Van De Nes PS, Bredenbeek PJ, Spaan WJM (2007) The coronavirus spike protein induces endoplasmic reticulum stress and upregulation of intracellular chemokine mRNA concentrations. *Journal of virology* 81: 10981-10990.
38. Miller S, Krijnse-Locker J (2008) Modification of intracellular membrane structures for virus replication. *Nature Reviews Microbiology* 6: 363-374.
39. David-Ferreira J, Manaker R (1965) An electron microscope study of the development of a mouse hepatitis virus in tissue culture cells. *The Journal of cell biology* 24: 57-78.
40. Angelini MM, Akhlaghpour M, Neuman BW, Buchmeier MJ (2013) Severe Acute Respiratory Syndrome Coronavirus Nonstructural Proteins 3, 4, and 6 Induce Double-Membrane Vesicles. *MBio* 4: e00524-00513.
41. van der Meer Y, Snijder EJ, Dobbe JC, Schleich S, Denison MR, et al. (1999) Localization of mouse hepatitis virus nonstructural proteins and RNA synthesis indicates a role for late endosomes in viral replication. *Journal of virology* 73: 7641-7657.

42. Prentice E, Jerome WG, Yoshimori T, Mizushima N, Denison MR (2004) Coronavirus replication complex formation utilizes components of cellular autophagy. *The Journal of biological chemistry* 279: 10136-10141.
43. Verheije MH, Raaben M, Mari M, Te Lintelo EG, Reggiori F, et al. (2008) Mouse hepatitis coronavirus RNA replication depends on GBF1-mediated ARF1 activation. *PLoS pathogens* 4: e1000088.
44. Snijder EJ, van der Meer Y, Zevenhoven-Dobbe J, Onderwater JJ, van der Meulen J, et al. (2006) Ultrastructure and origin of membrane vesicles associated with the severe acute respiratory syndrome coronavirus replication complex. *Journal of virology* 80: 5927-5940.
45. Reggiori F, Monastyrska I, Verheije MH, Cali T, Ulasli M, et al. (2010) Coronaviruses Hijack the LC3-I-positive EDEMosomes, ER-derived vesicles exporting short-lived ERAD regulators, for replication. *Cell host & microbe* 7: 500-508.
46. Cali T, Galli C, Olivari S, Molinari M (2008) Segregation and rapid turnover of EDEM1 by an autophagy-like mechanism modulates standard ERAD and folding activities. *Biochemical and biophysical research communications* 371: 405-410.
47. Cavanagh D, Davis PJ (1988) Evolution of avian coronavirus IBV: sequence of the matrix glycoprotein gene and intergenic region of several serotypes. *The Journal of general virology* 69: 621-629.
48. Jacobs L, van der Zeijst B, Horzinek M (1986) Characterization and translation of transmissible gastroenteritis virus mRNAs. *Journal of virology* 57: 1010-1015.
49. Nal B, Chan C, Kien F, Siu L, Tse J, et al. (2005) Differential maturation and subcellular localization of severe acute respiratory syndrome coronavirus surface proteins S, M and E. *Journal of general virology* 86: 1423-1434.
50. Charley B, Laude H (1988) Induction of alpha interferon by transmissible gastroenteritis coronavirus: role of transmembrane glycoprotein E1. *Journal of virology* 62: 8-11.
51. de Haan CA, de Wit M, Kuo L, Montalto-Morrison C, Haagmans BL, et al. (2003) The glycosylation status of the murine hepatitis coronavirus M protein affects the interferogenic capacity of the virus in vitro and its ability to replicate in the liver but not the brain. *Virology* 312: 395-406.
52. Laude H, Gelfi J, Lavenant L, Charley B (1992) Single amino acid changes in the viral glycoprotein M affect induction of alpha interferon by the coronavirus transmissible gastroenteritis virus. *Journal of virology* 66: 743-749.
53. Delmas B, Laude H (1990) Assembly of coronavirus spike protein into trimers and its role in epitope expression. *Journal of virology* 64: 5367-5375.
54. Han DP, Lohani M, Cho MW (2007) Specific asparagine-linked glycosylation sites are critical for DC-SIGN-and L-SIGN-mediated severe acute respiratory syndrome coronavirus entry. *Journal of virology* 81: 12029-12039.

55. Fukushi M, Yoshinaka Y, Matsuoka Y, Hatakeyama S, Ishizaka Y, et al. (2012) Monitoring S protein maturation in the endoplasmic reticulum by calnexin is important for the infectivity of severe acute respiratory syndrome-coronavirus. *Journal of virology*.
56. Szegezdi E, Logue SE, Gorman AM, Samali A (2006) Mediators of endoplasmic reticulum stress-induced apoptosis. *EMBO reports* 7: 880-885.
57. CLEMENS MJ, ELIA A (1997) The double-stranded RNA-dependent protein kinase PKR: structure and function. *Journal of interferon & cytokine research* 17: 503-524.
58. McEwen E, Kedersha N, Song B, Scheuner D, Gilks N, et al. (2005) Heme-regulated inhibitor kinase-mediated phosphorylation of eukaryotic translation initiation factor 2 inhibits translation, induces stress granule formation, and mediates survival upon arsenite exposure. *Journal of Biological Chemistry* 280: 16925-16933.
59. Sood R, Porter AC, Olsen DA, Cavener DR, Wek RC (2000) A mammalian homologue of GCN2 protein kinase important for translational control by phosphorylation of eukaryotic initiation factor-2 α . *Genetics* 154: 787-801.
60. Puthalakath H, O'Reilly LA, Gunn P, Lee L, Kelly PN, et al. (2007) ER stress triggers apoptosis by activating BH3-only protein Bim. *Cell* 129: 1337-1349.
61. Marciniak SJ, Yun CY, Oyadomari S, Novoa I, Zhang Y, et al. (2004) CHOP induces death by promoting protein synthesis and oxidation in the stressed endoplasmic reticulum. *Genes & development* 18: 3066.
62. He B (2006) Viruses, endoplasmic reticulum stress, and interferon responses. *Cell death and differentiation* 13: 393-403.
63. Gale Jr MJ, Korth MJ, Tang NM, Tan SL, Hopkins DA, et al. (1997) Evidence that hepatitis C virus resistance to interferon is mediated through repression of the PKR protein kinase by the nonstructural 5A protein. *Virology* 230: 217.
64. Lu Y, Wambach M, Katze MG, Krug RM (1995) Binding of the influenza virus NS1 protein to double-stranded RNA inhibits the activation of the protein kinase that phosphorylates the eIF-2 translation initiation factor. *Virology* 214: 222.
65. Baltzis D, Qu LK, Papadopoulou S, Blais JD, Bell JC, et al. (2004) Resistance to vesicular stomatitis virus infection requires a functional cross talk between the eukaryotic translation initiation factor 2 α kinases PERK and PKR. *Journal of virology* 78: 12747-12761.
66. Jordan R, Wang L, Graczyk TM, Block TM, Romano PR (2002) Replication of a cytopathic strain of bovine viral diarrhea virus activates PERK and induces endoplasmic reticulum stress-mediated apoptosis of MDBK cells. *Journal of virology* 76: 9588-9599.
67. Cheng G, Feng Z, He B (2005) Herpes simplex virus 1 infection activates the endoplasmic reticulum resident kinase PERK and mediates eIF-2 α dephosphorylation by the γ 134.5 protein. *Journal of virology* 79: 1379-1388.

68. Pavio N, Romano PR, Graczyk TM, Feinstone SM, Taylor DR (2003) Protein synthesis and endoplasmic reticulum stress can be modulated by the hepatitis C virus envelope protein E2 through the eukaryotic initiation factor 2 α kinase PERK. *Journal of virology* 77: 3578-3585.
69. Mulvey M, Arias C, Mohr I (2007) Maintenance of endoplasmic reticulum (ER) homeostasis in herpes simplex virus type 1-infected cells through the association of a viral glycoprotein with PERK, a cellular ER stress sensor. *Journal of virology* 81: 3377-3390.
70. Zorzitto J, Galligan CL, Ueng JJ, Fish EN (2006) Characterization of the antiviral effects of interferon- α against a SARS-like coronavirus infection in vitro. *Cell research* 16: 220-229.
71. Ye Y, Hauns K, Langland JO, Jacobs BL, Hogue BG (2007) Mouse hepatitis coronavirus A59 nucleocapsid protein is a type I interferon antagonist. *Journal of virology* 81: 2554-2563.
72. Bechill J, Chen Z, Brewer JW, Baker SC (2008) Coronavirus infection modulates the unfolded protein response and mediates sustained translational repression. *Journal of virology* 82: 4492-4501.
73. Krähling V, Stein DA, Spiegel M, Weber F, Mühlberger E (2009) Severe acute respiratory syndrome coronavirus triggers apoptosis via protein kinase R but is resistant to its antiviral activity. *Journal of virology* 83: 2298-2309.
74. Minakshi R, Padhan K, Rani M, Khan N, Ahmad F, et al. (2009) The SARS Coronavirus 3a protein causes endoplasmic reticulum stress and induces ligand-independent downregulation of the type 1 interferon receptor. *PloS one* 4: e8342.
75. Garlinghouse Jr L, Smith A, Holford T (1984) The biological relationship of mouse hepatitis virus (MHV) strains and interferon: in vitro induction and sensitivities. *Archives of virology* 82: 19-29.
76. Spiegel M, Pichlmair A, Martínez-Sobrido L, Cros J, García-Sastre A, et al. (2005) Inhibition of beta interferon induction by severe acute respiratory syndrome coronavirus suggests a two-step model for activation of interferon regulatory factor 3. *Journal of virology* 79: 2079-2086.
77. Roth-Cross JK, Martínez-Sobrido L, Scott EP, García-Sastre A, Weiss SR (2007) Inhibition of the alpha/beta interferon response by mouse hepatitis virus at multiple levels. *Journal of virology* 81: 7189-7199.
78. Ireland DD, Stohlman SA, Hinton DR, Atkinson R, Bergmann CC (2008) Type I interferons are essential in controlling neurotropic coronavirus infection irrespective of functional CD8 T cells. *Journal of virology* 82: 300-310.
79. Züst R, Cervantes-Barragan L, Habjan M, Maier R, Neuman BW, et al. (2011) Ribose 2 [prime]-O-methylation provides a molecular signature for the distinction of self and non-self mRNA dependent on the RNA sensor Mda5. *Nature immunology* 12: 137-143.

80. Roth-Cross JK, Bender SJ, Weiss SR (2008) Murine coronavirus mouse hepatitis virus is recognized by MDA5 and induces type I interferon in brain macrophages/microglia. *Journal of virology* 82: 9829-9838.
81. Zhao L, Jha BK, Wu A, Elliott R, Ziebuhr J, et al. (2012) Antagonism of the interferon-induced OAS-RNase L pathway by murine coronavirus ns2 protein is required for virus replication and liver pathology. *Cell host & microbe* 11: 607-616.
82. Zhao L, Rose KM, Elliott R, Van Rooijen N, Weiss SR (2011) Cell-type-specific type I interferon antagonism influences organ tropism of murine coronavirus. *Journal of virology* 85: 10058-10068.
83. Liao Y, Fung TS, Huang M, Fang SG, Zhong Y, et al. (2013) Up-regulation of CHOP/GADD153 during Coronavirus Infectious Bronchitis Virus Infection Modulates Apoptosis by Restricting Activation of the Extracellular Signal-Regulated Kinase Pathway. *Journal of virology*.
84. Wang X, Liao Y, Yap PL, Png KJ, Tam JP, et al. (2009) Inhibition of protein kinase R activation and upregulation of GADD34 expression play a synergistic role in facilitating coronavirus replication by maintaining de novo protein synthesis in virus-infected cells. *Journal of virology* 83: 12462-12472.
85. Cláudio N, Dalet A, Gatti E, Pierre P (2013) Mapping the crossroads of immune activation and cellular stress response pathways. *The EMBO journal* 32: 1214-1224.
86. Clavarino G, Cláudio N, Dalet A, Terawaki S, Couderc T, et al. (2012) Protein phosphatase 1 subunit Ppp1r15a/GADD34 regulates cytokine production in polyinosinic: polycytidylic acid-stimulated dendritic cells. *Proceedings of the National Academy of Sciences* 109: 3006-3011.
87. Gu M, Ouyang C, Lin W, Zhang T, Cao X, et al. (2014) Phosphatase Holoenzyme PP1/GADD34 Negatively Regulates TLR Response by Inhibiting TAK1 Serine 412 Phosphorylation. *The Journal of Immunology* 192: 2846-2856.
88. Perlman S, Dandekar AA (2005) Immunopathogenesis of coronavirus infections: implications for SARS. *Nature reviews Immunology* 5: 917-927.
89. Hayden MS, Ghosh S (2012) NF- κ B, the first quarter-century: remarkable progress and outstanding questions. *Genes & development* 26: 203-234.
90. Balachandran S, Beg AA (2011) Defining emerging roles for NF- κ B in antiviral responses: revisiting the interferon- β enhanceosome paradigm. *PLoS pathogens* 7: e1002165.
91. Basagoudanavar SH, Thapa RJ, Nogusa S, Wang J, Beg AA, et al. (2011) Distinct roles for the NF- κ B RelA subunit during antiviral innate immune responses. *Journal of virology* 85: 2599-2610.
92. Wang J, Basagoudanavar SH, Wang X, Hopewell E, Albrecht R, et al. (2010) NF- κ B RelA subunit is crucial for early IFN- β expression and resistance to RNA virus replication. *The Journal of Immunology* 185: 1720-1729.

93. Kunsch C, Rosen CA (1993) NF-kappa B subunit-specific regulation of the interleukin-8 promoter. *Molecular and cellular biology* 13: 6137-6146.
94. Libermann TA, Baltimore D (1990) Activation of interleukin-6 gene expression through the NF-kappa B transcription factor. *Molecular and cellular biology* 10: 2327-2334.
95. Wang W, Ye L, Ye L, Li B, Gao B, et al. (2007) Up-regulation of IL-6 and TNF- α induced by SARS-coronavirus spike protein in murine macrophages via NF- κ B pathway. *Virus research* 128: 1-8.
96. Dosch SF, Mahajan SD, Collins AR (2009) SARS coronavirus spike protein-induced innate immune response occurs via activation of the NF- κ B pathway in human monocyte macrophages in vitro. *Virus research* 142: 19-27.
97. Karin M, Ben-Neriah Y (2000) Phosphorylation meets ubiquitination: the control of NF- κ B activity. *Annual review of immunology* 18: 621-663.
98. Kanarek N, London N, Schueler-Furman O, Ben-Neriah Y (2010) Ubiquitination and Degradation of the Inhibitors of NF- κ B. *Cold Spring Harbor perspectives in biology* 2: a000166.
99. Jiang H-Y, Wek SA, McGrath BC, Scheuner D, Kaufman RJ, et al. (2003) Phosphorylation of the α subunit of eukaryotic initiation factor 2 is required for activation of NF- κ B in response to diverse cellular stresses. *Molecular and cellular biology* 23: 5651-5663.
100. Liao Y, Wang X, Huang M, Tam JP, Liu DX (2011) Regulation of the p38 mitogen-activated protein kinase and dual-specificity phosphatase 1 feedback loop modulates the induction of interleukin 6 and 8 in cells infected with coronavirus infectious bronchitis virus. *Virology* 420: 106-116.
101. Boutros T, Nantel A, Emadali A, Tzimas G, Conzen S, et al. (2008) The MAP Kinase Phosphatase - 1 MKP - 1/DUSP1 Is a Regulator of Human Liver Response to Transplantation. *American Journal of Transplantation* 8: 2558-2568.
102. Li B, Yi P, Zhang B, Xu C, Liu Q, et al. (2011) Differences in endoplasmic reticulum stress signalling kinetics determine cell survival outcome through activation of MKP-1. *Cellular signalling* 23: 35-45.
103. Gora S, Maouche S, Atout R, Wanherdrick K, Lambeau G, et al. (2010) Phospholipolyzed LDL induces an inflammatory response in endothelial cells through endoplasmic reticulum stress signaling. *The FASEB Journal* 24: 3284-3297.
104. Sun H, Charles CH, Lau LF, Tonks NK (1993) MKP-1 (3CH134), an immediate early gene product, is a dual specificity phosphatase that dephosphorylates MAP kinase in vivo. *Cell* 75: 487-493.
105. Franklin CC, Kraft AS (1997) Conditional expression of the mitogen-activated protein kinase (MAPK) phosphatase MKP-1 preferentially inhibits p38 MAPK and stress-activated protein kinase in U937 cells. *Journal of Biological Chemistry* 272: 16917-16923.

106. Xia Z, Dickens M, Raingeaud J, Davis RJ, Greenberg ME (1995) Opposing effects of ERK and JNK-p38 MAP kinases on apoptosis. *Science* 270: 1326-1331.
107. Bertolotti A, Zhang Y, Hendershot LM, Harding HP, Ron D (2000) Dynamic interaction of BiP and ER stress transducers in the unfolded-protein response. *Nature cell biology* 2: 326-332.
108. Calfon M, Zeng H, Urano F, Till JH, Hubbard SR, et al. (2002) IRE1 couples endoplasmic reticulum load to secretory capacity by processing the XBP-1 mRNA. *Nature* 415: 92-96.
109. Yoshida H, Matsui T, Yamamoto A, Okada T, Mori K (2001) XBP1 mRNA is induced by ATF6 and spliced by IRE1 in response to ER stress to produce a highly active transcription factor. *Cell* 107: 881-891.
110. Ng DT, Spear ED, Walter P (2000) The unfolded protein response regulates multiple aspects of secretory and membrane protein biogenesis and endoplasmic reticulum quality control. *The Journal of cell biology* 150: 77-88.
111. Lee AH, Iwakoshi NN, Glimcher LH (2003) XBP-1 regulates a subset of endoplasmic reticulum resident chaperone genes in the unfolded protein response. *Molecular and cellular biology* 23: 7448-7459.
112. Urano F, Wang X, Bertolotti A, Zhang Y, Chung P, et al. (2000) Coupling of stress in the ER to activation of JNK protein kinases by transmembrane protein kinase IRE1. *Science* 287: 664-666.
113. Yoneda T, Imaizumi K, Oono K, Yui D, Gomi F, et al. (2001) Activation of caspase-12, an endoplasmic reticulum (ER) resident caspase, through tumor necrosis factor receptor-associated factor 2-dependent mechanism in response to the ER stress. *The Journal of biological chemistry* 276: 13935-13940.
114. Ogata M, Hino S, Saito A, Morikawa K, Kondo S, et al. (2006) Autophagy is activated for cell survival after endoplasmic reticulum stress. *Molecular and cellular biology* 26: 9220-9231.
115. DeDiego ML, Nieto-Torres JL, Jiménez-Guardeño JM, Regla-Nava JA, Álvarez E, et al. (2011) Severe Acute Respiratory Syndrome Coronavirus Envelope Protein Regulates Cell Stress Response and Apoptosis. *PLoS pathogens* 7: e1002315.
116. Liao Y, Lescar J, Tam J, Liu D (2004) Expression of SARS-coronavirus envelope protein in *Escherichia coli* cells alters membrane permeability. *Biochemical and biophysical research communications* 325: 374-380.
117. DeDiego ML, Álvarez E, Almazán F, Rejas MT, Lamirande E, et al. (2007) A severe acute respiratory syndrome coronavirus that lacks the E gene is attenuated in vitro and in vivo. *Journal of virology* 81: 1701-1713.
118. Hollien J, Lin JH, Li H, Stevens N, Walter P, et al. (2009) Regulated Ire1-dependent decay of messenger RNAs in mammalian cells. *The Journal of cell biology* 186: 323-331.

119. Bhattacharyya S, Sen U, Vrati S (2014) Regulated IRE1-dependent decay pathway is activated during Japanese encephalitis virus-induced unfolded protein response and benefits viral replication. *Journal of General Virology* 95: 71-79.
120. Martinon F, Chen X, Lee A-H, Glimcher LH (2010) TLR activation of the transcription factor XBP1 regulates innate immune responses in macrophages. *Nature immunology* 11: 411-418.
121. Hu F, Yu X, Wang H, Zuo D, Guo C, et al. (2011) ER stress and its regulator X - box - binding protein - 1 enhance polyIC - induced innate immune response in dendritic cells. *European journal of immunology* 41: 1086-1097.
122. Tam AB, Mercado EL, Hoffmann A, Niwa M (2012) ER stress activates NF- κ B by integrating functions of basal IKK activity, IRE1 and PERK. *PloS one* 7: e45078.
123. Haze K, Yoshida H, Yanagi H, Yura T, Mori K (1999) Mammalian transcription factor ATF6 is synthesized as a transmembrane protein and activated by proteolysis in response to endoplasmic reticulum stress. *Molecular biology of the cell* 10: 3787-3799.
124. Yoshida H, Okada T, Haze K, Yanagi H, Yura T, et al. (2001) Endoplasmic reticulum stress-induced formation of transcription factor complex ERSF including NF-Y (CBF) and activating transcription factors 6 α and 6 β that activates the mammalian unfolded protein response. *Molecular and cellular biology* 21: 1239-1248.
125. Gotoh T, Oyadomari S, Mori K, Mori M (2002) Nitric oxide-induced apoptosis in RAW 264.7 macrophages is mediated by endoplasmic reticulum stress pathway involving ATF6 and CHOP. *Journal of Biological Chemistry* 277: 12343-12350.
126. Morishima N, Nakanishi K, Nakano A (2011) Activating transcription factor-6 (ATF6) mediates apoptosis with reduction of myeloid cell leukemia sequence 1 (Mcl-1) protein via induction of WW domain binding protein 1. *Journal of Biological Chemistry* 286: 35227-35235.
127. Nakanishi K, Sudo T, Morishima N (2005) Endoplasmic reticulum stress signaling transmitted by ATF6 mediates apoptosis during muscle development. *The Journal of cell biology* 169: 555-560.
128. Yu C, Achazi K, Niedrig M (2013) Tick-borne encephalitis virus triggers inositol-requiring enzyme 1 (IRE1) and transcription factor 6 (ATF6) pathways of unfolded protein response. *Virus research* 178: 471-477.
129. Galindo I, Hernaez B, Munoz-Moreno R, Cuesta-Geijo M, Dalmau-Mena I, et al. (2012) The ATF6 branch of unfolded protein response and apoptosis are activated to promote African swine fever virus infection. *Cell death & disease* 3: e341.
130. Ambrose RL, Mackenzie JM (2011) West Nile virus differentially modulates the unfolded protein response to facilitate replication and immune evasion. *Journal of virology* 85: 2723-2732.

131. Merquiol E, Uzi D, Mueller T, Goldenberg D, Nahmias Y, et al. (2011) HCV causes chronic endoplasmic reticulum stress leading to adaptation and interference with the unfolded protein response. *PloS one* 6: e24660.
132. Ambrose RL, Mackenzie JM (2013) ATF6 signaling is required for efficient West Nile virus replication by promoting cell survival and inhibition of innate immune responses. *Journal of virology* 87: 2206-2214.
133. Li S, Ye L, Yu X, Xu B, Li K, et al. (2009) Hepatitis C virus NS4B induces unfolded protein response and endoplasmic reticulum overload response-dependent NF- κ B activation. *Virology* 391: 257-264.
134. Asselah T, Bièche I, Mansouri A, Laurendeau I, Cazals - Hatem D, et al. (2010) In vivo hepatic endoplasmic reticulum stress in patients with chronic hepatitis C. *The Journal of pathology* 221: 264-274.
135. Sung SC, Chao CY, Jeng KS, Yang JY, Lai M (2009) The 8ab protein of SARS-CoV is a luminal ER membrane-associated protein and induces the activation of ATF6. *Virology* 387: 402-413.
136. Guan Y, Zheng B, He Y, Liu X, Zhuang Z, et al. (2003) Isolation and characterization of viruses related to the SARS coronavirus from animals in southern China. *Science* 302: 276-278.
137. Calfon M, Zeng H, Urano F, Till JH, Hubbard SR, et al. (2002) IRE1 couples endoplasmic reticulum load to secretory capacity by processing the XBP-1 mRNA. *Nature* 415: 92-96.

Figure legends

Figure 1

Schematic diagram showing the replication cycle of coronavirus and the stages in which ER stress may be induced during coronavirus infection. Infection starts with receptor binding and entry by membrane fusion. After uncoating, the genomic RNA is used as a template to synthesize progeny genomes and a nested set of subgenomic RNAs. The replication transcription centers are closely associated with double membrane vesicles, which are proposed to be adopted from the modified ER, possibly by the combined activities of non-structural proteins nsp3, nsp4 and nsp6. The S, E and M proteins are synthesized and anchored on the ER, whereas the N protein is translated in the cytosol. Assembly takes place in the ERGIC and mature virions are released via smooth-walled vesicles by exocytosis. The three stages that presumably induce ER stress are highlighted with numbered star signs, namely: 1. formation of double membrane vesicles, 2. massive production and modification of structural proteins and 3. depletion of ER membrane during budding.

Figure 2

Flowchart showing the induction of ER stress and its physiological outcomes during coronavirus infection. The integrated stress response pathways (including PERK) trigger translation shutdown and modulate apoptosis. The ATF6 pathway enhances the ER folding capacity, and the IRE1 pathway affects both ER folding and apoptosis induction. Pointed arrows indicate activation, and blunt-ended lines indicate inhibition. The dotted line suggests uncharacterized function of GCN2 and HRI during coronavirus infection.

Figure 3

Working model of PKR/PERK-eIF2 α -ATF4-GADD153 pathway activation during coronavirus infection, using IBV as an example. Phosphorylation of eIF2 α by PERK and PKR induces the expression of ATF4, ATF3, and GADD153. GADD153 exerts its pro-apoptotic activities via suppressing Bcl2 and ERKs by inducing TRIB3. The potential induction of DUSP1 by ATF3 may modulate phosphorylation of p38 and JNK, thus regulating IBV-induced apoptosis and cytokine production. The translation attenuation due to eIF2 α activation can also lead to reduced inhibition of I κ B α on NF- κ B, which in turn promote cytokine production. Pointed arrows indicate activation, and blunt-ended lines indicate inhibition. The question mark indicates hypothetical mechanism.

Figure 4

Working model of IRE1-XBP1 signaling pathway during coronavirus infection, using IBV as an example. IRE1 mediates XBP1 splicing, which up-regulates UPR target genes to restore ER stress, and the spliced XBP1 may also modulate the interferon and cytokine secretion. IRE1 activation modulates the phosphorylation of Akt and JNK, thus affecting IBV-induced apoptosis. IRE1 is also responsible for basal activity of IKK, which phosphorylates I κ B α to remove its inhibition on NF- κ B, thus facilitating the production of type I interferon and pro-inflammatory cytokines. Pointed arrows indicate activation, and blunt-ended lines indicate inhibition. The question mark indicates hypothetical mechanism.

Figure 1.TIF




Figure 1

Figure 2.TIF




Figure 2

Figure 3.TIF




Figure 3

Figure 4.TIF




Figure 4

Copyright of *Frontiers in Microbiology* is the property of Frontiers Media S.A. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.