


Accepted Manuscript

Transmission of SARS and MERS coronaviruses and influenza virus in healthcare settings: the possible role of dry surface contamination

J.A. Otter, C. Donskey, S. Yezli, S. Douthwaite, S.D. Goldenberg, D.J. Weber



PII: S0195-6701(15)00367-9

DOI: [10.1016/j.jhin.2015.08.027](https://doi.org/10.1016/j.jhin.2015.08.027)

Reference: YJHIN 4641

To appear in: *Journal of Hospital Infection*

Received Date: 24 July 2015

Accepted Date: 28 August 2015

Please cite this article as: Otter JA, Donskey C, Yezli S, Douthwaite S, Goldenberg SD, Weber DJ, Transmission of SARS and MERS coronaviruses and influenza virus in healthcare settings: the possible role of dry surface contamination, *Journal of Hospital Infection* (2015), doi: 10.1016/j.jhin.2015.08.027.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

J.A. Otter *et al.*

Review

Transmission of SARS and MERS coronaviruses and influenza virus in healthcare settings: the possible role of dry surface contamination[star]

J.A. Otter^{a,*}, C. Donskey^b, S. Yezli^c, S. Douthwaite^d, S.D. Goldenberg^d, D.J. Weber^e

^a*Imperial College Healthcare NHS Trust, London, UK*

^b*Cleveland Veterans Affairs Medical Center, Cleveland, OH, USA*

^c*Global Centre for Mass Gatherings Medicine, Riyadh, Saudi Arabia*

^d*Centre for Clinical Infection and Diagnostics Research (CIDR), Guy's and St Thomas NHS Foundation Trust & King's College London, UK*

^e*Division of Infectious Diseases, University of North Carolina, Chapel Hill, NC, USA*

[star]This work was presented in part at the Infection Prevention Society Conference, Glasgow, September 29th to October 1st, 2014.

*Corresponding author. Address: Centre for Clinical Infection and Diagnostics Research (CIDR), Department of Infectious Diseases, King's College London, and Guy's and St Thomas' Hospital NHS Foundation Trust, London SE1 9RT, UK. Tel.: +44 (0)207 188 7188. *E-mail address:* jonathan.otter@kcl.ac.uk (J.A. Otter).

SUMMARY

Viruses with pandemic potential including H1N1, H5N1, and H5N7 influenza viruses, and severe acute respiratory syndrome (SARS)/Middle East respiratory syndrome (MERS) coronaviruses (CoV) have emerged in recent years. SARS-CoV, MERS-CoV, and influenza virus can survive on surfaces for extended periods, sometimes up to months. Factors influencing the survival of these viruses on surfaces include: strain variation, titre, surface type, suspending medium, mode of deposition, temperature and relative humidity, and the method used to determine the viability of the virus. Environmental sampling has identified contamination in field-settings with SARS-CoV and influenza virus, although the frequent use of molecular detection methods may not necessarily represent the presence of viable virus. The importance of indirect contact transmission (involving contamination of inanimate surfaces) is uncertain compared with other transmission routes, principally direct contact transmission (independent of surface contamination), droplet, and airborne routes. However, influenza virus and SARS-CoV may be shed into the environment and be transferred from environmental surfaces to hands of patients and healthcare providers. Emerging data suggest that MERS-CoV also shares these properties. Once contaminated from the environment, hands can then initiate self-inoculation of mucous membranes of the nose, eyes or mouth.

Mathematical and animal models, and intervention studies suggest that contact transmission is the most important route in some scenarios. Infection prevention and control implications include the need for hand hygiene and personal protective equipment to minimize self-contamination and to protect against inoculation of mucosal surfaces and the respiratory tract, and enhanced surface cleaning and disinfection in healthcare settings.

Keywords:

Healthcare-associated infection

Influenza virus

MERS-CoV

SARS-CoV

Surface contamination

Transmission

Introduction

A number of viruses with pandemic potential have emerged in recent years. The 2002 emergence of severe acute respiratory syndrome coronavirus (SARS-CoV), 2009 pandemic of H1N1 influenza, continued circulation of influenza H5N1 and H5N7 strains, and the recent emergence of the Middle East respiratory syndrome coronavirus (MERS-CoV) illustrate the current threat of these viruses.¹⁻⁴

Despite fundamental differences in their structure and epidemiology, these pandemic viral threats share a number of important properties. They are zoonotic enveloped RNA respiratory viruses that rarely transmit between humans in their native form, but could mutate to allow more efficient human-to-human transmission. This was illustrated by the 2002–2003 SARS pandemic and the 2009 H1N1 influenza pandemic.^{3,4} Frequent and accepted transmission routes are ‘droplet transmission’, where droplets ($>5\ \mu\text{m}$ diameter, travelling $<1\ \text{m}$) containing viable viruses make contact with the nose, mouth, eyes, or upper respiratory tract, and ‘airborne transmission’, where droplet nuclei ($\leq 5\ \mu\text{m}$ diameter, which can travel $>1\ \text{m}$) are inhaled by susceptible individuals (Figure 1).⁵⁻⁸ The role of ‘direct contact transmission’ (not involving contaminated surfaces) and ‘indirect contact transmission’ (involving contaminated surfaces) in the spread of these viruses with pandemic potential has been controversial (Figure 1).⁶⁻⁸ However, several reviews and models have suggested that indirect contact transmission is the predominant transmission route for some respiratory viruses, including influenza, in some settings.⁷⁻⁹

Contaminated surfaces are an established route of transmission for important nosocomial pathogens including *Clostridium difficile*, methicillin-resistant *Staphylococcus aureus* (MRSA), vancomycin-resistant enterococci (VRE), *Acinetobacter baumannii* and

norovirus, which share the capacity to survive on surfaces for extended periods.^{10–12} There is a general perception that enveloped viruses, such as influenza and human coronaviruses including MERS-CoV and SARS-CoV, have a very limited capacity to survive on dry surfaces.^{13–15} However, several studies suggest that SARS-CoV, MERS-CoV and influenza virus have the capacity to survive on dry surfaces for a sufficient duration to facilitate onward transmission.^{16–18} SARS-CoV and surrogates, and influenza virus can also survive in environmental reservoirs such as water, on foods, and in sewage for extended periods.^{19–25} Here, we review the studies evaluating influenza and human coronavirus survival on dry surfaces, field investigations that have performed surface sampling for these viruses, and we consider the importance of contaminated surfaces in the transmission of these viruses.

Search strategy

PubMed searches without date or language restrictions were performed on November 22nd, 2014 using the following search terms: [coronavirus or influenza] survival surface OR fomite transmission OR surface contamination OR disinfection transmission. Studies evaluating contamination of any surface were included. A total of 254 articles were identified using these search terms (Appendix A). Articles were also identified by hand-searching of bibliographies and related articles on PubMed.

Survival on dry surfaces

Tables I and II summarize in-vitro studies evaluating the capacity of human coronaviruses (including SARS-CoV and MERS-CoV) and influenza to survive when inoculated on to dry surfaces. Important methodological differences include variation in the choice of virus species and strain, method used to detect virus, titre and volume applied, surface substrate, suspending medium, temperature and relative humidity (RH), and drying time. These differences mean that direct comparison of reported survival times between studies is often not meaningful. In some of the reviewed studies, these factors have been experimental variables, allowing comment on the influence of the method used to detect virus, species and strain, titre, substrate, suspending medium, and temperature/RH on drying time (Tables I and II).

Notwithstanding differences in methodology, some common themes emerge. Survival times for SARS-CoV, MERS-CoV, and surrogates such as transmissible gastroenteritis virus (TGEV) are generally measured in days, weeks, or months.^{16,26,28–30,43} Survival times for influenza virus are generally shorter, often measured in hours rather than days.^{16,32–34} However, some studies have reported considerably longer survival times for influenza virus, measured in days rather than hours.^{35,36,39,40–42} This apparent conflict is most likely explained by experimental factors. The difference in survival capacity between influenza virus and that

of SARS-CoV and MERS-CoV is best illustrated by van Doremalen *et al.* who tested both H1N1 influenza and MERS-CoV.¹⁶ Viable MERS-CoV was recovered after 48 h, with a half-life ranging from ~0.5 to 1 h. By contrast, no viable H1N1 was recovered after 1 h under any of the conditions tested.

SARS-CoV and MERS-CoV appear to have an unusual capacity to survive on dry surfaces compared with other human coronaviruses (229E, OC43, and NL63).^{17,28,27,31,44} SARS-CoV, like the non-enveloped adenovirus comparator, survived for more than six days when dried on to Petri dishes compared with human coronavirus HCoV-229E, which survived for less than 72 h.²⁸ Although data are limited, it appears that MERS-CoV may survive on surfaces for longer than most human coronaviruses.¹⁶ Since other human coronaviruses do not share the unusual survival properties of SARS-CoV, TGEV and mouse hepatitis virus (MHV) are often used as surrogates.^{26,43,45}

No study has tested more than one strain of SARS-CoV or MERS-CoV. However, some studies have tested more than one strain of influenza, highlighting considerable strain variation.^{18,35,39,42} Further work is necessary to evaluate the importance of strain variation in influenza and coronavirus survival.

There appears to be a 'dose response' in terms of survival, with more concentrated viral suspensions surviving longer than less concentrated suspensions.^{29,33,39} For example, SARS-CoV survived on disposable gowns for 1 h at 10^4 TCID₅₀/mL vs 2 days at 10^6 TCID₅₀/mL.²⁹ Similarly, H3N2 influenza survived on bank notes for 1 h at 1.1×10^5 TCID₅₀/mL vs 2 days at 8.9×10^5 TCID₅₀/mL.³⁹

Substantial variation in survival times is evident for coronaviruses and influenza on different surface substrates.^{30,34,37,41} Coronaviruses and influenza both have the capacity to survive on a wide range of porous and non-porous materials, including metals, plastics (such as light switches, telephones, perspex, latex, rubber, and polystyrene), woven and non-woven fabrics (including cotton, polyester, handkerchiefs, and disposable tissues), paper (including magazine pages), wood, glass, stethoscopes, tissue, Formica®, bank notes, tiles, eggs, feathers, and soft toys.^{16,27,31,32,33,34,39,41,43} The properties of different surfaces are likely to influence survival times. For example, the survival of influenza dried on to copper surfaces was considerably shorter than on stainless steel.⁴⁰

Several studies have evaluated the capacity for SARS-CoV (and the surrogate TGEV), and influenza virus to survive on materials widely used as personal protective equipment (PPE) such as gowns, gloves, and respirators.^{29,37,43} For example, TGEV survived on isolation gowns, nitrile, and latex gloves, N95 respirators and scrubs with a $<10^2$ reduction for >4 h, and was detected on some items after 24 h.⁴³ One study showed that H1N1 influenza virus

dried on to various materials could be transferred to the hands of volunteers for at least 24 h following inoculation on some surfaces, with clear implications for the acquisition of viable viruses on the hands of healthcare personnel during the removal of PPE.⁴² A more recent study identified viable pandemic H1N1 influenza after six days on coupons made from N95 respirators.¹⁸

The suspending medium used to dry the viruses on to surfaces is another important factor influencing survival times.^{18,28,39,46} For example, adding mucus increased the survival time of influenza dried on bank notes from hours to up to 17 days.³⁹ A related variable is the mode of deposition of the viruses. Most studies dried a small volume of a known concentration of virus in a cell culture medium. However, several studies have evaluated the use of deposited virus from clinical specimens, which may be more representative of the clinical scenario and tends to result in shorter survival times.^{32,33,39}

In all studies that tested varying temperature and RH, lower temperature and RH favoured the survival of both coronaviruses and influenza.^{16-18,26,35,36,38}

Different methods have been applied to detect virus – most often cell culture assays but also RNA detection using polymerase chain reaction (PCR) or indirect methods such as fluorescence or haemagglutinin assays.^{27,33,34,37,40} Intact viral RNA appears to remain detectable on surfaces for longer than viruses that retain the ability to infect cells.^{32,33,42} Since PCR assays only detect a small portion of RNA they cannot be used to replace culture-based methods in determining viability.

Experimental factors that have been shown to influence virus viability *in vitro* are likely to have important implications for virus survival on hospital surfaces. For example, the titre and volume of virus applied to surfaces will be influenced by the type and volume of respiratory secretion, as will the suspending medium. The temperature and RH of the hospital environment is likely to be controlled to comfortable levels, meaning that some of the extremes of temperature and relative humidity tested *in vitro* may not be so relevant in the field.

Survival in aerosols

Respiratory virus symptoms such as sneezing and coughing result in the generation of virus-containing particles, in a size continuum from 1 to 500 μm .^{47,48} Whereas the generation of small droplet nuclei has traditionally been associated with ‘aerosol-generating procedures’, several recent studies have identified aerosols (droplet nuclei, $\leq 5 \mu\text{m}$ diameter) in the vicinity of patients infected with influenza who are not undergoing recognized aerosol-generating procedures.⁴⁹⁻⁵¹ Coronaviruses especially have the ability to survive for long periods in aerosols. For example, HCoV-229E aerosol remained infectious for six days at 20°C and 50%

RH.⁵² One study has evaluated the survival of MERS-CoV aerosols, finding a 7% reduction over 10 min (at 40% RH).¹⁶ By contrast, H1N1 suffered a 95% reduction over the same time period, suggesting that influenza virus may be less robust as an aerosol than coronaviruses. However, other studies have shown extended survival times for influenza aerosols (surviving up to 36 h).^{53–55}

Environmental contamination in field settings

A number of studies have performed environmental sampling for influenza or SARS in field settings (Table III). No studies have yet been published evaluating MERS-CoV contamination in field settings.

The major limitation with field studies is the use of PCR to detect viral RNA, which is best seen as a marker of virus shedding rather than indicating the presence of viable virus on surfaces, which must be confirmed by the recovery of viruses able to infect cells. In a number of influenza virus studies, a considerably lower rate of detection was identified by viral culture than by PCR, and in one study no viable virus was detected by culture despite the detection of influenza virus RNA.^{56–58} Similarly, regarding SARS, two studies have detected environmental reservoirs of SARS-CoV RNA by PCR, but no viable virus by culture.^{44,63}

Three studies have evaluated influenza contamination of surfaces in healthcare settings. A UK study detected influenza virus RNA on two (0.5%) of 397 samples from surfaces around infected individuals, one of which grew viable influenza.⁵⁷ More than half of the patients in the study were receiving antiviral medication, which may have reduced shedding. Influenza virus RNA was recovered from 38.5% of 13 environmental surfaces around hospitalized patients in Mexico.⁶¹ In one case, one out of five surfaces (a bed rail) was positive from a patient's room 72 h after patient discharge and terminal cleaning. Pappas *et al.* sampled toys in the waiting room of a general paediatric practice, finding that only one out of 59 toys was contaminated with influenza RNA.⁵⁹ However, a higher proportion of toys was contaminated with picornavirus RNA (19.2%), including four out of 15 after cleaning. The identification of viral DNA on surfaces after cleaning and disinfection may be a marker of ineffective cleaning and disinfection.

Several studies have evaluated influenza RNA or viable influenza in homes, day-care centres and elementary schools.^{58,60,62} The proportion of sites contaminated with influenza virus RNA varied from 3% to >50% in these studies, with evidence of seasonal variation in the study by Boone *et al.*⁶² In Bangkok, households randomized to a handwashing intervention had a lower proportion of sites contaminated with influenza virus RNA than did control households (11.1% of 45 vs 24.4% of 45).⁵⁸

Influenza RNA was detected on 15% of the 1862 environmental samples collected from bird markets in Indonesia, and almost half of the markets (47%) were contaminated at one or more site(s).⁵⁶ Viable influenza was cultured from 4.6% of 280 samples tested. Markets that slaughtered birds, as well as one particular province, were associated with contamination, whereas zoning of poultry activities and daily disposal of solid waste were protective.

Two studies have evaluated SARS-CoV contamination. A study of areas used to care for patients with SARS in Bangkok and Taipei found that 38.1% of 63 sites were contaminated with SARS-CoV RNA.⁴⁴ Furthermore, 6.4% of 31 public areas were also contaminated with SARS-CoV RNA. A lower rate of contamination was identified at a Canadian hospital, where 3.5% of 85 surfaces in SARS units were contaminated with SARS-CoV RNA.⁶³ Viral culture did not detect viable SARS-CoV from any of the surfaces in these studies. A study of public surfaces in Jeddah Airport, Saudi Arabia, identified human coronavirus RNA from three (7.5%) of 40 surface samples. No viral culture was performed in the study.⁶⁴

Importance of contaminated surfaces in transmission

Direct and indirect contact transmission is an established transmission route for several respiratory and gastrointestinal viruses, including rhinovirus, respiratory syncytial virus, norovirus, and rotavirus.^{7,47,65–67} However, the importance of indirect contact transmission (contact transmission involving contaminated surfaces; Figure 1) in the spread of respiratory viruses, including influenza, SARS-CoV and MERS-CoV, compared with other transmission routes is uncertain.^{6–8,68}

For contaminated surfaces to play a role in transmission, a respiratory pathogen must be shed into the environment, have the capacity to survive on surfaces, transfer to hands or other equipment at a concentration above the infectious dose, and be able to initiate infection through contact with the eyes, nose or mouth.¹¹

Human coronaviruses and influenza are shed in respiratory secretions.^{14,69} They can also survive in the gastrointestinal tract and have been associated with diarrhoea, which causes widespread environmental dissemination.^{14,69–74} In the case of SARS-CoV, viral loads in nasopharyngeal (up to 10^6 /mL) and stool (up to 10^9 /g) specimens may be high.⁶⁹ Titres of influenza in nasopharyngeal specimens (generally ranging from 10^5 to 10^7 , but can be up to 10^{11} copies/mL) and stool specimens (up to 10^7 /g) exhibit a similar range.^{57,74–76} Emerging data suggest that MERS-CoV are shed in approximately equal quantities to SARS-CoV.^{77,78} By contrast with the high titre shed from the respiratory and gastrointestinal tracts, the

infectious dose may be low. For example, the infectious dose for influenza can be <1 TCID₅₀, and <20 plaque-forming units for SARS-CoV.^{13,79}

SARS-CoV, MERS-CoV and influenza virus can survive on dry surfaces for extended periods, particularly when suspended in human secretions (Tables I and II), and may contaminate hand-touch sites in the field (Table III).

Viral and bacterial surface contamination can be transferred to hands, and serial transfer to a number of surfaces from contaminated hands may occur.^{11,42,80–85} For example, Bean *et al.* calculated that an infectious dose of virus could be transmitted for at least 2 h and possibly up to 8 h from stainless steel surfaces to hands.⁴²

In order for the virus to initiate indirect contact transmission, oral inoculation or contact with mucous membranes must occur to transfer sufficient viruses. Nasal inoculation is a frequent route for establishing influenza and SARS infection.^{86–90} Whereas oral inoculation has not been reported for SARS, it may occur for influenza and other viruses.^{13,91,92}

Thus, the steps necessary to facilitate indirect contact transmission of both SARS-CoV and influenza are established. Although data are more limited for MERS-CoV, it appears to have the key properties to facilitate indirect contact transmission.

Determining which route is most important is challenging, but it seems that direct contact, indirect contact, droplet and airborne transmission do occur with both SARS-CoV and influenza viruses on occasion.^{8,68} Few data are available evaluating transmission routes for coronaviruses, but the relative importance of the various routes for influenza virus has been evaluated through mathematical models, animal models, and intervention studies.^{9,93,94}

Several mathematical models have been applied to SARS transmission, but none has considered an environmental route.^{93,95} However, some influenza transmission models have evaluated the relative importance of airborne, droplet, and contact influenza transmission.^{9,96,97} Two of these models conclude that contact transmission of influenza is at least as important as airborne or droplet spread, whereas one study found that contact transmission was negligible compared with other routes.^{9,96,97} However, it is important to note that the relative contribution of contact, droplet, and airborne transmission depends on a combination of viral factors (e.g. capacity to survive on surfaces), host factors (e.g. frequency of contact hand contact with the nose) and environmental factors (e.g. size of enclosure and density of shedders). Varying these and other parameters will change the relative contribution of the various transmission routes.⁹

Several influenza transmission models have compared the importance of indirect contact transmission (involving surface contamination) with direct contact transmission (that occurs independently of surface contamination).^{98,99} One model indicates that indirect

transmission via contaminated surfaces generates touch frequency-dependent patterns whereas transmission via the air generates human density-dependent patterns.⁹⁸ Another model compared the involvement of droplet-contaminated versus hand-contaminated surfaces.⁹⁹ Droplet-contaminated surfaces were more likely to be involved in transmission than hand-contaminated surfaces (~10-fold difference), and large surfaces (such as table tops) had a higher transmission potential than small surfaces (such as door handles). A number of simplifying assumptions were made, which may be unsound – for example, that people touch portions of the fomite homogeneously, and that pathogens on fomites are homogeneously distributed. Also, transportation of contamination from one type of fomite to another via human hands was not modelled. Notwithstanding these limitations, the study provides some useful data on indirect contact transmission of influenza.

An alternative approach is the use of animal models. For example, a guinea-pig model evaluated the relative contribution of airborne, droplet, and indirect contact transmission.⁹⁴ Indirect contact transmission was evaluated by placing uninfected animals in cages vacated by experimentally infected animals without changing bedding, food dishes, and water bottles. Animals were exposed to these cages for 24 h and tested for infection using nasal washings. Around a quarter of exposed guinea-pigs became infected, which was less efficient than transmission through airborne and droplet experiments (25–100% efficiency). Experimental contamination of surfaces in the cages was unable to establish infection. Another guinea-pig model showed that increasing the temperature to 30°C blocked aerosol but not contact transmission of influenza.¹⁰⁰ This provides further evidence that the relative importance of the various transmission routes is context dependent.

A small number of studies have demonstrated that interventions in field settings to improve surface or hand hygiene reduce influenza transmission, demonstrating the importance of contact transmission.^{63,101,102} For example, introducing regular cleaning using disinfectant wipes reduced the rate of respiratory and diarrhoeal disease in elementary schools.⁶⁰

Implications for cleaning and disinfection, and infection prevention and control in healthcare settings

The likely contribution of droplet, direct and indirect contact, and to a lesser extent the airborne route in the transmission of influenza, SARS and MERS dictates that each route must be separately addressed by infection prevention and control interventions. The use of a surgical mask will protect the respiratory tract from droplets, an N95 (FFP3) respirator will protect the respiratory tract from droplet nuclei, and gloves, gowns and eye protection will prevent contact with mucous membranes and contamination of clothing or hands for subsequent nasal inoculation.¹⁰³ Emerging literature suggests that doffing PPE presents a

challenging risk for the acquisition of important viruses on hands.^{104,105} Thus, protocols should be in place for minimizing the risk of contamination of hands and clothing, and hand hygiene should be performed following removal of PPE.

The extended survival of influenza virus, SARS-CoV and MERS-CoV on surfaces (Tables I and II) and some evidence of contamination in field settings (Table III) argue for enhanced disinfection, particularly at the time of patient discharge.^{59,61} A range of hospital disinfectants are active against SARS-CoV and surrogates, and influenza, including alcohol, hypochlorites (bleach), quaternary ammonium compounds, and hydrogen peroxide, although inactivation is time and concentration dependent and will be influenced by other factors such as type of contaminated surface, specific product, and protein load.^{28,45,106,107} However, in-vitro disinfectant effectiveness is a poor predictor for the elimination of contamination from surfaces if cleaning/disinfection is inadequate, which is often the case in hospitals.^{108,109} Thus, there may be a role for automated room disinfection (ARD) systems, such as hydrogen peroxide vapor and ultraviolet (UV) light, at the time of terminal discharge of patients known to be infected with pandemic influenza or coronaviruses.^{45,108}

There may be the potential for extended survival of an infectious viral aerosol in patients' rooms following their discharge. Using MERS-CoV as an illustrative example, infectious aerosol above the infectious dose could be present after the discharge of the patient for up to 26 h, assuming no air changes in the room and depending on the shed titre (Table IV). ARD systems address both contaminated air and surfaces, which may be important if infectious aerosol above the infectious dose remains following patient discharge.

Another consideration is the requirement for large quantities of N95 (FFP3) respirators in the event of a pandemic of influenza or MERS/SARS. Stockpiles of N95 respirators required for a pandemic are large, and stock shortages were acknowledged during the 2009 N1H1 influenza pandemic.¹¹⁰ Both influenza virus and SARS-CoV surrogates have been shown to survive for extended periods on N95 respirator material.^{18,37,43} This survival represents a barrier to the reuse of N95 respirators. One approach is to disinfect the N95 respirators. Several candidate technologies have been evaluated for the disinfection of N95 respirators; UV light, hydrogen peroxide vapour, and ethylene oxide show most promise.¹¹¹

Conclusion

We reviewed the capacity of viruses with pandemic potential, influenza SARS-CoV and MERS-CoV, to survive on dry surfaces. The experimental methods used to test survival are important, but it seems that surface survival of SARS/MERS-CoV is greater than that of influenza virus. Important factors that influence the survival of these viruses on surfaces include: strain variations, a 'dose-response' relationship between the titre applied and

survival time, the surface substrate (including the ability to survive on materials used to make PPE), the suspending medium (with the addition of mucus increasing substantially the survival time of influenza), the mode of deposition, temperature and RH, and the method used to determine the presence of the virus (specifically culture versus the use of PCR to detect viral RNA). All three viruses are able to survive in an aerosol for a considerable length of time (>24 h), which may have important infection control implications.

Environmental sampling has been performed for influenza virus and human coronaviruses (including SARS-CoV) in a number of field settings. Most studies have used PCR to detect viral RNA, which may not necessarily represent the presence of viable virus, but should be seen as a marker of virus shedding. Some studies have demonstrated the presence of viable influenza virus on surfaces using cell culture. There is a wide range in terms of the frequency of sites contaminated with influenza virus or SARS-CoV RNA, ranging from <5% to >50%, including hand-touch sites.

The importance of indirect contact transmission is uncertain compared with other transmission routes, principally direct contact transmission, droplet, and airborne routes. Influenza virus, SARS-CoV and probably MERS-CoV are shed into the environment at concentrations far in excess of the infective dose, they can survive for extended periods on surfaces, and sampling has identified contamination of hospital surfaces. Contaminated surfaces could result in onward contamination of hands or equipment, which could then initiate inoculation through contact with the nose, eyes, or mouth. Thus, the steps required for indirect contact transmission are established. Mathematical modelling, animal models, and intervention trials suggest that contact transmission may be the most important route for influenza, but that is context dependent.

The infection prevention and control implications of these findings include the need to wear appropriate PPE to account for contact, droplet and airborne routes, paying particular attention to the risk of contamination of hands and clothing during PPE removal. The potential for inadequate distribution and contact time during manual cleaning and disinfection, combined with the risk of extended survival of infectious aerosol, may argue for the use of ARD systems. These systems may also have a role in disinfection and reuse of N95/FFP3 respirators.

Viruses with pandemic potential including influenza, MERS-CoV, and SARS-CoV can survive for extended periods on dry surfaces, cause contamination in field settings and may require enhanced cleaning and disinfection to assure effective infection prevention and control.

Conflict of interest statement

J.A.O. is a consultant to Gama Healthcare. All other authors have no conflict to declare.

Funding sources

None.

Appendix A: PubMed searches

coronavirus survival surfaces (June 11th, 2013: 9 studies)

influenza survival surfaces (June 11th, 2013: 29 studies)

coronavirus fomite transmission (June 20th, 2013: 8 studies)

influenza virus fomite transmission (June 20th, 2013: 43 studies)

coronavirus surface contamination (June 20th, 2013: 4 studies)

influenza virus surface contamination (June 20th, 2013: 14 studies)

disinfection influenza transmission (June 04th, 2014: 112 studies)

disinfection SARS transmission (June 04th, 2014: 35 studies)

Updated May 21st, 2014

References

1. de Groot RJ, Baker SC, Baric RS, *et al.* Middle East Respiratory Syndrome Coronavirus (MERS-CoV); Announcement of the Coronavirus Study Group. *J Virol* 2013;**87**:7790–7792.
2. Zaki AM, van Boheemen S, Bestebroer TM, Osterhaus AD, Fouchier RA. Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J Med* 2012;**367**:1814–1820.
3. Fineberg HV. Pandemic preparedness and response – lessons from the H1N1 influenza of 2009. *N Engl J Med* 2014;**370**:1335–1342.
4. Hayden FG. Respiratory viral threats. *Curr Opin Infect Dis* 2006;**19**:169–178.
5. World Health Organization. Annex C: Respiratory droplets. In: Atkinson J, Chartier Y, Pessoa-Silva CL, *et al.* editors. *Natural ventilation for infection control in health-care settings*. Geneva: WHO; 2009.
6. Bridges CB, Kuehnert MJ, Hall CB. Transmission of influenza: implications for control in health care settings. *Clin Infect Dis* 2003;**37**:1094–1101.
7. Boone SA, Gerba CP. Significance of fomites in the spread of respiratory and enteric viral disease. *Appl Environ Microbiol* 2007;**73**:1687–1696.
8. Brankston G, Gitterman L, Hirji Z, Lemieux C, Gardam M. Transmission of influenza A in human beings. *Lancet Infect Dis* 2007;**7**:257–265.
9. Spicknall IH, Koopman JS, Nicas M, Pujol JM, Li S, Eisenberg JN. Informing optimal environmental influenza interventions: how the host, agent, and environment alter dominant routes of transmission. *PLoS Comput Biol* 2010;**6**:e1000969.

10. Otter JA, Yezli S, Salkeld JA, French GL. Evidence that contaminated surfaces contribute to the transmission of hospital pathogens and an overview of strategies to address contaminated surfaces in hospital settings. *Am J Infect Control* 2013;**41**:S6–S11.
11. Otter JA, Yezli S, French GL. The role played by contaminated surfaces in the transmission of nosocomial pathogens. *Infect Control Hosp Epidemiol* 2011;**32**:687–699.
12. Weber DJ, Rutala WA, Miller MB, Huslage K, Sickbert-Bennett E. Role of hospital surfaces in the transmission of emerging health care-associated pathogens: norovirus, *Clostridium difficile*, and *Acinetobacter* species. *Am J Infect Control* 2010;**38**:S25–S33.
13. Yezli S, Otter JA. Minimum infective dose of the major human respiratory and enteric viruses transmitted through food and the environment. *Food Environ Microbiol* 2011;**3**:1–30.
14. Geller C, Varbanov M, Duval RE. Human coronaviruses: insights into environmental resistance and its influence on the development of new antiseptic strategies. *Viruses* 2012;**4**:3044–3068.
15. Kramer A, Schwebke I, Kampf G. How long do nosocomial pathogens persist on inanimate surfaces? A systematic review. *BMC Infect Dis* 2006;**6**:130.
16. van Doremalen N, Bushmaker T, Munster VJ. Stability of Middle East respiratory syndrome coronavirus (MERS-CoV) under different environmental conditions. *Euro Surveill* 2013;**18**. pii: 20590.
17. Chan KH, Peiris JS, Lam SY, Poon LL, Yuen KY, Seto WH. The effects of temperature and relative humidity on the viability of the SARS Coronavirus. *Adv Virol* 2011;**734690**.
18. Coulliette AD, Perry KA, Edwards JR, Noble-Wang JA. Persistence of the 2009 pandemic influenza A (H1N1) virus on N95 respirators. *Appl Environ Microbiol* 2013;**79**:2148–2155.
19. Casanova L, Rutala WA, Weber DJ, Sobsey MD. Survival of surrogate coronaviruses in water. *Water Res* 2009;**43**:1893–1898.
20. Mullis L, Saif LJ, Zhang Y, Zhang X, Azevedo MS. Stability of bovine coronavirus on lettuce surfaces under household refrigeration conditions. *Food Microbiol* 2012;**30**:180–186.
21. Yepiz-Gomez MS, Gerba CP, Bright KR. Survival of respiratory viruses on fresh produce. *Food Environ Virol* 2013;doi 10.1007/s12560-013-9114-4.

22. Wang XW, Li J, Guo T, *et al.* Concentration and detection of SARS coronavirus in sewage from Xiao Tang Shan Hospital and the 309th Hospital of the Chinese People's Liberation Army. *Water Sci Technol* 2005;**52**:213–221.
23. Shigematsu S, Dublineau A, Sawoo O, *et al.* Influenza A virus survival in water is influenced by the origin species of the host cell. *Influenza Other Respir Viruses* 2014;**8**:123–130.
24. Chmielewski R, Swayne DE. Avian influenza: public health and food safety concerns. *Annu Rev Food Sci Technol* 2011;**2**:37–57.
25. Nazir J, Haumacher R, Ike A, Stumpf P, Bohm R, Marschang RE. Long-term study on tenacity of avian influenza viruses in water (distilled water, normal saline, and surface water) at different temperatures. *Avian Dis* 2010;**54**:720–724.
26. Casanova LM, Jeon S, Rutala WA, Weber DJ, Sobsey MD. Effects of air temperature and relative humidity on coronavirus survival on surfaces. *Appl Environ Microbiol* 2010;**76**:2712–2717.
27. Muller A, Tillmann RL, Muller A, Simon A, Schildgen O. Stability of human metapneumovirus and human coronavirus NL63 on medical instruments and in the patient environment. *J Hosp Infect* 2008;**69**:406–408.
28. Rabenau HF, Cinatl J, Morgenstern B, Bauer G, Preiser W, Doerr HW. Stability and inactivation of SARS coronavirus. *Med Microbiol Immunol* 2005;**194**:1–6.
29. Lai MY, Cheng PK, Lim WW. Survival of severe acute respiratory syndrome coronavirus. *Clin Infect Dis* 2005;**41**:e67–71.
30. Duan SM, Zhao XS, Wen RF, *et al.* Stability of SARS coronavirus in human specimens and environment and its sensitivity to heating and UV irradiation. *Biomed Environ Sci* 2003;**16**:246–255.
31. Sizun J, Yu MW, Talbot PJ. Survival of human coronaviruses 229E and OC43 in suspension and after drying on surfaces: a possible source of hospital-acquired infections. *J Hosp Infect* 2000;**46**:55–60.
32. Zuo Z, de Abin M, Chander Y, Kuehn TH, Goyal SM, Pui DY. Comparison of spike and aerosol challenge tests for the recovery of viable influenza virus from non-woven fabrics. *Influenza Other Respi Viruses* 2013;doi 10.1111/irv.12095.
33. Mukherjee DV, Cohen B, Bovino ME, Desai S, Whittier S, Larson EL. Survival of influenza virus on hands and fomites in community and laboratory settings. *Am J Infect Control* 2012;**40**:590–594.

34. 34Greatorex JS, Digard P, Curran MD, *et al.* Survival of influenza A (H1N1) on materials found in households: implications for infection control. *PLoS One* 2011;**6**:e27932.
35. Dublineau A, Batejat C, Pinon A, Burguiere AM, Leclercq I, Manuguerra JC. Persistence of the 2009 pandemic influenza A (H1N1) virus in water and on non-porous surface. *PLoS One* 2011;**6**:e28043.
36. Wood JP, Choi YW, Chappie DJ, Rogers JV, Kaye JZ. Environmental persistence of a highly pathogenic avian influenza (H5N1) virus. *Environ Sci Technol* 2010;**44**:7515–7520.
37. Sakaguchi H, Wada K, Kajioka J, *et al.* Maintenance of influenza virus infectivity on the surfaces of personal protective equipment and clothing used in healthcare settings. *Environ Health Prev Med* 2010;**15**:344–349.
38. McDevitt J, Rudnick S, First M, Spengler J. Role of absolute humidity in the inactivation of influenza viruses on stainless steel surfaces at elevated temperatures. *Appl Environ Microbiol* 2010;**76**:3943–3947.
39. Thomas Y, Vogel G, Wunderli W, *et al.* Survival of influenza virus on banknotes. *Appl Environ Microbiol* 2008;**74**:3002–3007.
40. Noyce JO, Michels H, Keevil CW. Inactivation of influenza A virus on copper versus stainless steel surfaces. *Appl Environ Microbiol* 2007;**73**:2748–2750.
41. Tiwari A, Patnayak DP, Chander Y, Parsad M, Goyal SM. Survival of two avian respiratory viruses on porous and nonporous surfaces. *Avian Dis* 2006;**50**:284–287.
42. Bean B, Moore BM, Sterner B, Peterson LR, Gerding DN, Balfour HH, Jr. Survival of influenza viruses on environmental surfaces. *J Infect Dis* 1982;**146**:47–51.
43. Casanova L, Rutala WA, Weber DJ, Sobsey MD. Coronavirus survival on healthcare personal protective equipment. *Infect Control Hosp Epidemiol* 2010;**31**:560–561.
44. Dowell SF, Simmerman JM, Erdman DD, *et al.* Severe acute respiratory syndrome coronavirus on hospital surfaces. *Clin Infect Dis* 2004;**39**:652–657.
45. Goyal SM, Chander Y, Yezli S, Otter JA. Evaluating the virucidal efficacy of hydrogen peroxide vapour. *J Hosp Infect* 2014;**86**:255–259.
46. Parker ER, Dunham WB, MacNeal WJ. Resistance of the Melbourne strain of influenza virus to desiccation. *J Lab Clin Med* 1944;**29**:37–42.
47. Musher DM. How contagious are common respiratory tract infections? *N Engl J Med* 2003;**348**:1256–1266.
48. Gerone PJ, Couch RB, Keefer GV, Douglas RG, Derrenbacher EB, Knight V. Assessment of experimental and natural viral aerosols. *Bacteriol Rev* 1966;**30**:576–588.

49. Tran K, Cimon K, Severn M, Pessoa-Silva CL, Conly J. Aerosol generating procedures and risk of transmission of acute respiratory infections to healthcare workers: a systematic review. *PLoS ONE* 2012;**7**:e35797.
50. Bischoff WE, Swett K, Leng I, Peters TR. Exposure to influenza virus aerosols during routine patient care. *J Infect Dis* 2013;**207**:1037–1046.
51. Thompson KA, Pappachan JV, Bennett AM, *et al.* Influenza aerosols in UK hospitals during the H1N1 (2009) pandemic – the risk of aerosol generation during medical procedures. *PLoS One* 2013;**8**:e56278.
52. Ijaz MK, Brunner AH, Sattar SA, Nair RC, Johnson-Lussenburg CM. Survival characteristics of airborne human coronavirus 229E. *J Gen Virol* 1985;**66**(Pt 12):2743–2748.
53. Schaffer FL, Soergel ME, Straube DC. Survival of airborne influenza virus: effects of propagating host, relative humidity, and composition of spray fluids. *Arch Virol* 1976;**51**:263–273.
54. Mitchell CA, Guerin LF. Influenza A of human, swine, equine and avian origin: comparison of survival in aerosol form. *Can J Comp Med* 1972;**36**:9–11.
55. Tellier R. Review of aerosol transmission of influenza A virus. *Emerg Infect Dis* 2006;**12**:1657–1662.
56. Indriani R, Samaan G, Gultom A, *et al.* Environmental sampling for avian influenza virus A (H5N1) in live-bird markets, Indonesia. *Emerg Infect Dis* 2010;**16**:1889–1895.
57. Killingley B, Greatorex J, Cauchemez S, *et al.* Virus shedding and environmental deposition of novel A (H1N1) pandemic influenza virus: interim findings. *Health Technol Assess* 2010;**14**:237–354.
58. Simmerman JM, Suntarattiwong P, Levy J, *et al.* Influenza virus contamination of common household surfaces during the 2009 influenza A (H1N1) pandemic in Bangkok, Thailand: implications for contact transmission. *Clin Infect Dis* 2010;**51**:1053–1061.
59. Pappas DE, Hendley JO, Schwartz RH. Respiratory viral RNA on toys in pediatric office waiting rooms. *Pediatr Infect Dis J* 2010;**29**:102–104.
60. Bright KR, Boone SA, Gerba CP. Occurrence of bacteria and viruses on elementary classroom surfaces and the potential role of classroom hygiene in the spread of infectious diseases. *J Sch Nurs* 2010;**26**:33–41.
61. Macias AE, de la Torre A, Moreno-Espinosa S, Leal PE, Bourlon MT, Ruiz-Palacios GM. Controlling the novel A (H1N1) influenza virus: don't touch your face! *J Hosp Infect* 2009;**73**:280–281.

62. Boone SA, Gerba CP. The occurrence of influenza A virus on household and day care center fomites. *J Infect* 2005;**51**:103–109.
63. Booth TF, Kournikakis B, Bastien N, *et al.* Detection of airborne severe acute respiratory syndrome (SARS) coronavirus and environmental contamination in SARS outbreak units. *J Infect Dis* 2005;**191**:1472–1477.
64. Memish ZA, Almasri M, Assirri A, *et al.* Environmental sampling for respiratory pathogens in Jeddah airport during the 2013 Hajj season. *Am J Infect Control* 2014;**42**:1266–1269.
65. Gwaltney JM, Jr, Hendley JO. Transmission of experimental rhinovirus infection by contaminated surfaces. *Am J Epidemiol* 1982;**116**:828–833.
66. Abad FX, Pinto RM, Bosch A. Survival of enteric viruses on environmental fomites. *Appl Environ Microbiol* 1994;**60**:3704–3710.
67. Hall CB. Respiratory syncytial virus: its transmission in the hospital environment. *Yale J Biol Med* 1982;**55**:219–223.
68. Chan PK, Tang JW, Hui DS. SARS: clinical presentation, transmission, pathogenesis and treatment options. *Clin Sci (Lond)* 2006;**110**:193–204.
69. Hung IF, Cheng VC, Wu AK, *et al.* Viral loads in clinical specimens and SARS manifestations. *Emerg Infect Dis* 2004;**10**:1550–1557.
70. Zhang XM, Herbst W, Kousoulas KG, Storz J. Biological and genetic characterization of a hemagglutinating coronavirus isolated from a diarrhoeic child. *J Med Virol* 1994;**44**:152–161.
71. Vabret A, Dina J, Gouarin S, Petitjean J, Corbet S, Freymuth F. Detection of the new human coronavirus HKU1: a report of 6 cases. *Clin Infect Dis* 2006;**42**:634–639.
72. Peiris JS, Chu CM, Cheng VC, *et al.* Clinical progression and viral load in a community outbreak of coronavirus-associated SARS pneumonia: a prospective study. *Lancet* 2003;**361**:1767–1772.
73. Pinsky BA, Mix S, Rowe J, Ikemoto S, Baron EJ. Long-term shedding of influenza A virus in stool of immunocompromised child. *Emerg Infect Dis* 2010;**16**:1165–1167.
74. Chan MC, Lee N, Chan PK, Leung TF, Sung JJ. Fecal detection of influenza A virus in patients with concurrent respiratory and gastrointestinal symptoms. *J Clin Virol* 2009;**45**:208–211.
75. Kaiser L, Fritz RS, Straus SE, Gubareva L, Hayden FG. Symptom pathogenesis during acute influenza: interleukin-6 and other cytokine responses. *J Med Virol* 2001;**64**:262–268.

76. Hall CB, Douglas RG, Jr, Geiman JM, Meagher MP. Viral shedding patterns of children with influenza B infection. *J Infect Dis* 1979;**140**:610–613.
77. Drosten C, Seilmaier M, Corman VM, *et al*. Clinical features and virological analysis of a case of Middle East respiratory syndrome coronavirus infection. *Lancet Infect Dis* 2013;**13**:745–751.
78. Guery B, Poissy J, el Mansouf L, *et al*. Clinical features and viral diagnosis of two cases of infection with Middle East Respiratory Syndrome coronavirus: a report of nosocomial transmission. *Lancet* 2013;**381**:2265–2272.
79. Watanabe T, Bartrand TA, Weir MH, Omura T, Haas CN. Development of a dose–response model for SARS coronavirus. *Risk Anal* 2010;**30**:1129–1138.
80. Oelberg DG, Joyner SE, Jiang X, Laborde D, Islam MP, Pickering LK. Detection of pathogen transmission in neonatal nurseries using DNA markers as surrogate indicators. *Pediatrics* 2000;**105**:311–315.
81. Barker J, Vipond IB, Bloomfield SF. Effects of cleaning and disinfection in reducing the spread of Norovirus contamination via environmental surfaces. *J Hosp Infect* 2004;**58**:42–49.
82. Guerrero DM, Nerandzic MM, Jury LA, Jinno S, Chang S, Donskey CJ. Acquisition of spores on gloved hands after contact with the skin of patients with *Clostridium difficile* infection and with environmental surfaces in their rooms. *Am J Infect Control* 2012;**40**:556–558.
83. Rusin P, Maxwell S, Gerba C. Comparative surface-to-hand and fingertip-to-mouth transfer efficiency of gram-positive bacteria, gram-negative bacteria, and phage. *J Appl Microbiol* 2002;**93**:585–592.
84. Jiang X, Dai X, Goldblatt S, *et al*. Pathogen transmission in child care settings studied by using a cauliflower virus DNA as a surrogate marker. *J Infect Dis* 1998;**177**:881–888.
85. Rheinbaben F, Schunemann S, Gross T, Wolff MH. Transmission of viruses via contact in a household setting: experiments using bacteriophage straight phiX174 as a model virus. *J Hosp Infect* 2000;**46**:61–66.
86. Nagata N, Iwata N, Hasegawa H, *et al*. Pathology and virus dispersion in cynomolgus monkeys experimentally infected with severe acute respiratory syndrome coronavirus via different inoculation routes. *Int J Exp Pathol* 2007;**88**:403–414.
87. McAuliffe J, Vogel L, Roberts A, *et al*. Replication of SARS coronavirus administered into the respiratory tract of African Green, rhesus and cynomolgus monkeys. *Virology* 2004;**330**:8–15.

88. Henle W, Henle G, Stokes J, Maris EP. Experimental exposure of human subjects to viruses of influenza. *J Immunol* 1946;**52**:145–165.
89. Frankova V. Inhalatory infection of mice with influenza A0/PR8 virus. I. The site of primary virus replication and its spread in the respiratory tract. *Acta Virol* 1975;**19**:29–34.
90. Qin C, Wang J, Wei Q, *et al.* An animal model of SARS produced by infection of *Macaca mulatta* with SARS coronavirus. *J Pathol* 2005;**206**:251–259.
91. Quan FS, Compans RW, Kang SM. Oral vaccination with inactivated influenza vaccine induces cross-protective immunity. *Vaccine* 2012;**30**:180–188.
92. VanDalen KK, Franklin AB, Mooers NL, Sullivan HJ, Shriner SA. Shedding light on avian influenza H4N6 infection in mallards: modes of transmission and implications for surveillance. *PLoS One* 2010;**5**:e12851.
93. van Kleef E, Robotham JV, Jit M, Deeny SR, Edmunds WJ. Modelling the transmission of healthcare associated infections: a systematic review. *BMC Infect Dis* 2013;**13**:294.
94. Mubareka S, Lowen AC, Steel J, Coates AL, Garcia-Sastre A, Palese P. Transmission of influenza virus via aerosols and fomites in the guinea pig model. *J Infect Dis* 2009;**199**:858–865.
95. Kwok KO, Leung GM, Lam WY, Riley S. Using models to identify routes of nosocomial infection: a large hospital outbreak of SARS in Hong Kong. *Proc Biol Sci* 2007;**274**:611–617.
96. Atkinson MP, Wein LM. Quantifying the routes of transmission for pandemic influenza. *Bull Math Biol* 2008;**70**:820–867.
97. Nicas M, Jones RM. Relative contributions of four exposure pathways to influenza infection risk. *Risk Anal* 2009;**29**:1292–1303.
98. Li S, Eisenberg JN, Spicknall IH, Koopman JS. Dynamics and control of infections transmitted from person to person through the environment. *Am J Epidemiol* 2009;**170**:257–265.
99. Zhao J, Eisenberg JE, Spicknall IH, Li S, Koopman JS. Model analysis of fomite mediated influenza transmission. *PLoS One* 2012;**7**:e51984.
100. Lowen AC, Steel J, Mubareka S, Palese P. High temperature (30 degrees C) blocks aerosol but not contact transmission of influenza virus. *J Virol* 2008;**82**:5650–5652.
101. Cowling BJ, Chan KH, Fang VJ, *et al.* Facemasks and hand hygiene to prevent influenza transmission in households: a cluster randomized trial. *Ann Intern Med* 2009;**151**:437–446.

102. Apisarnthanarak A, Apisarnthanarak P, Cheevakumjorn B, Mundy LM. Intervention with an infection control bundle to reduce transmission of influenza-like illnesses in a Thai preschool. *Infect Control Hosp Epidemiol* 2009;**30**:817–822.
103. Seto WH, Tsang D, Yung RW, *et al.* Effectiveness of precautions against droplets and contact in prevention of nosocomial transmission of severe acute respiratory syndrome (SARS). *Lancet* 2003;**361**:1519–1520.
104. Beam EL, Gibbs SG, Boulter KC, Beckerdite ME, Smith PW. A method for evaluating health care workers' personal protective equipment technique. *Am J Infect Control* 2011;**39**:415–420.
105. Johnson DW, Sullivan JN, Piquette CA, *et al.* Lessons learned: critical care management of patients with Ebola in the United States. *Crit Care Med* 2015;**43**:1157–1164.
106. Hulkower RL, Casanova LM, Rutala WA, Weber DJ, Sobsey MD. Inactivation of surrogate coronaviruses on hard surfaces by health care germicides. *Am J Infect Control* 2011;**39**:401–407.
107. Jeong EK, Bae JE, Kim IS. Inactivation of influenza A virus H1N1 by disinfection process. *Am J Infect Control* 2010;**38**:354–360.
108. Otter JA, Yezli S, Perl TM, Barbut F, French GL. Is there a role for “no-touch” automated room disinfection systems in infection prevention and control? *J Hosp Infect* 2013;**83**:1–13.
109. Carling PC, Parry MM, Rupp ME, Po JL, Dick B, Von Beheren S. Improving cleaning of the environment surrounding patients in 36 acute care hospitals. *Infect Control Hosp Epidemiol* 2008;**29**:1035–1041.
110. Anonymous. Interim guidance on infection control measures for 2009 H1N1 influenza in healthcare settings, including protection of healthcare personnel. *Miss RN* 2009;**71**:13–18.
111. Viscusi DJ, Bergman MS, Eimer BC, Shaffer RE. Evaluation of five decontamination methods for filtering facepiece respirators. *Ann Occup Hyg* 2009;**53**:815–827.

Table I

Survival of SARS-CoV, MERS-CoV, and surrogates on dry surfaces

| Study | Year | Location | Test virus | Load applied | Substrate(s) | Suspending medium | Volume applied (μL) | Temperature (°C)/RH (%) | Drying time (min) for time 0 sample | Results |
|---|------|-----------|------------|-----------------|-------------------|--------------------------|---------------------|-------------------------|-------------------------------------|---|
| van Doremalen <i>et al.</i> ¹⁶ | 2013 | USA | MERS-CoV | 10 ⁵ | Steel and plastic | Cell culture medium only | 100 | Variable | 10 | Viable virus detected after 48 h at 20°C/40% RH. Less survival at 30°C/80% RH (8 h) and 30°C/30% RH (24 h). Half-life ranged from ~0.5 to 1 h. |
| Chan <i>et al.</i> ¹⁷ | 2011 | Hong Kong | SARS-CoV | 10 ⁵ | Plastic | Cell culture medium only | 10 | Variable | Until visibly dry | SARS-CoV survived for 5 days with <10-fold reduction in titre at room temperature and humidity, and was viable for >20 days. The virus was more stable at lower |

| | | | | | | | | | | |
|--------------------------------------|------|-----|-----------|---------|--|--------------------------|----|----------|-------------------|--|
| | | | | | | | | | | temperatures (28 vs 38°C) and lower humidity (80–89% vs >95%). The reduction in viral titre was similar in suspension compared with virus dried on surfaces. |
| Casanova <i>et al.</i> ²⁶ | 2010 | USA | TGEV | $>10^3$ | Latex/nitrile gloves, N95 respirator, hospital scrubs, isolation gowns | Cell culture medium only | 10 | 20/50 | 0 | TGEV survived with $<10^2$ reduction on all items after 4 h and was detected on some items after 24 h |
| Casanova <i>et al.</i> ¹⁹ | 2009 | USA | TGEV, MHV | 10^5 | Stainless steel discs | Cell culture medium only | 10 | Variable | Until visibly dry | Both TGEV and MHV could survive in excess of 28 days under some conditions, with lower temperature and relative humidity |

| | | | | | | | | | | |
|-------------------------------------|------|---------|---|-------------------------------------|---|--|---------------|---------|-------------------|---|
| | | | | | | | | | | resulting in improved survival. TGEV and MHV did not differ significantly in their survival properties. |
| Muller <i>et al.</i> ²⁷ | 2008 | Germany | HCoV-NL63, human metapneumovirus | Not specified | Latex gloves, thermometer caps, stethoscopes, plastic table | Cell culture medium only | Not specified | Ambient | Not specified | Viable virus not detected after drying; viral RNA detectable for up to 7 days |
| Rabenau <i>et al.</i> ²⁸ | 2005 | Germany | SARS-CoV, HCoV-229E, herpes simplex virus, adenovirus | 10^6 – 10^7 | Polystyrene Petri dish | Cell culture medium \pm 20% fetal calf serum | 500 | Ambient | Until visibly dry | SARS-CoV, adenovirus and herpes simplex virus survived >6 days. HCoV-229E survived for <72 h. The addition of FCS made little impact on survival times. |
| Lai <i>et al.</i> ²⁹ | 2005 | China | SARS-CoV | Dilution series (10^2 – 10^4) | Paper, disposable gowns, | Cell culture medium + | 5 | Ambient | Until visibly dry | There was a dose–response in terms of survival |

| | | | | | | | | | | |
|----------------------------------|------|-------|----------|-----------------|---|-----------------------------------|-----|---------|---------------------|--|
| | | | | | cotton gowns | 2% fetal calf serum | | | | times of all materials, with more concentrated inocula surviving longer. Survival times ranged from 5 min (10 ² load on a cotton gown) to 2 days (10 ⁴ load on disposable gown). |
| Duan <i>et al.</i> ³⁰ | 2003 | China | SARS-CoV | 10 ⁶ | Wood board, glass, mosiac, metal, cloth, paper, filter paper, plastic | Cell culture medium only | 300 | Ambient | No time 0 sample | Viability was assessed semiquantitatively and SARS-CoV was able to survive, albeit with reduced infectivity, for >72 h on all surfaces tested, and for >120 h on metal, cloth and filter paper. Additionally, virus survived for >72 h on cotton cloth |

in an experimentally
dried enclosure.

| | | | | | | | | | | |
|-----------------------------------|------|--------|-------------------------|-----------------|--|-----------------------------------|----|---------|---|--|
| Sizun <i>et al.</i> ³¹ | 2000 | Canada | HCoV-229E, HCoV-OC43 | 10 ³ | Aluminium, cotton gauze, latex gloves | Cell culture medium only | 10 | Ambient | Until visibly dry (15–45 min) | Viability fell to below detectable levels after 6 h for 229E and 2 h for OC43. |
|-----------------------------------|------|--------|-------------------------|-----------------|--|-----------------------------------|----|---------|---|--|

SARS, severe acute respiratory syndrome; CoV, human coronavirus; MERS, Middle East respiratory syndrome; RH, relative humidity; TGEV, transmissible gastroenteritis coronavirus; MHV, mouse hepatitis virus.

Table II

Survival of influenza viruses on dry surfaces

| Study | Year | Location | Test virus | Load applied | Substrate (s) | Suspending medium | Volume applied (μL) | Temp (°C)/RH (%) | Drying time (min) for time 0 sample | Results |
|--|------|----------|------------------------|-----------------|------------------------------|----------------------------------|---------------------|------------------|-------------------------------------|--|
| van Doremale <i>et al.</i> ¹⁶ | 2013 | USA | H1N1 (human isolate) | 10 ⁵ | Steel and plastic | Cell culture medium only | 100 | Variable | 10 | No viable virus recovered after 4 h. No difference between plastic and steel. |
| Coulliette <i>et al.</i> ¹⁸ | 2013 | USA | H1N1 (pandemic strain) | 10 ⁴ | Coupons from N95 respirators | Cell culture medium/2% FBS/mucin | 100 | Variable | 60 | 10 ² TCID ₅₀ per coupon recovered from time 0 samples (after drying). Viable virus was recovered after 6 days with a 10-fold reduction. Viral survival was |

| | | | | | | | | | | |
|---------------------------------|------|-----|----------------------|--|-------------------------|--------------------------|----|---------|--|--|
| | | | | | | | | | | longer in FBS and mucin compared with cell culture medium. Lower absolute humidity favoured longer survival. |
| Zuo <i>et al.</i> ³² | 2013 | USA | Avian influenza H9N9 | Liquid spike (10 ⁴ –10 ⁵) | Three non-woven fabrics | Cell culture medium only | 20 | Ambient | 0 min; until visibly dry; 30 min after visibly dry | Viable virus survival for >1 h on each of the materials tested; survival times varied significantly by material. Survival on hydrophilic nylon lower than on hydrophobic materials. Choice of eluent did not significantly affect recovery. Virus recovery following |

| | | | | | | | | | | |
|---------------------------------------|------|-----|--|--|---|--------------------------|-----|---------|-----|--|
| | | | | | | | | | | deposition as an aerosol was considerably lower. |
| Mukherjee <i>et al.</i> ³³ | 2012 | USA | Field study of 20 influenza-infected individuals | Participants coughed or sneezed on hands then touched surfaces | Door handle, telephone, pillowcase, cotton handkerchief | n/a | n/a | Ambient | n/a | Virus RNA recovered from three door handles and one telephone; no samples were tissue culture positive. |
| | | | H1N1 (recovered from two participants) | Dilution series (10–10 ⁵) | Formica, vinyl, stainless steel, cotton pillowcase, facial tissue | Cell culture medium only | 20 | Ambient | 5 | Viable virus detected by tissue culture from some hard surfaces at higher applied load for up to 1 h; no viable virus detectable by tissue culture after 1 h; virus RNA detectable after 1 h |

on some surfaces.

| | | | | | | | | | | |
|--|----------|----|--|-----------------|--|--|----|-----------------|---|---|
| Greatorex <i>et al.</i> ³⁴ | 201 1 | UK | H1N1 (PR8) | 10 ⁶ | Common porous and non-porous household materials | Cell culture medium plus 1% bovine serum albumin | 10 | 17–21/23– 24 | 0 (drying times ranged from 5 min to 7 h) | Viral RNA detected with minimal reduction on most surfaces over 24 h; viral infectivity falls away more rapidly, with infective virus at low titre detectable from most surfaces at 4 h but from only stainless steel at 9 h |
| | | | H1N1 (AH04): recent clinical isolate | 10 ⁴ | | Cell culture medium only | | | | Semiquantitative fluorescence assay indicated the presence of virus at 4–24 h on hard surfaces but <4 h on porous surfaces. |

| | | | | | | | | | | |
|---------------------------------------|------|-------|------------------------------------|-----------------|---------------------------------------|--------------------------|-----|------------|--------|--|
| Dublineau <i>et al.</i> ³⁵ | 2011 | Paris | H1N1 seasonal and pandemic strains | 10^5 – 10^6 | Watch glass | Cell culture medium only | 50 | Variable | 5–17 h | Both viruses survived for >3 days under all conditions tested; pandemic H1N1 survived for >7 days at 35°C and 2 months at 4°C. |
| Wood <i>et al.</i> ³⁶ | 2010 | USA | H5N1 | 10^6 | Glass and galvanized steel | Cell culture medium only | 100 | 4/variable | 60 | Influenza stable at low temperature, regardless of humidity, with 13-day survival and reduction by factor of <1 on both substrates. Surface survival not tested at room temperature. |
| Sakaguchi <i>et al.</i> ³⁷ | 2010 | Japan | H1N1 | 10^4 | Personal protective equipment: rubber | Cell culture medium only | 500 | 25.2/55 | 0 | The haemagglutinin titre of the virus remained stable on |

| | | | | | | | | | | |
|---|----------|-----|------------|----------------------------------|--|----------------------------------|----|----------|---|--|
| | | | | | gloves, N95 mask, surgical mask, Tyvek gown, coated wood, steel | | | | | all surfaces up to 24 h. The virus remained infective by TCID ₅₀ on all materials up to 8 h, and on rubber for up to 24 h. |
| McDevitt <i>et al.</i> ³⁸ | 201 0 | USA | H1N1 (PR8) | 10 ⁴ –10 ⁵ | Stainless steel | Purchased virus suspension | 50 | Variable | Until visibly dry (~30 min) | Virus survival assessed at 15, 30 and 60 mins at variable temperature 55–65°C) and relative humidity (25–75%). Virus survived for >60 min with a 10 ^{1.5} reduction at the lowest temperature/humid ity combination (55°C/25%). |

| | | | | | | | | | | |
|---------------------------------------|----------|-------------|--|-----------------|------------|--------------------------|----|-----------------------|---|--|
| | | | | | | | | | | Linear association between increasing humidity and logarithmic reduction. |
| Thomas <i>et al.</i> ³⁹ | 200 8 | Switzerland | H3N2 (2 strains), H1N1 and influenza B | 10^3 – 10^8 | Bank notes | Cell culture medium only | 50 | 21–28 (avg. 22)/30–50 | Dried under laminar airflow; time not specified | Survival varied by strain from 3 h to 3 days, depending on the virus tested. |
| | | | Spiked pooled negative nasopharyngeal secretions | | Bank notes | Cell culture medium only | 50 | 21–28 (avg. 22)/30–50 | Dried under laminar airflow; time not specified | Higher inocula survived for longer on surfaces; the addition of respiratory mucus significantly increased survival, usually from hours to up to 17 days. |

| | | | | | | | | | | |
|------------------------------------|----------|-----|--|-----------------|--|-----------------------------------|----|-----------------------------|---|---|
| | | | Influenza- positive nasopharyngeal secretions | | Bank notes | Cell culture medium only | 50 | 21–28 (avg. 22)/30–50 | Dried under laminar airflow; time not specifie d | Infective influenza recovered from 7/14 (50%) of notes after 24 h, 5/14 (36%) of notes after 48 h, and in one case, after 12 days. |
| Noyce <i>et al.</i> ⁴⁰ | 200 7 | UK | H1N1 | 10 ⁶ | Stainless steel or copper | Cell culture medium only | 20 | 20–24/50– 60 | Not specifie d | 10 ⁵ viable virus recovered from stainless steel after 24 h vs 10 ² viable virus on copper after 6 h |
| Tiwari <i>et al.</i> ⁴¹ | 200 6 | USA | Avian influenza virus, avian metapneumovir us | 10 ⁴ | Steel, wood, tile, tire, gumboot, feather, egg shell, egg tray, plastic, latex, cotton and | Cell culture medium only | 10 | Ambient | Until visibly dry (~30–4 0 min) | Both viruses survive for up to 72 h on most surfaces tested. Influenza survived for up to 6 days on latex and feather. |

| | | | | | | | | | | |
|----------------------------------|----------|-----|--|----------------------------------|---|-----------------------------------|-----|-----------------|----------------|--|
| | | | | | polyester | | | | | |
| Bean <i>et al.</i> ⁴² | 198 2 | USA | H1N1 and influenza B clinical isolates | 10 ² –10 ⁴ | Steel, plastic, cotton handkerchie f, paper tissue, magazine page, cotton panamas | Cell culture medium only | 100 | 26–29/35– 56 | Up to 1.5 h | Viruses survived for 48–72 h on non-porous surfaces (steel and plastic) and for shorter periods on porous surfaces. Influenza A survived significantly longer than influenza B. Viruses dried on to surfaces could be transferred to hands from all surfaces for 15 min, and from steel for 24 h. |

FBS, fetal bovine serum; TCID, tissue culture infectious dose; avg., average.

Table III

Field sampling for influenza and human coronaviruses including SARS-CoV environmental contamination

| Study | Year | Setting and location | Sites sampled | Sampling method | No. of samples | No. positive (%) | Notes |
|--|------|--|--|---|----------------|------------------|--|
| Influenza | | | | | | | |
| Indriani <i>et al.</i> ⁵⁶ | 2010 | Live-bird markets, Indonesia | 27 sites were sampled at 83 live-bird markets for avian influenza (H5N1) | Cotton swabs; PCR for viral RNA and viral culture | 1862 (PCR) | 280 (15) | 39 (47%) markets contaminated at one or more site. Structured questionnaire to assess risk factors for contamination. One province and markets that slaughtered birds associated with contamination; zoning of poultry activities and daily disposal of solid waste were protective. |
| | | | | | 280 (culture) | 13 (4.6) | |
| Killingley <i>et al.</i> ⁵⁷ | 2010 | Influenza-infected adults in hospital and community settings in and around Nottingham, | 19 patients (daily) and their immediate environment (every other day) | Moistened cotton swabs; PCR for viral RNA and viral culture | 397 | 2 (0.5) | Live virus recovered from 1/2 positive surfaces. 54% of subjects took an antiviral drug, which may have influenced shedding. Duration of virus shedding had a |

| | | | | | | | |
|---------------------------------------|------|---|---|---|-----|-----------|---|
| | | UK | were sampled. | | | | mean of 6.2 days and a range of 3–10 days. |
| Simmerman <i>et al.</i> ⁵⁸ | 2010 | 90 children with influenza in Bangkok, Thailand. Households were randomized to obtain handwashing education or not. | Six household items in 90 households | Moistened rayon tipped swabs; PCR for viral RNA and viral culture | 540 | 18 (3.3) | 16 (17.8%) of the 90 households had one or more sample positive for influenza by PCR. Nine TV remotes, six toys, two bathroom knobs and one light switch had positive results. No viable virus was detected by culture. |
| Pappas <i>et al.</i> ⁵⁹ | 2010 | Toys in the waiting room of a general paediatric practice in Virginia, USA | Hard surfaces and fabric toy samples on three separate occasions | Moistened swab; samples tested for picornavirus, RSV and influenza by PCR | 52 | 1 (1.9) | 19.2% of the toys were contaminated with picornavirus RNA. |
| Bright <i>et al.</i> ⁶⁰ | 2010 | Surfaces in three elementary school classrooms in Seattle, Washington, USA | Standardized surfaces sampled in the morning, at midday and in the afternoon. | Moistened swabs; PCR for viral RNA | 54 | 13 (24.1) | Also, norovirus RNA was found on 16.4% of 55 surfaces sampled. |
| Macias <i>et al.</i> ⁶¹ | 2009 | Hospital in Mexico City, Mexico | Samples collected from hands and surfaces in the rooms of patients | Swabs; PCR for viral RNA | 13 | 5 (38.5) | In one case, 1/5 surfaces (a bed rail) was positive from a patient's room 72 h after patient discharge and terminal cleaning. 5/6 |

| | | | | | | | | |
|------------------------------------|------|--|--|---|-----|--------------|--|--|
| Human coronavirus | | | with confirmed influenza | | | | | samples from patient hands were positive for influenza. |
| | | | Samples from eight homes | | | | | None of 33 surfaces sampled during summer months vs 59% of 59 samples during March. |
| | | | Samples from 14 day-care centres | | | | | Influenza was detected on 23% of surfaces during the autumn and 53% during the spring. |
| Boone and Gerba ⁶² | 2005 | Homes and day-care centres in Tucson, Arizona, USA | | Moistened swabs; PCR for viral RNA | 92 | 35 (38.0) | | |
| | | | | | 218 | – | | |
| Human coronavirus | | | 19 rooms in SARS units and ‘control’ areas not housing SARS patients | Moistened swabs; PCR for viral RNA and viral culture | 85 | 3 (3.5) | | Positive sites were a bed table, a television remote control and a refrigerator handle in a nurses’ medication station. All swabs were culture negative. Two (5%) of 40 air-slit samples were positive for SARS-CoV. |
| | | | | | | | | |
| | | | | | | | | |
| Booth <i>et al.</i> ⁶³ | 2005 | Hospitals in Toronto, Canada | | | | | | |
| Human coronavirus | | | SARS-infected patient areas (patient rooms, nursing stations, emergency department) | Moistened swabs; PCR for viral RNA and viral culture | 63 | 24 (38.1) | | All swabs were culture negative. |
| | | | | | | | | |
| | | | | | | | | |
| Dowell <i>et al.</i> ⁴⁴ | 2004 | Hospitals in Bangkok, Thailand and Taipei, Taiwan | | | | | | |

| | | | | | | | |
|------------------------------------|------|---------------------------------|--|---|----|---------|---|
| | | | Public areas | | 31 | 2 (6.4) | |
| Memish <i>et al.</i> ⁶⁴ | 2014 | Jeddah airport, Saudi Arabia | Various frequently touched items in public areas | Moistened swabs; PCR panel for viral culture | 40 | 3 (7.5) | Human coronavirus (OC43/HKU1) RNA was identified from surfaces. Influenza B virus RNA was identified from 1/18 air samples, but was not identified on surfaces. |

SARS-CoV, severe acute respiratory syndrome coronavirus; PCR, polymerase chain reaction.

Table IV


Calculating the time that an infectious aerosol shed by a patient infected with Middle East respiratory syndrome coronavirus could survive

| Shed titre | Time to reach 20 virus particles |
|------------|----------------------------------|
| 1,000,000 | 26 h |
| 100,000 | 20 h |
| 10,000 | 15 h |
| 1000 | 9 h |
| 100 | 4 h |

The calculation assumes an infectious dose equal to severe acute respiratory syndrome coronavirus (<20 plaque-forming units) and a decay rate of 7% over 10 min in a room with no air changes.^{13,16}

The calculation used the following equation: $P(t) = P_0 e^{-rt}$, where $P(t)$ = the amount of some quantity at time t , P_0 = initial amount at time $t = 0$, r = the decay rate, t = time (number of periods).

Figure 1. Transmission routes: droplet, airborne, direct contact, and indirect contact. (Indirect contact: routes involving a combination of hand and surface.) Definitions of ‘droplet’ and ‘droplet nuclei’ are from Atkinson *et al.*⁵



* Transmission routes involving a combination of hand & surface = indirect contact.