

INDEX

Note: Page numbers followed by “*f*” indicate figures, and “*t*” indicate tables.

A

Antibody-dependent enhancement of infectivity (ADEI), 270–271
Antiviral selection, CoV enzymatic activities, 273 inhibitors, vacuolar acidification, 273–274 p38 MAPK, 274

C

Cap-dependent translation, 170–171
Cellular proteins, CoV replication BCoV, 150 DDX5 and DDX1, 151 hnRNP A1, 149–150 PABP, 150 TGEV, 149–150 Circulating virulent-avirulent FCoV hypothesis, 203–204 *ds*-acting RNA elements, CoV cellular proteins, role of BCoV, 150 DDX5 and DDX1, 151 hnRNP A1, 149–150 PABP, 150 TGEV, 149–150 *Coronaviridae* classification, 128 DI RNAs definition, 131–132 identification and characterization, 132 leader switching, 132 genome replication and transcription 3'-5' exoribonuclease, 129–130 ORF, 131 RdRp, 129–130 Sg minus-strand RNAs, 130–131 transcription-regulating sequences (TRSs), 130–131 nsp, 128–129 ORF1a and ORF1b, 128–129 RNA elements and genome packaging, 148–149

subgenomic messenger RNAs (sg mRNAs), 128–129
3'-terminal functional roles, 145–148 structural features, 142–145
5'-terminal functional roles, 136–141 structural features, 133–136 zoonotic coronaviruses, 128
Coronavirus (CoV)
eIF2 α phosphorylation activation, 182–183 ER stress and unfolded protein response, 180–181 genome organization and gene expression strategy discontinuous extension, 170 eIF4F and the 43S preinitiation complex, 169–170 ORFs, 169 subgenomic minus-strand RNAs, 170 transcription regulatory sequences (TRS), 169 HCoV 229E-infected cells, 170–171, 184 ORF, 174–175 SARS-CoV and MERS-CoV, 168 m⁶A-modification, 184 MERS-CoV, 168 MHV-infected cells, 184–185 mRNA translation cap-dependent translation, 170–171 IRES-mediated translation, 175–176 leaky scanning translation mechanism, internal ORFs, 174–175 N protein-mediated enhancement, 177–178 poly(A) tail length changes, 172 ribosomal frameshift, 172–173 ribosomal shunting mechanism, 174 structural features, 184 upstream ORFs (uORF), 176

Coronavirus (CoV) (*Continued*)

- viral enzymes, 171–172
- viral UTRs, 177
- PERK activation, 182–183
- PKR activation, 182–183
- protein suppressors, translation
 - 7a protein, 180
 - eIF3f, 180
 - elongation factor 1 alpha (eEF1 α), 180
 - MERS-CoV nsp1, 179
 - SARS-CoV nsp1, 179
 - 40S ribosomal subunits, 178–179
- SARS-CoV, 168
- stress granules and processing bodies, 181–182
- structural proteins, 168–169
- translation mechanism, animal cells
 - cap-dependent translation initiation, 167–168
 - eukaryotic cells, 166–167
 - eukaryotic initiation factors (eIFs), 167
 - IRES, 167–168
- viral gene expression, 183–184
- CoV S protein. *See* Spike protein (S) and tropism changes, CoV
- Cross-species transmission, 31

D

- Defective interfering RNAs (DI RNAs)
 - definition, 131–132
 - identification and characterization, 132
 - leader switching, 132
- Dendritic cells (DCs)
 - cytokines and chemokines, 273
 - immune cells, SARS-CoV and MERS-CoV, 271, 272*t*
 - MERS-CoV interaction, 272
 - SARS-CoV, 272–273
 - virus replication, 271–272

E

- Envelope protein
 - SARS-CoV, 10
 - structural features, 10–11
 - VLP, 9–10
- Eosinophilia induction, 270–271
- Epithelial cells, 271

F

- Feline coronaviruses (FCoVs)
 - accessory genes and proteins
 - ORF 3a, 196
 - ORF7a, 196–197
 - ORF 3b, 196
 - ORF7b, 196–197
 - ORF 3c, 196
 - S and E genes, 195–196
 - sgRNA3, 196
 - TGEV, 195–196
 - discriminatory mutations, 210
 - FECV
 - seropositivity rates, 199
 - symptomless persistent infections, 199–200
 - transmission, 200
 - FIPV
 - characteristic lesions, 200–201
 - clinical forms, 201–202
 - immunosuppression, 202
 - incidence of, 203
 - incubation time, 202
 - origin of, 203–204
 - molecular pathogenesis, FECV *vs.* FIPV
 - 7a gene, 204
 - 7b gene, 204–205
 - 3c gene, 204
 - furin cleavage site, 206
 - heptad repeat 1 (HR1) region, 205–206
 - mutations, 205
 - reverse genetics
 - bacterial artificial chromosome (BAC), 207–208
 - cDNA, 208–209
 - FCoV Black sequence, 209–210
 - recombinant viruses, 207
 - targeted RNA recombination, 206–207
 - role of accessory proteins, 211
 - serotypes and cellular receptor usage
 - feline aminopeptidase N (fAPN), 198–199
 - serotype I, 197–198
 - serotype II, 198–199
 - S gene mutations, 210
 - taxonomy and genome organization, 194–195, 195*f*

Feline enteric coronavirus (FECV)

vs. FIPV
7a gene, 204
7b gene, 204–205
3c gene, 204
furin cleavage site, 206
heptad repeat 1 (HR1) region, 205–206
mutations, 205
seropositivity rates, 199
symptomless persistent infections, 199–200
transmission, 200

Feline infectious peritonitis (FIP), 204–206.

See also Feline infectious peritonitis virus (FIPV)

Feline infectious peritonitis virus (FIPV)
characteristic lesions, 200–201
clinical forms, 201–202
vs. FECV
7a gene, 204
7b gene, 204–205
3c gene, 204
furin cleavage site, 206
heptad repeat 1 (HR1) region, 205–206
mutations, 205
immunosuppression, 202
incidence of, 203
incubation time, 202
origin of, 203–204

G**Genome replication and transcription**

3'-5' exoribonuclease, 129–130
ORF, 131
RdRp, 129–130
Sg minus-strand RNAs, 130–131
transcription-regulating sequences (TRSSs), 130–131

Genomic packaging signal (PS)
conservation and lineage, 148–149
discovery, 148–149
role of, 149
TGEV, 149

H

Host gene expression, 178–179
Host translational shutoff, 181–182

Human CoV (HCoV)

229E-infected cells, 170–171, 184
IFN response
human MxA, 227–228
IFITMs, 227–228
ISG, 228
2'-5' oligoadenylate synthetase (OAS), 227–228
ORF, 174–175
SARS-CoV and MERS-CoV, 168

I**IFN-gamma, 223**

IFN-induced transmembrane (IFITMs), 227–228

IFN-lambda 1–4, 223

Inactivated whole virus, 256

Innate immunity modulators, 261–263, 261f

Interferon (IFN) response

antiviral action, human CoV
human MxA, 227–228
IFITMs, 227–228
IFN-stimulated genes (ISG), 228
2'-5' oligoadenylate synthetase (OAS), 227–228

CoV genome

genome organization, 220–221, 221f

life cycle, 221–222

RTC, 221–222

sgRNA, 221–222

induction, inhibition of

dsRNA, 228–229

IRF3, 229–231

mechanisms and factors, 229–231, 230–231*t*

MHV-A59, 229

ORF4a and ORF4b protein, 231–232
pDCs, 229

resistance, 233–234, 233*t*

signaling, inhibition of, 232–233, 232*t*

type I

induction of, 224–225

stimulated gene expression, 225–226

types and signaling pathways, 222–223

Internal mutation hypothesis, 203–204

Internal ribosome entry site

(IRES)-mediated translation, 175–176

L

Leaky scanning translation mechanism,
internal ORFs, 174–175
Live-attenuated vaccines
engineer attenuated CoVs, 258
IFN sensitivity, 259–261
innate immunity modulators, 261–263,
261*f*
MERS-CoV genes, 267–269
SARS-CoV genes, 263–267
virulence of, 258–259

M

Macrophages (MØ)
cytokines and chemokines, 273
immune cells, SARS-CoV and
MERS-CoV, 271, 272*t*
MERS-CoV interaction, 272
SARS-CoV, 272–273
virus replication, 271–272
Membrane protein
appearance and characteristics, 7–8, 7*f*
endodomains, 6
MLONG–RNP interactions, 6
virus-like particles (VLP) assembly, 6–8
Middle East respiratory syndrome
(MERS) CoV
animal model, 253
antigenic complexity, 250–251
antiviral interferon response
(*see* Interferon (IFN) response)
cryo-EM reconstruction, 2–4, 3*f*
DCs, interaction, 272–273
DPP4, 252
as drug target, 89–90
envelope protein, 10
E protein, 10–11
ExoN, 102–103
HCoV, 168
hDPP4, 253–254
HR1 mutation, 46–47
immune cells, 271, 272*t*
interactions, 151
live-attenuated vaccines, 263–269
M protein, 9–10
NiRAN, 79
N protein, 8

nsp1, 179
nsp7, 67–68, 68*f*
nsp8, 68–70
nsp9, 70–71
nsp10, 71–72
nsp12, 76–77
nsp13, 81–82, 82*f*
nsp14, 103–104
nsp15, 106–109
nsp16, 97–98, 100–101
origin, 30–31
p38 MAPK, 274
protein receptor preference, 39
proteolytic cleavage sites, 45–46
RBM, 37–38
S₁ CTR changes, 43–44
SL3, 133–135
spike protein, 12
survival time, 5
5'-terminal genome, 136–137
transmission, 252

Molecular pathogenesis, FIP, 204–206
mRNA translation, CoV
cap-dependent translation, 170–171
IRES-mediated translation, 175–176
leaky scanning translation mechanism,
internal ORFs, 174–175
N protein-mediated enhancement,
177–178
poly(A) tail length changes, 172
ribosomal frameshift, 172–173
ribosomal shunting mechanism, 174
structural features, 184
upstream ORFs (uORF), 176
viral enzymes, 171–172
viral UTRs, 177

N

Nidovirus RdRp-associated
nucleotidyltransferase (NiRAN)
domain, 78–80
Nonstructural protein (nsp) 7, 67–68, 68*f*
Nonstructural protein (nsp) 9, 70–71
Nonstructural protein (nsp)10, 71–72
Nonstructural protein (nsp) 10–13–14–16
cap structure, 91–92
interferon-stimulated genes (ISGs), 90–91
7-methylguanosine (^{m7}G), 90

- mRNA, 90–91
nsp14 N7-methyl transferase, 94–97,
 95–96^f
nsp16 2'-O-methyl transferase, 97–101,
 99–100^f
nsp13 RNA 5' triphosphatase, 92–93
RNA guanylyl transferase (GTase), 93
- Nonstructural protein (nsp)12
 characterization, 72–73
 inhibitors, 77–78
 initiation mechanism, 74–77, 75^f
NiRAN domain, 78–80
RdRp domain, 73–74
virus-encoded RdRp, 72–73
- Nonstructural protein (nsp)13
 drug target, 89–90
 functional characterization, nidovirus
 helicase, 86–89
 HEL1, 81–83, 82^f
 helicase-associated ZBD, 83–84
 nidovirus helicase structural biology,
 84–86
 superfamilies (SFs), 80–81
- Nonstructural protein (nsp) 15
 conserved domain, 108
 hexameric form, 109
 hexamerization, 109
 His residues, 110–111
 interrelated by topological permutation,
 109
 Mn²⁺ ions, 107–108
 nidovirus endoU domains, 106–107, 107^f
 recombinant forms, 107–108
 RNase A, 110
 XendoU, *Xenopus laevis*, 106–107, 107^f
- Nonstructural protein (nsp) 8 and nsp7–nsp8
 complexes, 68–70
- Nonstructural protein (nsp) 14 ExoN
 DEDD exonucleases, 102
 genome size and ExoN conservation,
 102–103
 metal ion-coordinating residues, 105–106
 Mg²⁺ or Mn²⁺ ions, 104
 N7-MTase domain, 103–104
 nsp10–nsp14 complexes, 104–105
 SARS-CoV, 102–103
 zinc finger, 105
- N protein-mediated enhancement, 177–178
- Nucleoprotein
 conservation of, 9, 9^f
 C-terminal domain (CTD), 8–9
 kinases, 8
 N-terminal domain, 8–9
- P**
- Pathogen recognition receptors (PRRs),
 224–225
- Protein kinase, RNA activated (PKR),
 224–225
- Protein suppressors, CoV
 7a protein, 180
 eIF3f, 180
 elongation factor 1 alpha (eEF1 α), 180
 MERS-CoV nsp1, 179
 SARS-CoV nsp1, 179
 40S ribosomal subunits, 178–179
- Proteolytic cleavage and conformational
 changes
 prefusion conformation, 40–41
 receptor mediated endocytosis, 39–40
 virus-cell fusion, 39–40
- R**
- Receptor-binding domain (RBD), 37–38
- Receptor interactions, S₁ subunit
 C-terminal region (CTR) changes, 43–44
 N-terminal region (NTR) changes,
 42–43
- Replication and transcription complex
 (RTC)
 IFN response, 221–222
 RNA synthesis and processing, 64, 112
- Reverse genetics, FCoVs
 bacterial artificial chromosome (BAC),
 207–208
 cDNA, 208–209
 FCoV Black sequence, 209–210
 recombinant viruses, 207
 targeted RNA recombination, 206–207
- Ribosomal frameshift, 172–173
- Ribosomal shunting mechanism, 174
- RIG-I-like receptors (RLRs), 225
- RNA-dependent RNA polymerase (RdRp)
 genome replication and transcription,
 129–130
 inhibitors, 77–78

- RNA-dependent RNA polymerase (RdRp)
(Continued)
initiation mechanism, 74–77, 75*f*
recombinant expression, 74
- RNA elements, CoV. *See cis*-acting RNA elements, CoV
- RNA synthesis and processing, CoVs
accessory proteins, 64–65
antibody availability, 67
cell culture models, 66–67
3C-like protease (3CL^{pro}), 61–63
core replicase domains, 61
error rate and genomic plasticity, 113
genome organization and expression strategy, 61–63, 62*f*
- nidoviruses, 61
- nsp 7, 67–68, 68*f*
- nsp 9, 70–71
- nsp 10, 71–72
- nsp 10–13–14–16
cap structure, 91–92
interferon-stimulated genes (ISGs), 90–91
7-methylguanosine (^{m7}G), 90
mRNA, 90–91
nsp14 N7-methyl transferase, 94–97, 95–96*f*
- nsp16 2'-O-methyl transferase, 97–101, 99–100*f*
- nsp13 RNA 5' triphosphatase, 92–93
- RNA guanylyl transferase (GTase), 93
- nsp12
characterization, 72–73
inhibitors, 77–78
initiation mechanism, 74–77, 75*f*
- NiRAN domain, 78–80
- RdRp domain, 73–74
- virus-encoded RdRp, 72–73
- nsp13
drug target, 89–90
functional characterization, nidovirus helicase, 86–89
HEL1, 81–83, 82*f*
helicase-associated ZBD, 83–84
nidovirus helicase structural biology, 84–86
superfamilies (SFs), 80–81
- nsp15
conserved domain, 108
hexameric form, 109
hexamerization, 109
His residues, 110–111
interrelated by topological permutation, 109
Mn²⁺ ions, 107–108
nidovirus endoU domains, 106–107, 107*f*
recombinant forms, 107–108
RNase A, 110
XendoU, *Xenopus laevis*, 106–107, 107*f*
- nsp8 and nsp7–nsp8 complexes, 68–70
- nsp 14 ExoN
DEDD exonucleases, 102
genome size and ExoN conservation, 102–103
metal ion-coordinating residues, 105–106
Mg²⁺ or Mn²⁺ ions, 104
N7-MTase domain, 103–104
nsp10–nsp14 complexes, 104–105
SARS-CoV, 102–103
zinc finger, 105
- nsp interactome, 65–66
- open reading frames (ORFs), 61–63
- papain-like proteases (PL^{pro}), 61–63
- RTC, 64, 112
- subgenomic (sg) mRNAs, 64–65
- traditional genetic methods, 112–113
- transcriptionregulatory sequences (TRSs), 65
- transmission, 60–61
- ubiquitin, 61–63
- S**
- Severe acute respiratory syndrome (SARS) CoV
eEF α , 180
ER stress, 181
HCoV, 168
host mRNA decay, 184
IRES-driven translation, 179
nsp1, 179
stimulatory structure, 173
- Spike protein
HR1 and HR2, 12–13
MERS-CoV, 12

- S1 and S2 regions, 11–12
SARS-CoV, 12
Spike protein (S) and tropism changes, CoV
 cross-species transmission, 31
 MERS-CoV, 30–31
 mutations
 FIPV, 41–42
 S₂ mutations, altered tropism, 45–47
 S₁ receptor interactions, 42–44
 origin, 30
proteolytic cleavage and conformational changes
 prefusion conformation, 40–41
 receptor mediated endocytosis, 39–40
 virus-cell fusion, 39–40
SARS-CoV, 30–31
spike–receptor interactions
 protein receptor preference, 39
 RBD, 37–38
structural proteins, 30
structure of
 S₁ subunit, 32–34, 33–36*f*
 S₂ subunit, 33–34*f*, 34–37
 trimers, 31–32
Spike–receptor interactions
 protein receptor preference, 39
 RBD, 37–38
S₁ subunit
 RBD, 37–38
 receptor interactions
 C-terminal region (CTR) changes, 43–44
 N-terminal region (NTR) changes, 42–43
 structure of, 32–34, 33–36*f*
S₂ subunit
 mutations
 altered tropism, 46–47
 proteolytic cleavage sites, changes in, 45–46
 structure of, 33–34*f*, 34–37
Subunit vaccines, 254–256
Supramolecular architecture
 evolution, structural proteins
 CTD, 15
 M and N, 13–14
 nidovirus structural proteins, 14–15, 14*f*
pleomorphic nidovirus, 15–16
VLP, 14–15
packaged host-encoded protein, 1–2
S–E–M–N genes, 2–4
structure and organization, 2–4, 3*f*
viral genome, 1–2
viral proteins
 envelope protein, 9–11
 membrane protein, 5–8
 nucleoprotein, 8–9
 spike protein, 11–13
virion structure and durability, 4–5
- T**
- 3'-terminal *cis*-acting RNA elements
functional roles
 BSL and pseudoknot, 146–147
 hypervariable region (HVR), 147
 3'-terminal poly(A) tail, 147–148
- structural features
 bulged stem-loop (BSL), 143
 hairpin-type pseudoknot (PK) structure, 143–144
 hypervariable region (HVR), 145
 MHV genome, 143–145
- TGEV and FCoV, 141–142
- 5'-terminal *cis*-acting RNA elements
functional roles
 DI RNA replication, 136–137
 SARS-CoV, 136–137
 stem-loop 3, 138–139
 stem-loop 4, 139–140
 stem-loop 5, 140–141
 stem-loops 1 and 2, 137–138
- structural features
 MHV, 133, 134*f*
 RNA structure, 135–136
 SARS-CoV, 133–135
 selective 2'-hydroxyl acylation and primer extension (SHAPE) method, 135–136
 stem-loops (SLs), 133
- Translation mechanism, CoV
 cap-dependent translation initiation, 167–168
 eukaryotic cells, 166–167

- Translation mechanism, CoV (*Continued*)
 eukaryotic initiation factors
 (eIFs), 167
 internal ribosome entry site (IRES),
 167–168
- Transmissible gastroenteritis virus (TGEV),
 195–196
- Tropism changes, S protein mutations
 canine CoVs (CCoVs), 41–42
 feline CoV (FECV), 41–42
 FIPV, 41–42
 S₂ mutations
 altered tropism, 46–47
 proteolytic cleavage sites, changes in,
 45–46
- S₁ receptor interactions
 C-terminal region (CTR) changes,
 43–44
 N-terminal region (NTR) changes,
 42–43
- Type I IFN system
 induction of, 224–225
 stimulated gene expression, 225–226
 types and signaling pathways
 IFN-gamma, 223
 IFN-lambda 1–4, 223
 ISG, 222
- U**
 Upstream ORFs (uORF), 176
- V**
 Vectored vaccines, 256–257
 Viral and cellular mRNA translation, CoV.
See mRNA translation, CoV
 Viral enzymes, 171–172
 Viral proteins, assembly and fusion
 envelope protein
 SARS-CoV, 10
 structural features, 10–11
 VLP, 9–10
 membrane protein
 appearance and characteristics,
 7–8, 7f
 endodomains, 6
 MLONG–RNP interactions, 6
 virus-like particles (VLP) assembly,
 6–8
- nucleoprotein
 conservation of, 9, 9f
 C-terminal domain (CTD), 8–9
 kinases, 8
 N-terminal domain, 8–9
- spike protein
 HR1 and HR2, 12–13
 MERS-CoV, 12
 S1 and S2 regions, 11–12
 SARS-CoV, 12
- Viral UTRs, 177
- Virulence and vaccine development, CoV
 ADEI and eosinophilia induction,
 270–271
- animal models
 CRISPR–Cas9 system, 252–254
 dipeptidyl peptidase-4
 (DPP4), 252
 HCoV-OC43, 251–252
 hDPP4, 253
 MERS-CoV, 252
 SARS-CoV, 253
- antigenic complexity, SARS- and
 MERS-CoV, 250–251
- antiviral selection
 enzymatic activities, 273
 inhibitors, vacuolar acidification,
 273–274
 p38 MAPK, 274
- B- and T-cell responses, requirement of,
 249–250
- dendritic cells (DCs)
 cytokines and chemokines, 273
 immune cells, SARS-CoV and
 MERS-CoV, 271, 272t
 MERS-CoV interaction, 272
 SARS-CoV, 272–273
 virus replication, 271–272
- epithelial cells, 271
- genome structure and protein
 composition, 248–249, 248f
- human CoVs, 271
- inactivated whole virus, 256
- live-attenuated vaccines
 engineer attenuated CoVs, 258
 IFN sensitivity, 259–261
 innate immunity modulators,
 261–263, 261f

- MERS-CoV genes, 267–269
SARS-CoV genes, 263–267
virulence of, 258–259
macrophages (MØ)
 cytokines and chemokines, 273
 immune cells, SARS-CoV
 and MERS-CoV, 271, 272*t*
MERS-CoV interaction, 272
- SARS-CoV, 272–273
virus replication, 271–272
MERS-CoV, 247–248
pathogenesis, 246–247
prevalence, 246–247
SARS-CoV, 247–248
subunit vaccines, 254–256
vectored vaccines, 256–257