



ELSEVIER

<http://www.elsevier.com/locate/jiph>

Clinical spectrum of the Middle East respiratory syndrome coronavirus (MERS-CoV)

Mikiko Senga^{a,*}, Yaseen M. Arabi^{b,c,d}, Robert A. Fowler^{e,f}

^a Department of Pandemic and Epidemic Diseases, World Health Organization, Geneva, Switzerland

^b Intensive Care Department, King Abdulaziz Medical City, Riyadh, Saudi Arabia

^c King Saud bin Abdulaziz University for Health Sciences, Riyadh, Saudi Arabia

^d King Abdullah International Medical Research Center, Riyadh, Saudi Arabia

^e Institute of Health Policy Management and Evaluation, University of Toronto, Toronto, Canada

^f Department of Critical Care Medicine and Department of Medicine, Sunnybrook Health Sciences Center, Toronto, Canada

Received 29 March 2016; accepted 2 April 2016

Introduction

Middle East respiratory syndrome coronavirus (MERS-CoV) was first recognized as a cause of severe acute respiratory infection in the Kingdom of Saudi Arabia in 2012. As of February 2, 2016, 1638 laboratory-confirmed cases have been reported to the World Health Organization, including at least 587 deaths [1]. While MERS-CoV has affected 26 countries, the majority of cases have been reported in the Middle East [2]. Transmission has most commonly been associated with nosocomial outbreaks [3–7] and is occasionally due to animal-to-human (most commonly

camel-to-human) transmission and sporadic human-to-human transmission in the community [3,8–12]. Although the number of symptomatic cases resulting from secondary and tertiary transmission in human populations appears to be limited, little is known about potential human-to-human spread among asymptomatic or minimally symptomatic individuals [3,6,11,13].

Epidemiology of patients

The majority of MERS-CoV infections have been observed in adults (98%), with a slight male predominance (65%) [14]. Older adults and those with comorbidities, such as diabetes, renal disease, respiratory disease, and immunosuppression, have been more commonly represented among those with severe illness [13,14]. Younger and previously healthy individuals appear more likely

* Corresponding author at: Department of Pandemic and Epidemic Diseases, 20 Avenue Appia, 1211 Geneva 27, Switzerland. Tel.: +41 22 791 3964; fax: +41 22 791 4878.

E-mail address: sengam@who.int (M. Senga).

<http://dx.doi.org/10.1016/j.jiph.2016.04.008>

1876-0341/© 2016 Published by Elsevier Limited on behalf of King Saud Bin Abdulaziz University for Health Sciences.

to exhibit mild symptoms or be asymptomatic [5,6,11,13,15,16]. The true incidence of symptomatic and asymptomatic MERS-CoV cases remains unclear. Mild and asymptomatic cases may be misdiagnosed, which can lead to underreporting and misclassification of the disease and may also result in a possible detection bias depending on the type of health facility/unit that the patients seek treatment from. Health care workers appear to be at an increased risk of being infected with MERS-CoV, and this is likely mediated by inadequate infection prevention and control (IPC) measures and the amplification of cases amidst nosocomial outbreaks.

Clinical characteristics

After an incubation period ranging from 2 to 14 days, the majority (>70%) of symptomatic patients present with fever, chills, cough, and dyspnea. Approximately, one-third have myalgia, malaise, and gastrointestinal symptoms [6,17]. Among those with severe illness, the most prominent clinical manifestations have been acute, rapidly progressive viral pneumonia and pneumonitis, leading to hypoxic lung injury and acute respiratory distress syndrome [3,4,13,17–21]. For those who become severely ill, the median time from the onset of symptoms to intensive care unit (ICU) admission is only 5 days [14].

Approximately, half of the known adult symptomatic patients have been admitted to an ICU, with 40–70% requiring mechanical ventilation [3–5,12,13,17–28] and 4–17% needing ‘rescue’ oxygenation or ventilation therapy [3,17,20,29], such as extracorporeal membrane oxygenation and high frequency oscillatory or prone ventilation [13,21,29]. Renal replacement therapy was required in as many as 70% of critically ill patients [3,17,18,21]. However, the incidence of renal failure may be over-estimated due to nosocomial outbreaks involving patients with pre-existing renal disease [3]. In another hospital outbreak that was not epidemiologically related, renal dysfunction was reported in only 22% of cases [4]. Circulatory shock requiring intravenous vasoactive medication has been described in as many as 92% of critically ill patients [4,21,28].

Although not commonly observed, severe neurological syndrome characterized by varying degrees of consciousness, ataxia, focal motor deficit, and bilateral hyperintense lesions has been described in 3 patients [30]. MERS-CoV has not been isolated from the cerebrospinal fluid or the brain tissue of any of these patients infected with the virus, but radiological evidence suggests that MERS-CoV

associated vasculopathy may be an underlying cause [30].

Both community- and nosocomial-acquired bacterial and viral co-infections have been reported in patients with severe illness and have been associated with increased mortality [20,24,25,31–33]. This underscores the importance of preventive measures and the need to consider a broad range of pathogens when treating patients with either suspected or confirmed MERS-CoV.

Liver inflammation, evidenced by elevated aspartate aminotransferase and alanine aminotransferase levels, is common, especially among those who are critically ill [3,17–21]. Other laboratory findings include lymphopenia and leukopenia, as well as renal injury and coagulopathy in patients with severe illness and multi-system organ dysfunction [4,20,21].

Chest computed tomography has typically revealed unilateral or bilateral infiltrates, ground-glass opacities, occasional pleural effusions, and intralobular thickening consistent with acute respiratory distress syndrome and viral pneumonitis [13,21–23].

Among all patients, the median time from symptom onset to death was 11.5 days (ranging from 4 to 298 days) [6,14]. The median length of stay in the hospital was 41 days (ranging from 8 to 96 days) [21]. Higher viral load determination in the upper respiratory tract has been associated with increased severity of the disease, admission to the ICU and increased mortality [34].

Treatment

Since there are no specific treatments that have proven to be effective for MERS-CoV, case management has largely relied on organ-supportive therapy and prevention of complications. While co-infections can be treated with specific antimicrobial and antiviral medications, potential agents to treat MERS-CoV infection have been derived from experience with non-influenza respiratory viral infections, including severe acute respiratory syndrome (SARS). Case studies have reported on the outcomes of patients treated with oral ribavirin and interferon α 2b. However, such studies are at a high risk of selection bias and confounding by indication because the inclusion of patients has largely depended on the severity of the illness. Such non-randomized allocations cannot separate the influence of many other potentially confounding conditions and co-interventions. Previous studies may also have been affected by the immortal time bias, as the patients needed to remain alive long

enough to receive a particular therapy. All of these forms of bias and confounding variables preclude any valid estimation of the true treatment effect.

Currently, a phase II, open-label, single group assignment, safety/efficacy study of convalescent plasma therapy (plasma derived from survivors of MERS-CoV infection) is underway at the King Abdulrah International Medical Research Center in Saudi Arabia, and it is expected to be completed by June 2017 [36]. A recombinant modified vaccinia virus Ankara vaccine expressing the full-length MERS-CoV spike glycoprotein has been developed for preventative purposes, and it has been endorsed by the German Center for Infection Research (DIFZ) for phase I clinical trials in humans. However, there is not yet any human data regarding the potential effectiveness of the vaccine [37,38].

Little is known about the long-term outcomes of patients with MERS-CoV. However, according to the largest cohort of confirmed MERS-CoV patients ($n=14$) who were intubated and received invasive mechanical ventilation, those ($n=5$) who recovered from the acute infection and its complications have survived to the one year follow-up period [39].

Conclusion

In the short time since MERS-CoV has been discovered, significant progress has been made in understanding the epidemiology, pathophysiology, clinical manifestations, and potential treatment options. However, many important questions remain. Although exposure to camels has been implicated in human MERS-CoV cases, all potential animal reservoirs have not yet been identified. Effective treatments and vaccines have also yet to be developed. Because of the sporadic nature of MERS-CoV outbreaks, a collaborative research effort across the most affected countries will be necessary to better assess the potential animal reservoirs, epidemiology, risk factors, and transmission of MERS-CoV to develop controlled and ideally blinded treatment evaluations. Research protocols and data collection tools should be prepared, vetted, approved, and pre-positioned across the Middle East and elsewhere to facilitate the timely implementation of research activities and close many of the remaining knowledge gaps.

Funding

No funding sources.

Competing interests

None declared.

Ethical approval

Not required.

Disclaimer

Mikiko Senga is a staff member of the World Health Organization. This author alone is responsible for the views expressed in this article, and they do not necessarily represent the decisions, policy or views of the World Health Organization.

References

- [1] WHO. Disease Outbreak News: Middle East respiratory syndrome coronavirus (MERS-CoV) – Saudi Arabia. Available at: <http://who.int/csr/don/2-february-2016-mers-saudi-arabia/en/> [accessed 26.02.16].
- [2] WHO. Middle East respiratory syndrome coronavirus (MERS-CoV). Available at: <http://who.int/emergencies/mers-cov/en/> [accessed 28.02.16].
- [3] Assiri A, McGeer A, Perl TM, Price CS, Al Rabeeah AA, Cummings DA, et al. Hospital outbreak of Middle East respiratory syndrome coronavirus. *N Engl J Med* 2013;369(5):407–16.
- [4] Al-Abdallat MM, Payne DC, Alqasrawi S, Rha B, Tohme RA, Abedi GR, et al. Hospital-associated outbreak of Middle East respiratory syndrome coronavirus: a serologic, epidemiologic, and clinical description. *Clin Infect Dis* 2014;59:1225–33.
- [5] Oboho IK, Tomczyk SM, Al-Asmari AM, Banjar AA, Al-Mugti H, Aloraini MS, et al. 2014 MERS-CoV outbreak in Jeddah – a link to health care facilities. *N Engl J Med* 2015;372(9):846–54.
- [6] Korea Centers for Disease Control Prevention. Middle East respiratory syndrome coronavirus outbreak in the Republic of Korea, 2015. *Osong Public Health Res Perspect* 2015;6(4):269–78.
- [7] Park HY, Lee EJ, Ryu YW, Kim Y, Kim H, Lee H, et al. Epidemiological investigation of MERS-CoV spread in a single hospital in South Korea, May to June 2015. *Euro Surveill* 2015;20(25):1–6.
- [8] Memish ZA, Cotten M, Meyer B, Watson SJ, Alshafiqi AJ, Al Rabeeah AA, et al. Human infection with MERS coronavirus after exposure to infected camels, Saudi Arabia, 2013. *Emerg Infect Dis* 2014;20(6):1012–5.
- [9] Azhar EI, El-Kafrawy SA, Farraj SA, Hassan AM, Al-Saeed MS, Hashem AM, et al. Evidence for camel-to-human transmission of MERS coronavirus. *N Engl J Med* 2014;370(26):2499–505.
- [10] Reusken CB, Farag EA, Haagmans BL, Mohran KA, Godeke G, Raj S, et al. Occupational exposure to dromedaries and risk for MERS-CoV infection, Qatar, 2013–2014. *Emerg Infect Dis* 2015;21(8):1422–5.
- [11] Drosten C, Meyer B, Muller MA, Corman VM, Al-Masri M, Hosain R, et al. Transmission of MERS-coronavirus in household contacts. *N Engl J Med* 2014;371(9):828–35.
- [12] Memish ZA, Zumla AI, Al-Hakeem RF, Al-Rabeeah AA, Stephens GM. Family cluster of Middle East respiratory syndrome coronavirus infections. *N Engl J Med* 2013;368(26):2487–94.
- [13] Who Mers-Cov Research G. State of knowledge and data gaps of Middle East respiratory syndrome coronavirus (MERS-CoV) in humans. *PLoS Curr* 2013:5.

- [14] Zumla A, Hui DS, Perlman S. Middle East respiratory syndrome. *Lancet* 2015;386(9997):995–1007.
- [15] Memish ZA, Al-Tawfiq JA, Assiri A, et al. Middle East respiratory syndrome coronavirus disease in children. *Pediatr Infect Dis J* 2014;33:904–6.
- [16] Memish ZA, Al-Tawfiq JA, Makhdoom HQ, Al-Rabeeah AA, Assiri A, Alhakeem RF, et al. Screening for Middle East respiratory syndrome coronavirus infection in hospital patients and their healthcare worker and family contacts: a prospective descriptive study. *Clin Microbiol Infect* 2014;20(5):469–74.
- [17] Assiri A, Al-Tawfiq JA, Al-Rabeeah AA, Al-Rabiah FA, Al-Hajjar S, Al-Barrak A, et al. Epidemiological, demographic, and clinical characteristics of 47 cases of Middle East respiratory syndrome coronavirus disease from Saudi Arabia: a descriptive study. *Lancet Infect Dis* 2013;13(9):752–61.
- [18] Al-Tawfiq JA, Hinedi K, Ghandour J, Khairalla H, Musleh S, Ujayli A, et al. Middle East respiratory syndrome coronavirus: a case-control study of hospitalized patients. *Clin Infect Dis* 2014;59(2):160–5.
- [19] Al-Tawfiq JA, Momattin H, Dib J, Memish ZA. Ribavirin and interferon therapy in patients infected with the Middle East respiratory syndrome coronavirus: an observational study. *Int J Infect Dis* 2014;20:42–6.
- [20] Saad M, Omrani AS, Baig K, Bahloul A, Elzein F, Matin MA, et al. Clinical aspects and outcomes of 70 patients with Middle East respiratory syndrome coronavirus infection: a single-center experience in Saudi Arabia. *Int J Infect Dis* 2014;29:301–6.
- [21] Arabi YM, Arifi AA, Balkhy HH, Najm H, Aldawood AS, Ghabashi A, et al. Clinical course and outcomes of critically ill patients with Middle East respiratory syndrome coronavirus infection. *Ann Intern Med* 2014;160(6):389–97.
- [22] Ajlan AM, Ahyad RA, Jamjoom LG, Alharthy A, Madani TA. Middle East respiratory syndrome coronavirus (MERS-CoV) infection: chest CT findings. *AJR Am J Roentgenol* 2014;203(4):782–7.
- [23] Das KM, Lee EY, Enani MA, E AlJawder S, Singh R, Bashir S, et al. CT correlation with outcomes in 15 patients with acute Middle East respiratory syndrome coronavirus. *AJR Am J Roentgenol* 2015:1–7.
- [24] Drosten C, Seilmaier M, Corman VM, Hartmann W, Scheible G, Sack S, et al. Clinical features and virological analysis of a case of Middle East respiratory syndrome coronavirus infection. *Lancet Infect Dis* 2013;13(9):745–51.
- [25] Guery B, Poissy J, el Mansouf L, Sejourne C, Ettahar N, Lemaire X, et al. Clinical features and viral diagnosis of two cases of infection with Middle East Respiratory Syndrome coronavirus: a report of nosocomial transmission. *Lancet* 2013;381(9885):2265–72.
- [26] Hijawi B, Abdallat M, Sayaydeh A, Alqasrawi S, Haddadin A, Jaarour N, et al. Novel coronavirus infections in Jordan, April 2012: epidemiological findings from a retrospective investigation. *East Mediterr Health J* 2013;19(Suppl. 1):S12–8.
- [27] Kraaij-Dirkzwager M, Timen A, Dirksen K, Gelinck L, Leyten E, Groeneveld P, et al. Middle East respiratory syndrome coronavirus (MERS-CoV) infections in two returning travellers in the Netherlands, May 2014. *Euro Surveill* 2014;19(21).
- [28] Omrani AS, Saad MM, Baig K, Bahloul A, Abdul-Matin M, Alaidaroos AY, et al. Ribavirin and interferon alfa-2a for severe Middle East respiratory syndrome coronavirus infection: a retrospective cohort study. *Lancet Infect Dis* 2014;14(11):1090–5.
- [29] Cha RH, Joh JS, Jeong I, Lee JY, Shin HS, Kim G, et al. Renal complications and their prognosis in Korean patients with Middle East respiratory syndrome-coronavirus from the central MERS-CoV designated hospital. *J Korean Med Sci* 2015;30(12):1807–14.
- [30] Arabi YM, Harthi A, Hussein J, Bouchama A, Johani S, Hajeer AH, et al. Severe neurologic syndrome associated with Middle East respiratory syndrome coronavirus (MERS-CoV). *Infection* 2015;43:495–501.
- [31] Zaki AM, van Boheemen S, Bestebroer TM, Osterhaus AD, Fouchier RA. Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J Med* 2012;367(19):1814–20.
- [32] Bermingham A, Chand MA, Brown CS, Aarons E, Tong C, Langrish C, et al. Severe respiratory illness caused by a novel coronavirus, in a patient transferred to the United Kingdom from the Middle East, September 2012. *Euro Surveill* 2012;17(40):20290.
- [33] Health Protection Agency UKNCIt. Evidence of person-to-person transmission within a family cluster of novel coronavirus infections, United Kingdom, February 2013. *Euro Surveill* 2013;18(11):20427.
- [34] Feikin DR, Alraddadi B, Qutub M, Shabouni O, Curns A, Oboho IK, et al. Association of higher MERS-CoV virus load with severe disease and death, Saudi Arabia, 2014. *Emerg Infect Dis* 2015;21(11):2029–35.
- [35] Arabi YM. Anti-MERS-COV Convalescent Plasma Therapy. *ClinicalTrials.gov Identifier: NCT02190799*.
- [36] Volz A, Kupke A, Song F, Jany S, Fux R, Shams-Eldin H, et al. Protective efficacy of recombinant modified vaccinia virus ankara delivering Middle East respiratory syndrome coronavirus spike glycoprotein. *J Virol* 2015;89(16):8651–6.
- [37] Ludwig-Maximilians-Universität. Candidate vaccine gears up for clinical trials. Available at: <https://www.en.uni-muenchen.de/news/newsarchiv/2015/mers-sutter.html> [accessed March 4].
- [38] Khalid I, Alraddadi BM, Dairi Y, Khalid TJ, Kadri M, Alshukairi AN, et al. Acute management and long-term survival among subjects with severe Middle East respiratory syndrome coronavirus pneumonia and ARDS. *Respir Care* 2016;61(3):340–8.

Available online at www.sciencedirect.com

ScienceDirect