

in South Korea during the care of patients with known status of MERS infection⁸ whereas there was no seroconversion case among the 443 HCP with adequate personal protective equipment during the 2015 MERS outbreak in South Korea.⁹

In summary, there was no evidence of MERS-CoV infection among the HCP who participated in the care of 30 patients in NMC although a substantial proportion of HCP reported that they experienced MERS-like symptoms during the patient care period. Our results suggest that risk of MERS acquisition among HCP is low under stringent infection control measures.

ACKNOWLEDGMENTS

We thank Yu Mi Jung for technical assistance in data analysis, and we appreciate Professor Sara Gianella for her stimulating advice during the manuscript preparation.

Financial support. NMC Research Institute (grant NMC2015-MS-03).

Potential conflicts of interest. All authors report no conflicts of interest relevant to this article.

Ji Yeon Lee, MD;¹
Gayeon Kim, MD;²
Dong-Gyun Lim, MD;³
Hyeon-Gun Jee, PhD;³
Yunyoung Jang, RN;⁴
Joon-Sung Joh, MD;¹
Ina Jeong, MD;¹
Yeonjae Kim, MD;²
Eunhee Kim, RN;⁴
Bum Sik Chin, MD, PhD²

Affiliations: 1. Division of Pulmonary and Critical Care Medicine, Department of Internal Medicine, National Medical Center, Seoul, Republic of Korea; 2. Center for Infectious Diseases, National Medical Center, Seoul, Republic of Korea; 3. Center for Chronic Diseases, Research Institute, National Medical Center, Seoul, Republic of Korea; 4. Infection Control Unit, National Medical Center, Seoul, Republic of Korea.

Address correspondence to Bum Sik Chin, MD, PhD, Center for Infectious Diseases, National Medical Center, 245, Euljiro, Jung-gu, Seoul 100-799, Republic of Korea (moberr@nmc.or.kr).

Received July 7, 2016; accepted August 25, 2016; electronically published October 4, 2016.

Infect. Control Hosp. Epidemiol. 2016;37:1513–1514

© 2016 by The Society for Healthcare Epidemiology of America. All rights reserved. 0899-823X/2016/3712-0021. DOI: 10.1017/ice.2016.221

REFERENCES

1. Korea Centers for Disease Control and Prevention. Middle East respiratory syndrome coronavirus outbreak in the Republic of Korea, 2015 [published correction appears in *Osong Public Health Res Perspect* 2016;7:138]. *Osong Public Health Res Perspect* 2015;6:269–278.
2. Zumla A, Hui DS. Infection control and MERS-CoV in health-care workers. *Lancet* 2014;383:1869–1871.
3. Liu S, Chan TC, Chu YT, et al. Comparative epidemiology of human infections with Middle East respiratory syndrome

and severe acute respiratory syndrome coronaviruses among healthcare personnel. *PLOS ONE* 2016;11:e0149988.

4. Memish ZA, Al-Tawfiq JA, Makhdoom HQ, et al. Screening for Middle East respiratory syndrome coronavirus infection in hospital patients and their healthcare worker and family contacts: a prospective descriptive study. *Clin Microbiol Infect* 2014;20:469–474.
5. Yu IT, Xie ZH, Tsoi KK, et al. Why did outbreaks of severe acute respiratory syndrome occur in some hospital wards but not in others? *Clin Infect Dis* 2007;44:1017–1025.
6. Hall AJ, Tokars JI, Badreddine SA, et al. Health care worker contact with MERS patient, Saudi Arabia. *Emerg Infect Dis* 2014;20:2148–2151.
7. Oboho IK, Tomczyk SM, Al-Asmari AM, et al. 2014 MERS-CoV outbreak in Jeddah—a link to health care facilities. *N Engl J Med* 2015;372:846–854.
8. Park GE, Ko JH, Peck KR, et al. Control of an outbreak of Middle East respiratory syndrome in a tertiary hospital in Korea. *Ann Intern Med* 2016;165:87–93.
9. Kim C-J, Choi WS, Jung Y, et al. Surveillance of the MERS coronavirus infection in healthcare workers after contact with confirmed MERS patients: incidence and risk factors of MERS-CoV seropositivity [published online July 27, 2016]. *Clin Microbiol Infect* 2016.

Middle East Respiratory Syndrome Coronavirus Transmission in Dialysis Unit and Infection Control Interventions in Korea

In May 2015, a Korean was diagnosed with Middle East respiratory syndrome (MERS) coronavirus (CoV) infection after travel to the Arabian Peninsula. Within 1 month, there was the largest outbreak outside the Middle East with 186 laboratory-confirmed MERS-CoV infections resulting in 36 fatalities.¹ There were relatively few cases of MERS-CoV infection in patients requiring chronic hemodialysis. Here we report the precaution measures, hemodialysis methods, and outcomes of the contacted patients in our dialysis unit (DU).

At the time of the MERS outbreak in 2015, precaution measures were applied in our hospital. All visitors had their body temperatures monitored and were encouraged to perform hand hygiene before entering the hospital. If a person reported respiratory symptoms, such as cough, sputum, or dyspnea, a surgical mask was applied. Surgical masks were applied to all patients in the DU regardless of respiratory symptoms.

During the outbreak, 1 hemodialysis patient in our hospital was confirmed to have MERS-CoV infection. The beds in the DU are spaced approximately 1.2 meters apart without screens. During the hemodialysis sessions, 1 nurse usually cares for 6 patients, making both the patients and the healthcare providers (HCPs) vulnerable to the transmission of infectious diseases.

The patients and HCPs in the DU were isolated. Contact was classified according to the closeness and the timing of the contact. Grade 1 contact means that the person stayed within 2 meters of the index patient. Grade 2 contact means that the person stayed in the DU while the index patient was undergoing hemodialysis. Grade 3 contact means that the person stayed in the DU at different times but possibly contacted the index patient indirectly.

All hemodialysis patients were hospitalized in isolation rooms. For individual isolation hemodialysis (IIH), 25 hemodialyzers were installed in the inpatient wards. In the DU, 7–8 patients underwent cohort hemodialysis (CH) at 1 session. HCPs caring for those patients utilized contact and droplet precautions with level D personal protection equipment in accordance with World Health Organization recommendations,² including waterproof disposable gowns, gloves, face shields or goggles, and N95 masks. After each hemodialysis sessions, DU and IIH rooms were disinfected.

Sputum or throat swab specimens were obtained for real-time reverse-transcriptase polymerase chain reaction (RT-PCR) testing. RT-PCR tests were performed for surveillance at the beginning and end of the isolation, and when the patient had symptoms possibly related to MERS. At 2 and 4 weeks after exposure, blood samples were collected for serologic testing for MERS-CoV.

A total of 104 patients and 18 HCPs were exposed to MERS-CoV in the DU. There were 92 patients undergoing regular hemodialysis and 12 peritoneal dialysis patients visiting the DU. Fifty patients underwent IIH and 42 patients underwent CH. During the CH sessions, the distances between the beds were extended to 2.5 meters. The patient characteristics are summarized in Table 1.

During the isolation, 23 patients (22.1%) developed symptoms possibly related to MERS. Two patients died of aspiration pneumonia during the isolation. RT-PCR test results of 23 patients with symptoms were all negative.

The results of RT-PCR surveillance were all negative. Serologic testing was performed in 84 patients who consented to the test and the results were negative in all patients.

Because hemodialysis patients must continue hemodialysis in the DU, complete isolation is more difficult and the risk of exposure to infectious diseases is increased. As a result, when one patient is diagnosed with an infection such as MERS, there is a high risk of transmission through possible continuous exposure within the DU. To our knowledge, ours is the first case of isolating hemodialysis patients with direct or indirect contact with MERS-CoV.

In our hospital, there were no additional MERS-CoV infections among 104 dialysis-dependent patients. We believe this was because the precaution measures and isolations were effective. During the MERS outbreak, we practiced intensified precaution measures for dialysis patients, given their increased susceptibility to infection.^{3,4} In a report from Saudi Arabia in 2013, there were 9 additional MERS-CoV infections in the DU from 1 confirmed patient without precaution measures.⁵

TABLE 1. Characteristics of 104 Patients in Study of MERS-CoV in Dialysis Unit

Variable	Value
Age, mean \pm SD, y	60.5 \pm 13.7
Male sex	55 (52.9)
Type of dialysis	
HD	92 (88.5)
PD	12 (11.5)
Grade of exposure	
Grade 1	14 (13.5)
Grade 2	14 (13.5)
Grade 3	76 (73.1)
Admission	75 (72.1)
Patients with symptoms	23 (22.1)
Diarrhea	9/23 (39.1)
Fever	8/23 (34.8)
Cough	6/23 (26.1)
Sputum	6/23 (26.1)
Chills	1/23 (4.3)
Myalgia	1/23 (4.3)
Dyspnea	1/23 (4.3)
Sore throat	1/23 (4.3)
MERS PCR	
Negative at baseline	92/92 (100.0)
Negative after 2 weeks of isolation	90/90 (100.0)

NOTE. Data are no. (%) of patients unless otherwise specified. CoV, coronavirus; HD, hemodialysis; MERS, Middle East respiratory syndrome; PCR, polymerase chain reaction; PD, peritoneal dialysis.

There are some reports on the management of SARS in hemodialysis patients.^{6–8} But there are no data on the effect of precaution measures for hemodialysis patients who have contacted a SARS case. One study showed that surgical masks would be helpful for preventing transmission of SARS in patients in the DU.⁹ Though the World Health Organization recommended the N95 mask in the precaution measures for MERS,² it can cause physiological stress. One Taiwanese study showed that wearing the N95 mask for 4 hours during hemodialysis significantly reduced P_aO_2 and increased adverse respiratory outcomes in hemodialysis patients.⁹

With a large number of patients exposed to an infection in the DU, isolation and IIH are practically impossible owing to limited space as well as device and HCP availability. We isolated and classified all patients. IIH was performed for patients with grade 1 exposure or those with symptoms possibly related to MERS. CH was performed for patients with grade 2 or 3 exposure and no symptoms possibly related to MERS. Because fewer patients stayed in the DU, we could maintain more space between the patients during CH, reducing the possibility of transmission.

IIH and CH were performed to prevent further transmission of MERS. Because there was no further infection, we could not determine whether this measure was effective or not.

During a MERS outbreak, surgical masks, appropriate hand hygiene, and body temperature monitoring would be useful as precaution measures for hemodialysis patients. In the case of confirmed MERS in the DU, IIH and CH would be the means of maximum isolation minimizing possible secondary transmission with limited facilities and manpower.

ACKNOWLEDGMENTS

Financial support. None reported.

Potential conflicts of interest. All authors report no conflicts of interest relevant to this article.

Soo-youn Moon, MD;
Jun Seong Son, MD, PhD;
Yu Ho Lee, MD;
Young-Tae Kwak, MD;
Ho Yeon Chung, MD;
Hyung-Lae Lee, MD;
Woo-In Lee, MD;
Sang Ho Lee, MD;
Ju-Young Moon, MD;
Yang-Gyun Kim, MD

Affiliations: Division of Infectious Diseases, Kyung Hee University Hospital at Gangdong, Seoul, Korea.

Address correspondence to Jun Seong Son, MD, PhD, Division of Infectious Diseases, Kyung Hee University Hospital at Gangdong, 892, Dongnam-ro, Gangdong-gu, Seoul, 134-727, Korea (isonjs@naver.com).

Presented in part: American Society of Microbiology Microbe 2016; Boston, Massachusetts; June 17, 2016.

Received July 1, 2016; accepted August 4, 2016; electronically published August 30, 2016.

Infect Control Hosp Epidemiol 2016;37:1514–1516

© 2016 by The Society for Healthcare Epidemiology of America. All rights reserved. 0899-823X/2016/3712-0022. DOI: 10.1017/ice.2016.201

REFERENCES

- Oh MD, Choe PG, Oh HS, et al. Middle East respiratory syndrome coronavirus superspreading event involving 81 persons, Korea 2015. *J Korean Med Sci* 2015;30:1701–1705.
- World Health Organization (WHO). Infection prevention and control during health care for probable or confirmed cases of Middle East respiratory syndrome coronavirus (MERS-CoV) infection. WHO website. http://www.who.int/csr/disease/coronavirus_infections/ipc-mers-cov/en/. Published 2015. Accessed August 14, 2016.
- Eleftheriadis T, Liakopoulos V, Leivaditis K, Antoniadis G, Stefanidis I. Infections in hemodialysis: a concise review—part 1: bacteremia and respiratory infections. *Hippokratia* 2011;15:12–17.
- Pitcher D, Rao A, Caskey F, et al. UK Renal Registry 17th annual report: chapter 12 epidemiology of reported infections amongst patients receiving dialysis for established renal failure in England in 2012 to 2013: a joint report from Public Health England and the UK Renal Registry. *Nephron* 2015;129:257–265.
- Assiri A, McGeer A, Perl TM, et al. Hospital outbreak of Middle East respiratory syndrome coronavirus. *N Engl J Med* 2013;369:407–416.
- Wong PN, Mak SK, Lo KY, et al. Clinical presentation and outcome of severe acute respiratory syndrome in dialysis patients. *Am J Kidney Dis* 2003;42:1075–1081.
- Kwan BC, Leung CB, Szeto CC, Wang AY, Li PK. Severe acute respiratory syndrome in a hemodialysis patient. *Am J Kidney Dis* 2003;42:1069–1074.
- Kwan BC, Leung CB, Szeto CC, et al. Severe acute respiratory syndrome in dialysis patients. *J Am Soc Nephrol* 2004;15:1883–1888.
- Kao TW, Huang KC, Huang YL, Tsai TJ, Hsieh BS, Wu MS. The physiological impact of wearing an N95 mask during hemodialysis as a precaution against SARS in patients with end-stage renal disease. *J Formos Med Assoc* 2004;103:624–628.

Standardizing Direct Observation for Assessing Compliance to a Daily Chlorhexidine Bathing Protocol Among Hospitalized Patients

An efficacious intervention for preventing health care-associated infections is daily bathing with chlorhexidine gluconate (CHG).¹ Consequently, many hospitals in the United States have implemented CHG bathing in their intensive care units (ICUs) and non-ICU units.² With the increasing implementation of CHG bathing in healthcare facilities, it is important to monitor compliance to ensure that CHG baths are appropriately conducted and to identify potential opportunities for improving the process.

Most studies on compliance with CHG bathing procedures have used bathing product purchasing data³ or inventory assessments as proxy measures of compliance.⁴ Direct observations of CHG bathing may be a preferred method for assessing compliance and for understanding the overall process. Direct observation is an effective method for collecting real-time, naturalistic behavioral information about a specific process,⁵ and this method is commonplace in infection prevention.

Herein, we describe our experience training observers to conduct CHG bathing observations, and we present findings from pilot observations.

METHODS

Training of New Observers

An experienced observer trained 2 new observers using a CHG bathing training manual (with components for both ICU and non-ICU CHG bathing) created by a multidisciplinary team. The training manual is available on our website (<http://cqpi.wisc.edu/1758.htm>). New observers (trainees) were given 3 days to read the training manual; then they met with the experienced observer, who reviewed each item on the data