

Contents lists available at [ScienceDirect](#)

Vaccine

journal homepage: www.elsevier.com/locate/vaccine

DNA vaccine encoding Middle East respiratory syndrome coronavirus S1 protein induces protective immune responses in mice

Hang Chi^a, Xuexing Zheng^{a,b}, Xiwen Wang^a, Chong Wang^a, Hualei Wang^{a,c}, Weiwei Gai^a, Stanley Perlman^d, Songtao Yang^{a,c,*}, Jincun Zhao^{e,*}, Xianzhu Xia^{a,c,*}

^a Key Laboratory of Jilin Province for Zoonosis Prevention and Control, Institute of Military Veterinary, Academy of Military Medical Science, Changchun, China

^b School of Public Health, Shandong University, Jinan, China

^c Jiangsu Co-innovation Center for Prevention and Control of Important Animal Infectious Diseases and Zoonoses, Yangzhou, China

^d Department of Microbiology, University of Iowa, Iowa City, IA, USA

^e State Key Laboratory of Respiratory Diseases, Guangzhou Institute of Respiratory Disease, The First Affiliated Hospital of Guangzhou Medical University, Guangzhou, China

ARTICLE INFO

Article history:

Received 10 June 2016

Received in revised form 13 February 2017

Accepted 28 February 2017

Available online xxx

Keywords:

MERS-CoV

DNA vaccine

Spike protein

ABSTRACT

The Middle East respiratory syndrome coronavirus (MERS-CoV), is an emerging pathogen that continues to cause outbreaks in the Arabian peninsula and in travelers from this region, raising the concern that a global pandemic could occur. Here, we show that a DNA vaccine encoding the first 725 amino acids (S1) of MERS-CoV spike (S) protein induces antigen-specific humoral and cellular immune responses in mice. With three immunizations, high titers of neutralizing antibodies (up to 1: 10⁴) were generated without adjuvant. DNA vaccination with the MERS-CoV S1 gene markedly increased the frequencies of antigen-specific CD4⁺ and CD8⁺ T cells secreting IFN- γ and other cytokines. Both pcDNA3.1-S1 DNA vaccine immunization and passive transfer of immune serum from pcDNA3.1-S1 vaccinated mice protected Ad5-hDPP4-transduced mice from MERS-CoV challenge. These results demonstrate that a DNA vaccine encoding MERS-CoV S1 protein induces strong protective immune responses against MERS-CoV infection.

© 2017 Published by Elsevier Ltd.

1. Introduction

Middle East respiratory syndrome (MERS)-coronavirus (MERS-CoV), an emerging zoonotic virus, is the causative agent of MERS. MERS-CoV was first identified in Saudi Arabia in 2012 and MERS cases have been reported in 27 countries since then [1,2]. As of February 10, 2017, 1905 laboratory-confirmed cases, including 677 deaths related to MERS-CoV, had been reported to WHO (~36% mortality). Several family clusters and nosocomial clusters cases have been reported, revealing the human-to-human transmissibility of MERS-CoV, and raising the concern of a MERS-CoV global pandemic [3–5]. Currently, no licensed therapeutic or vaccine is available, which highlights the need for efficient vaccines against MERS-CoV.

To date, several vaccine candidates have been developed, such as viral vector-based recombinants [6–11], subunit vaccines [12–19], DNA vaccines [20], DNA prime/protein-boost vaccines [21] and a reverse genetics-constructed recombinant coronavirus vaccine [22]. Among them, DNA vaccines present a range of unique advantages such as proper antigen protein folding, rapid design and production, cost-effectiveness, and stability at non-refrigerated temperatures for convenient storage and shipping [23]. Furthermore, it has been reported that DNA vaccines can induce both humoral and cellular immune responses against MERS-CoV and SARS-CoV infection [20,24,25].

MERS-CoV is the first lineage of *Betacoronavirus* known to infect humans [26]. The genome of MERS-CoV encodes four structural proteins – spike (S), envelope (E), membrane (M) and nucleocapsid (N) [27]. The S protein, a class I fusion protein forming protruding spikes on the virus surface, is composed of an N-terminal S1 subunit and a C-terminal S2 subunit [28]. It has been reported that MERS-CoV binds to host cell receptor dipeptidyl peptidase 4 (DPP4) through an independently folded receptor binding domain (RBD) localized within the S1 subunit [29,30]. Moreover, S protein has been identified as the most immunogenic antigen of MERS-CoV. It plays an important role in the induction of neutralizing antibody and anti-viral T-cell responses [28]. Thus, S protein is

* Corresponding authors at: Department of Virology, Institute of Military Veterinary, Academy of Military Medical Sciences, 666 Liuying West Road, Changchun, Jilin 130012, China (S. Yang and X. Xia). State Key Laboratory of Respiratory Diseases, Guangzhou Institute of Respiratory Disease, The First Affiliated Hospital of Guangzhou Medical University, Guangzhou 510120, China (J. Zhao).

E-mail addresses: yst62041@163.com (S. Yang), zhaojincun@gird.cn (J. Zhao), xiaxzh@cae.cn (X. Xia).

the major target for current vaccines development to protect against MERS [8,10,28]. However, previous studies have demonstrated that vaccines based on full-length S potentially induce harmful side effects caused by non-neutralizing epitopes [27,31]. In contrast, RBD protein-based subunit vaccines are able to induce both neutralizing antibody and anti-viral T-cell responses against MERS-CoV infection, with the additional superiority of safety [28]. Nevertheless, to improve the immunogenicity of these subunit vaccines, it has been found necessary to use an appropriate adjuvant or even adjuvant combinations, or immune enhancers (e.g., human IgG Fc), and optimized delivery routes and doses [12–17]. An ideal MERS vaccine should induce potent neutralizing antibody response without inducing harmful immune effects such as virus-enhancing antibody or immunopathology [28,32]. Based on the established background and our previous research results, we selected S1 protein as the target for our DNA vaccine development.

In the present study, we designed and constructed a DNA vaccine encoding the S1 subunit of MERS-CoV (pcDNA3.1-S1), and evaluated antigen-specific humoral and cellular immune responses induced by this DNA vaccine in mice. Further, we investigated the protective efficacy of pcDNA3.1-S1 DNA vaccine in an Ad5-hDPP4-transduced mouse model following MERS-CoV challenge. Vaccinated mice and mice receiving immune serum before infection were found to have significantly decreased virus loads in their lungs.

2. Material and methods

2.1. Mice, virus and cells

Six-to eight-week-old specific pathogen-free female BALB/c mice were purchased from the Changchun Institute of Biological Products Co., Ltd (Changchun, China) or the National Cancer Institute and Jackson Laboratories (Maine, USA). The EMC/2012 strain of MERS-CoV (passage 8, designated MERS-CoV) was kindly provided by Bart Haagmans and Ron Fouchier (Erasmus Medical

Center, Rotterdam, The Netherlands). Vero 81 cells (derived from African Green monkey kidney) [ATCC No. CCL81] were grown in DMEM (Gibco, San Diego, CA, USA) supplemented with 10% fetal bovine serum (FBS) (Gibco, San Diego, CA, USA). MERS-CoV EMC/2012 was passaged once in Vero 81 cells and titrated by plaque assay in the same cell line.

2.2. Construction of the recombinant plasmids expressing MERS-CoV spike protein

The gene sequence encoding amino acid 1–1353 (S) of the spike protein of the Al-Hasa_15_2013 strain of MERS-CoV (GenBank accession No. KF600645.1) was synthesized by Sangon Biotech Company (Shanghai, China). The synthetic full-length S, Δ CD (S without the entire cytoplasmic domain), and S1 fragment were respectively subcloned into the mammalian expression vector pcDNA3.1 (+) (Invitrogen, San Diego, CA, USA) to generate recombinant plasmid pcDNA3.1-S, pcDNA3.1- Δ CD, and pcDNA3.1-S1 (Fig. 1A). The recombinant plasmid was then amplified in *Escherichia coli* HST08 (TaKaRa, Dalian, China) and purified using the Endo-Free Plasmid Maxi Kit (QIAGEN GmbH, Shanghai, China). The recombinant plasmid was dissolved in PBS at a final concentration of 1 μ g/ μ L for *in vitro* transfection and *in vivo* animal immunization.

2.3. Western blot analysis of spike protein expression in vitro

A 6-well plate was seeded with 293T cells which were grown to 80–90% confluence. Cells were respectively transfected with the recombinant plasmids and pcDNA3.1 empty vector using Lipofectamine 3000 Transfection Reagent (Invitrogen, San Diego, CA, USA) according to the manufacturer's instructions. Cells were harvested at 48 h post-transfection. Cell lysates were prepared using RIPA Lysis buffer (Solarbio LIFE SCIENCES, Beijing, China) according to the manufacturer's instructions, then separated on a 12% polyacrylamide gel and transferred onto a 0.45 μ m nitrocellulose blotting membrane (GE Healthcare Life Sciences, Freiburg, Germany) for Western blotting analysis using mouse anti-MERS-S1 mono-




Fig. 1. Construction and verification of DNA vaccine. Schematic diagrams of the construction of DNA vaccines encoding different fragments of MERS-CoV spike protein (A). Western blot analyses of MERS-CoV spike protein expression *in vitro*. Lysates from pcDNA3.1-S, pcDNA3.1- Δ CD, pcDNA3.1-S1 transfected 293T cells (lane 1–3) and lysates from pcDNA3.1-Empty transfected 293T cells (lane 4) were incubated with mouse anti-MERS-S1 monoclonal antibodies and mouse anti- β -tubulin monoclonal antibodies (B). The schematic of the experiment (C).

clonal antibodies (Sino biologicals, Beijing, China) and mouse anti- β -tubulin monoclonal antibodies (Ray antibody biotech, Beijing, China).

2.4. Animal immunizations

Mice were randomly divided into two groups. Mice in the experimental group were injected intramuscularly (i.m.) in the quadriceps muscle with 100 μ g recombinant plasmid in 100 μ L PBS on week 0, 3, 6 (Fig. 1C). Mice in the control group received either the same volume of PBS or pcDNA3.1 empty vector at the same time points.

2.5. ELISA measurement of MERS-CoV S-specific IgG

At weeks 1, 2, 4, 5, 7 and 8 following the primary immunization, 6 mice from each group were randomly selected for collection of serum. Blood samples were collected by retro-orbital plexus puncture. Anti-MERS-S antibody levels in serum were measured by indirect ELISA using purified RBD protein (10 μ g/mL) as the coating antigen as previously described [19]. Absorbance was read at 450 nm. Values 2-fold higher than the control group were considered positive.

2.6. Plaque reduction neutralizing test

One week following the third immunization, serum samples were harvested and were 4-fold serially diluted in DMEM (Gibco, San Diego, CA, USA) and mixed 1:1 with 80 PFU MERS-CoV EMC/2012. After a 1 h incubation at 37 °C, the mixture was added to Vero 81 cells for an additional 1 h to permit absorption. Cells were then overlaid with 1.2% agarose (containing 2% FBS, DMEM). After a further incubation of 3 days, agarose plugs were removed for collection of virus. The remaining plaques were visualized by 0.1% crystal violet staining.

2.7. IFN- γ and IL-4 ELISpot assays

Two weeks following the second immunization, 3 mice from each group were randomly selected and euthanized. Spleens were harvested into a tissue culture dish and teased apart into single-cell suspensions by pressing through a 3 ml syringe. Cells were cultured in RPMI 1640 medium (Gibco, San Diego, CA, USA) containing 10% FBS (Gibco, San Diego, CA, USA), then stimulated with or without recombinant MERS-CoV RBD (10 μ g/mL). The protein was prokaryotically expressed and purified by Ni-NTA affinity chromatography (Thermo, USA). After passing through a endotoxin removal spinning column, the endotoxin level was measured to be less than 0.04 EU/ml using a gel-clot limulus amoebocyte lysate assay. Following incubation at 37 °C in 5% CO₂ for 24 h, splenocytes producing IFN- γ and IL-4 were measured using mouse enzyme-linked immunospot (ELISpot) kits (Mabtech AB, Stockholm, Sweden) according to the manufacturer's instructions. Spot-forming cells (SFCs) were enumerated by an automated ELISpot reader (AID ELISPOT reader-iSpot, AID GmbH, GER).

2.8. Intracellular cytokine staining

Two weeks following the second immunization, splenocytes from 3 mice of each group were isolated, cultured (1×10^6 cells/mL) and stimulated at 37 °C in 5% CO₂ for 6 h, as described above, in the presence of protein transport inhibitor containing monensin (BD Biosciences, Franklin, VA, USA). Cells were then labelled with equal volumes of 1:250 dilutions of anti-CD4-FITC (Clone #RM4-5) and anti-CD8-PE (Clone #53-6.7) monoclonal antibodies (BD Biosciences, Franklin, VA, USA), then fixed and permeabilized by

Fixation/Permeabilization solution (BD Biosciences, Franklin, VA, USA) and labelled for 30 min at 4 °C with equal volumes of 1:250 dilutions of anti-IFN- γ PE-Cy7 (Clone #XMG1.2) and anti-IL-4-APC (Clone # 11B11) monoclonal antibodies (BD Biosciences, Franklin, VA, USA). Labelled cells were analyzed in a FACSAria™ Cell Sorter (BD Biosciences, Franklin, VA, USA).

2.9. ELISA measurement of cytokines

Two weeks following the second immunization, splenocytes from 3 mice of each group were isolated, cultured (1×10^6 cells/mL) and stimulated as described above, then incubated at 37 °C in 5% CO₂. After 48 h, cell-free culture supernatants were harvested. Levels of IL-2, IL-4, IL-10 and IFN- γ were measured using mouse enzyme-linked immunosorbent assays (ELISA) development kits (Mabtech AB, Stockholm, Sweden) according to the manufacturer's instructions.

2.10. MERS-CoV infection of mice

Mice were sensitized to MERS-CoV infection after prior transduction with adenovirus 5 expressing human DPP4 (Ad5-hDPP4) as previously described [33]. One week following the third immunization, DNA vaccine immunized mice or mice given 200 μ L immune serum (harvested 1 week following the third immunization) were transduced with Ad5-hDPP4 5 days before intranasal challenge with 1×10^5 PFU MERS-CoV. Lungs from 3 mice of each group were removed into PBS at days 3 and 5 post-infection and manually homogenized. Virus titers of clarified supernatants were assayed in Vero 81 cells and expressed as PFU/g tissue.

2.11. Laboratory facilities and ethics statement

All BALB/c mice were handled in compliance with the guidelines for the Welfare and Ethics of Laboratory Animals of China, and protocols were approved by the Animal Welfare and Ethics Committee of the Veterinary Institute at the Academy of Military Medical Sciences. BALB/c mice used for the MERS-CoV challenge experiments were maintained in the animal care facility at the University of Iowa and all protocols in the related experiments were approved by the University of Iowa Institutional Animal Care and Use Committee. Experiments with the MERS-CoV EMC/2012 strain were conducted in a biosafety level 3 (BSL3) laboratory and were approved by the University of Iowa.

3. Results

3.1. Construction and verification of DNA vaccine

Recombinant plasmids expressing the different fragments (full-length S, Δ CD and S1) of MERS-CoV were obtained and verified by restriction enzyme digestion and sequencing. Expression of MERS-CoV spike protein in 293T cells respectively transfected with the above recombinant plasmids was confirmed by Western blot (Fig. 1B). The expression level of S1 protein was significantly higher than S and Δ CD. We considered that the differences in expression level had an influence on the immune response to the various constructs.

3.2. DNA vaccine-induced neutralizing antibody against MERS-CoV

Antibody responses to MERS-CoV were evaluated by indirect ELISA, and shown as end-point dilution titers. Of the three DNA vaccines constructed, pcDNA3.1-S1 DNA vaccine elicited the highest antibody titer in immunized mice (Fig. 2A) and thus was

selected for further experiments. The sera from pcDNA3.1-S1 immunized mice strongly reacted with MERS-CoV RBD protein after receiving the second and third immunizations, reaching end-point titers up to 1:1280 (Fig. 2B). As shown in Fig. 2B, no significant differences were observed between samples harvested 1 week and 2 weeks post the third immunization, indicating that the antibody response reached the plateau. To determine if the antibodies in the immune serum could neutralize MERS-CoV infection *in vitro*, a plaque reduction neutralizing assay was performed using serially diluted serum samples. The serum samples efficiently neutralized MERS-CoV infection *in vitro* even after 1:10⁴ dilution (Fig. 2C). These results demonstrate that DNA vaccine encoding MERS-CoV S1 gene induced a potent neutralizing antibody response.

3.3. DNA vaccine-induced antigen-specific cellular immune responses

After confirming that pcDNA3.1-S1 successfully induced antibody responses in mice, antigen-specific cellular immune responses were evaluated by ELISpot assays and intracellular cytokine staining (ICS) assays. Splenocytes were harvested at two weeks post the second immunization. We chose this time because the RBD-specific antibody response was first detected 1–2 weeks after the second immunization (Fig. 2B). We speculated that T cell responses were also generated at the same time. As expected, significantly more SFCs of both IFN- γ and IL-4 were detected in splenocytes from pcDNA3.1-S1 treated mice (Fig. 3A and B) than controls. The frequencies of IFN- γ -expressing CD4⁺ and CD8⁺ T cells in the mice injected with pcDNA3.1-S1 was significantly

higher after MERS RBD stimulation (Fig. 3C and D), and similar results were observed for IL-4-expressing CD4⁺ and CD8⁺ T cells (Fig. 3E and F). These results demonstrate that the pcDNA3.1-S1 DNA vaccine markedly increased the frequencies of antigen-specific CD4⁺ and CD8⁺ T cells.

3.4. DNA vaccine-enhanced splenocyte cytokine secretion

To further investigate the antigen-specific cellular immune responses induced by pcDNA3.1-S1 DNA vaccine, cytokines secreted by splenocytes were assayed by ELISA. Levels of IL-2, IL-4, IL-10 and IFN- γ of splenocytes in pcDNA3.1-S1 immunized group were all significantly higher than those in the controls (Fig. 4A–D). These data demonstrate that pcDNA3.1-S1 DNA vaccine enhanced the secretion of both type 1 cytokines such as IL-2 and IFN- γ , and type 2 cytokines such as IL-4 and IL-10 in splenocytes.

3.5. Protection of MERS-CoV infected Ad5-hDPP4-transduced mice by DNA vaccine or immune serum transfer

The Ad5-hDPP4-transduced mouse model was used to evaluate the protective immunity of the DNA vaccine and the efficacy of immune serum containing neutralizing antibodies against MERS-CoV as determined by virus load in the infected lungs. Both the pcDNA3.1-S1 DNA vaccine and immune serum from pcDNA3.1-S1 vaccinated mice accelerated virus clearance. By day 3, virus titers had decreased 1–2 logs and by day 5, virus had been cleared in both groups (Fig. 5A and B).




Fig. 2. DNA vaccine-induced neutralizing antibody against MERS-CoV. Serum samples were collected by retro-orbital plexus puncture at weeks 1, 2, 4, 5, 7 and 8. Anti-MERS-S antibody levels in serum were assessed by indirect ELISA with the purified RBD protein as the detection antigen, and shown as end-point dilution titers. The horizontal dotted line indicates limit of determination (LOD). n = 6 mice/group/time point. The ELISA titers of serum samples from pcDNA3.1-S, pcDNA3.1-S Δ CD, pcDNA3.1-S1 at weeks 2, 5 and 8. Data are shown as the means \pm SDs and were analyzed by one-way ANOVA. (*****P* < 0.0001) (A). The ELISA titers of serum samples from pcDNA3.1-S1 treated mice at the indicated time (B). Serum samples were harvested 1-week post the third immunization, and serially diluted in DMEM and mixed 1:1 with 80 PFU MERS-CoV EMC/2012. Neutralizing antibodies were measured by plaque reduction neutralizing assay. n = 3 mice/group/time point (C). Data are shown as the means \pm SDs.




Fig. 3. DNA vaccine-induced antigen-specific cellular immune responses. Splenocytes were isolated two weeks following the second immunization and stimulated with or without the purified RBD protein. The S1-specific IFN- γ and IL-4 activities in splenocytes were evaluated using commercial ELISpot kits. SFCs secreting IL-4 (A) and IFN- γ (B) were enumerated in an automated ELISpot reader. The ability of the pcDNA3.1-S1 DNA vaccine to induce IFN- γ - and IL-4-expression in antigen-specific CD4⁺ and CD8⁺ T cells was analyzed by intracellular cytokine staining. Cells were stained with combined mouse anti-CD4-FITC and anti-CD8-PE, anti-IFN- γ -PE-Cy7 and anti-IL-4-PE-Cy3 monoclonal antibodies. CD4⁺ T cells expressing IFN- γ (C) and IL-4 (D) and CD8⁺ T cells expressing IFN- γ (E) and IL-4 (F) were analyzed in a FACSAria™ Cell Sorter. $n = 3$ mice/group/time point. Data are shown as the means \pm SDs and were analyzed by unpaired Student's t test. (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$).

4. Discussion

Here, we aimed to develop a new vaccine able to elicit potent immune responses against MERS-CoV infection. Considering that




Fig. 4. DNA vaccine-enhanced splenocyte cytokine secretion. Splenocytes were isolated two weeks following the second immunization and stimulated with the purified RBD protein for 48 h. Levels of IL-2 (A), IL-4 (B), IL-10 (C) and IFN- γ (D) secreted by splenocytes were measured using commercial ELISA kits. $n = 3$ mice/group/time point. Data are shown as the means \pm SDs and were analyzed by unpaired Student's t test. (** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$).




Fig. 5. Protection of MERS-CoV infected Ad5-hDPP4-transduced mice by DNA vaccine or immune serum transfer. Mice were injected intramuscularly with 100 μ g pcDNA3.1-Empty or pcDNA3.1-S1 in 100 μ L PBS on week 0, 3, 6. Serum samples were harvested 1-week post the third immunization. DNA vaccine immunized mice (A) or mice receiving 200 μ L of immune serum one day before infection (B) were transduced with Ad5-hDPP4 and infected intranasally with 1×10^5 PFU MERS-CoV. Virus titers in the lungs were measured at the indicated time points. Titers are expressed as PFU/g tissue. $n = 3$ mice/group/time point. Data are shown as the means \pm SEM and were analyzed by unpaired Student's t test. (* $P < 0.05$).

currently no studies have compared the immunogenicity of different S gene fragments in MERS DNA vaccines, we choose three mutants of MERS-CoV S protein as antigens: full-length S, Δ CD, and extracellular domain S1. Of the three DNA vaccines (pcDNA3.1-S, pcDNA3.1- Δ CD, and pcDNA3.1-S1) constructed, pcDNA3.1-S1 DNA vaccine was selected for further study since it elicited the highest antibody titer in immunized mice and contained major neutralizing epitopes [27], which made it an effective

and safe target for MERS vaccine development. Similar to our DNA-based vaccine, an adenovirus 5 (Ad5) vector-based vaccine, Ad5-MERS-S1 expressing the MERS-CoV S1 extracellular domain, induced stronger neutralizing antibody responses when compared to the vector expressing full-length S [8]. This may be because the S1 fragment that can induce humoral immune responses more efficiently than full length S since it is soluble and can easily be taken up by B cells in lymph node follicles [34–38]. Of note, pcDNA3.1- Δ CD immunization induced a slightly higher antibody response in mice than did pcDNA3.1-S, possibly because the Δ CD mutant still contains the transmembrane region anchoring the S protein to the membrane. Previous studies have shown that partial or complete removal of the SARS-CoV S cytoplasmic domain, or removal of the transmembrane domain along with the cytoplasmic domain from a DNA vaccine candidate increased the neutralizing antibody response in mice, indicating that removal of the cytoplasmic domain may result in a more native and more functionally relevant structure *in vivo* [38]. We considered that besides the influence of expression level differences, such modifications of the MERS-CoV S protein may also be responsible for the increased generation of a neutralizing antibody response.

Our data show that the pcDNA3.1-S1 DNA vaccine induced antigen-specific immune responses (IgG production, neutralizing antibodies generation, and cytokines secretion) in mice. High levels of neutralizing antibodies were generated following three immunizations without adjuvant. Furthermore, both pcDNA3.1-S1 DNA vaccination and administration of immune serum from pcDNA3.1-S1 vaccinated mice accelerated virus clearance in the lungs, suggesting that neutralizing antibodies against MERS-CoV S1 protein were protective and the immune serum transfer did not mediate an antibody-dependent enhancement of infection in this Ad5-hDPP4-transduced mouse model (Fig. 5B). We chose 12 days post the third immunization to challenge our mice because we just had limited access to BSL-3 labs, and we speculated that since the antibody response reached the plateau and T cell response is probably at the peak at this time point as well, they would not diminish so quickly after the third immunization. However, this could be a potential limitation. Long-term protection experiments are still required to evaluate the efficacy of this vaccine.

Since the emergence of MERS in 2012, some adaptive evolution of MERS-CoV strains has been reported [39,40]. In the current study, the MERS-CoV S gene sequence from the Al-Hasa_15_2013 strain was selected for its high homology with other published strains. It is worth noting that in challenge experiments, DNA vaccine immunization protected mice infected with the MERS-CoV EMC strain, indicating that the pcDNA3.1-S1 DNA vaccine did indeed induce protective immunity against different MERS-CoV strains.

Overall, we constructed and examined a DNA vaccine encoding MERS-CoV S1 protein in this study. Our data clearly demonstrate that the pcDNA3.1-S1 DNA vaccine induced a potent and protective immune response in mice, with the vaccinated animals showing no visible signs of adverse effects. While the protective efficacy evaluation of pcDNA3.1-S1 DNA vaccine in non-human primates as well as camels must be considered in a future study, our results strongly support the use of the S1 protein of MERS-CoV for gene-based vaccine development, as an effective target able to elicit antigen-specific humoral and cellular immune responses.

Author contributions

SY, JZ and XX designed the experiments. HC, XW, JZ, CW, WG and SP performed the experiment. HC, JZ, HW and SP analyzed

the data. HC and XW wrote the manuscript. HC, XZ, JZ, CW and HW reviewed the manuscript.

Conflict of interest statement

The authors declared no conflict of interest.

Acknowledgments

This work was supported by the open project of the State Key Laboratory of Respiratory Disease, China (Grant No. 2014SKRD-001); the Municipal Healthcare Joint-Innovation Major Project of Guangzhou, China (Grant No. 201604020011 to J.Z.) and the National Institutes of Health, United States (Grant No. RO1 AI091322 and PO1 AI060699 to S.P.); and the National Science and Technology Pillar Program during the Twelfth Five-year Plan Period, China (Grant No. 2013BAD12B04).

References

- [1] Zaki AM, van Boheemen S, Bestebroer TM, Osterhaus AD, Fouchier RA. Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J Med* 2012;367(19):1814–20.
- [2] de Groot RJ, Baker SC, Baric RS, Brown CS, Drosten C, Enjuanes L, et al. Middle East respiratory syndrome coronavirus (MERS-CoV): announcement of the coronavirus study group. *J Virol* 2013;87(14):7790–2.
- [3] Assiri A, McGeer A, Perl TM, Price CS, Al Rabeeah AA, Cummings DA, et al. Hospital outbreak of Middle East respiratory syndrome coronavirus. *N Engl J Med* 2013;369(5):407–16.
- [4] Ki M. 2015 MERS outbreak in Korea: hospital-to-hospital transmission. *Epidemiol Health* 2015;37:e2015033.
- [5] Vijay R, Perlman S. Middle East respiratory syndrome and severe acute respiratory syndrome. *Curr Opin Virol* 2016;16:70–6.
- [6] Song F, Fux R, Provacia LB, Volz A, Eickmann M, Becker S, et al. Middle East respiratory syndrome coronavirus spike protein delivered by modified vaccinia virus Ankara efficiently induces virus-neutralizing antibodies. *J Virol* 2013;87(21):11950–4.
- [7] Volz A, Kupke A, Song F, Jany S, Fux R, Shams-Eldin H, et al. Protective efficacy of recombinant modified vaccinia virus Ankara delivering Middle East respiratory syndrome coronavirus spike glycoprotein. *J Virol* 2015;89(16):8651–6.
- [8] Kim E, Okada K, Kenniston T, Raj VS, AlHajri MM, Farag EA, et al. Immunogenicity of an adenoviral-based Middle East respiratory syndrome coronavirus vaccine in BALB/c mice. *Vaccine* 2014;32(45):5975–82.
- [9] Guo X, Deng Y, Chen H, Lan J, Wang W, Zou X, et al. Systemic and mucosal immunity in mice elicited by a single immunization with human adenovirus type 5 or 41 vector-based vaccines carrying the spike protein of Middle East respiratory syndrome coronavirus. *Immunology* 2015;145(4):476–84.
- [10] Malczyk AH, Kupke A, Prüfer S, Scheuplein VA, Hutzler S, Kreuz D, et al. A highly immunogenic and protective Middle East respiratory syndrome coronavirus vaccine based on a recombinant measles virus vaccine platform. *J Virol* 2015;89(22):11654–67.
- [11] Haagmans BL, van den Brand JM, Raj VS, Volz A, Wohlsein P, Smits SL, et al. An orthopoxvirus-based vaccine reduces virus excretion after MERS-CoV infection in dromedary camels. *Science* 2016;351(6268):77–81.
- [12] Ma C, Wang L, Tao X, Zhang N, Yang Y, Tseng CT, et al. Searching for an ideal vaccine candidate among different MERS coronavirus receptor-binding fragments—the importance of immunofocusing in subunit vaccine design. *Vaccine* 2014;32(46):6170–6.
- [13] Ma C, Li Y, Wang L, Zhao G, Tao X, Tseng CT, et al. Intranasal vaccination with recombinant receptor-binding domain of MERS-CoV spike protein induces much stronger local mucosal immune responses than subcutaneous immunization: implication for designing novel mucosal MERS vaccines. *Vaccine* 2014;32(18):2100–8.
- [14] Lan J, Deng Y, Chen H, Lu G, Wang W, Guo X, et al. Tailoring subunit vaccine immunity with adjuvant combinations and delivery routes using the Middle East respiratory coronavirus (MERS-CoV) receptor-binding domain as an antigen. *PLoS ONE* 2014;9(11):e112602.
- [15] Zhang N, Channappanavar R, Ma C, Wang L, Tang J, Garron T, et al. Identification of an ideal adjuvant for receptor-binding domain-based subunit vaccines against Middle East respiratory syndrome coronavirus. *Cell Mol Immunol* 2016;13(2):180–90.
- [16] Lan J, Yao Y, Deng Y, Chen H, Lu G, Wang W, et al. Recombinant receptor binding domain protein induces partial protective immunity in rhesus macaques against Middle East respiratory syndrome coronavirus challenge. *EBioMedicine* 2015;2(10):1438–46.
- [17] Tang J, Zhang N, Tao X, Zhao G, Guo Y, Tseng CT, et al. Optimization of antigen dose for a receptor-binding domain-based subunit vaccine against MERS coronavirus. *Hum Vaccin Immunother* 2015;11(5):1244–50.

- [18] Coleman CM, Liu YV, Mu H, Taylor JK, Massare M, Flyer DC, et al. Purified coronavirus spike protein nanoparticles induce coronavirus neutralizing antibodies in mice. *Vaccine* 2014;32:3169–74.
- [19] Wang C, Zheng X, Gai W, Zhao Y, Wang H, Wang H, et al. MERS-CoV virus-like particles produced in insect cells induce specific humoral and cellular immunity in rhesus macaques. *Oncotarget* 2016.
- [20] Muthumani K, Falzarano D, Reuschel EL, Tingey C, Flingai S, Villarreal DO, et al. A synthetic consensus anti-spike protein DNA vaccine induces protective immunity against Middle East respiratory syndrome coronavirus in nonhuman primates. *Sci Transl Med* 2015;7(301). 301ra132.
- [21] Wang L, Shi W, Joyce MG, Modjarrad K, Zhang Y, Leung K, et al. Evaluation of candidate vaccine approaches for MERS-CoV. *Nat Commun* 2015;6:7712.
- [22] Almazán F, DeDiego ML, Sola I, Zuñiga S, Nieto-Torres JL, Marquez-Jurado S, et al. Engineering a replication-competent, propagation-defective Middle East respiratory syndrome coronavirus as a vaccine candidate. *Mbio* 2013;4(5). e00650-13.
- [23] Liu MA. DNA vaccines: a review. *J Intern Med* 2003;253(4):402–10.
- [24] Martin JE, Louder MK, Holman LA, Gordon IJ, Enama ME, Larkin BD, et al. A SARS DNA vaccine induces neutralizing antibody and cellular immune responses in healthy adults in a Phase I clinical trial. *Vaccine* 2008;26(50):6338–43.
- [25] Lu B, Tao L, Wang T, Zheng Z, Li B, Chen Z, et al. Humoral and cellular immune responses induced by 3a DNA vaccines against severe acute respiratory syndrome (SARS) or SARS-like coronavirus in mice. *Clin Vaccine Immunol* 2009;16(1):73–7.
- [26] Banik GR, Khandaker G, Rashid H. Middle East respiratory syndrome coronavirus “MERS-CoV”: current knowledge gaps. *Paediatr Respir Rev* 2015;16(3):197–202.
- [27] Du L, Tai W, Zhou Y, Jiang S. Vaccines for the prevention against the threat of MERS-CoV. *Expert Rev Vaccines* 2016;15(9):1123–34.
- [28] Zhang N, Jiang S, Du L. Current advancements and potential strategies in the development of MERS-CoV vaccines. *Expert Rev Vaccines* 2014;13(6):761–74.
- [29] Raj VS, Mou H, Smits SL, Dekkers DH, Müller MA, Dijkman R, et al. Dipeptidyl peptidase 4 is a functional receptor for the emerging human coronavirus-EMC. *Nature* 2013;495(7440):251–4.
- [30] Gao J, Lu G, Qi J, Li Y, Wu Y, Deng Y, et al. Structure of the fusion core and inhibition of fusion by a heptad-repeat peptide derived from the S protein of MERS-CoV. *J Virol* 2013;87(24):13134–40.
- [31] Du L, He Y, Zhou Y, Liu S, Zheng BJ, Jiang S. The spike protein of SARS-CoV –a target for vaccine and therapeutic development. *Nat Rev Microbiol* 2009;7(3):226–36.
- [32] Du L, Jiang S. Middle East respiratory syndrome: current status and future prospects for vaccine development. *Expert Opin Biol Ther* 2015;15(11):1647–51.
- [33] Zhao J, Li K, Wohlford-Lenane C, Agnihothram SS, Fett C, Zhao J, et al. Rapid generation of a mouse model for Middle East respiratory syndrome. *Proc Natl Acad Sci U S A* 2014;111(13):4970–5.
- [34] Pape KA, Catron DM, Itano AA, Jenkins MK. The humoral immune response is initiated in lymph nodes by B Cells that acquire soluble antigen directly in the follicles. *Immunity* 2007;26(4):491–502.
- [35] Bajénoff M, Germain RN. B-cell follicle development remodels the conduit system and allows soluble antigen delivery to follicular dendritic cells. *Blood* 2009;114(24):4989–97.
- [36] De Becker G, Moulin V, Tielemans F, De Mattia F, Urbain J, Leo O, et al. Regulation of T helper cell differentiation *in vivo* by soluble and membrane proteins provided by antigen-presenting cells. *Eur J Immunol* 1998;28(10):3161–71.
- [37] Enriquez-Rincon F, Klaus GG. Differing effects of monoclonal anti-hapten antibodies on humoral responses to soluble or particulate antigens. *Immunology* 1984;52(1):129–36.
- [38] Yang ZY, Kong WP, Huang Y, Roberts A, Murphy BR, Subbarao K, et al. A DNA vaccine induces SARS coronavirus neutralization and protective immunity in mice. *Nature* 2004;428(6982):561–4.
- [39] Forni D, Cagliani R, Mozzi A, Pozzoli U, Al-Daghri N, Clerici M, et al. Extensive positive selection drives the evolution of nonstructural proteins in lineage C *Betacoronaviruses*. *J Virol* 2016;90(7):3627–39.
- [40] Kim DW, Kim YJ, Park SH, Yun MR, Yang JS, Kang HJ, et al. Variations in spike glycoprotein gene of MERS-CoV, South Korea, 2015. *Emerg Infect Dis* 2016;22(1):100–4.