

# Chemokine CXCL10 and Coronavirus-Induced Neurologic Disease

Dominic Skinner,<sup>1</sup> Brett S. Marro,<sup>2,\*</sup> and Thomas E. Lane<sup>1,3,4,\*</sup>

## Abstract

Chemokines (chemotactic cytokines) are involved in a wide variety of biological processes. Following microbial infection, there is often robust chemokine signaling elicited from infected cells, which contributes to both innate and adaptive immune responses that control growth of the invading pathogen. Infection of the central nervous system (CNS) by the neuroadapted John Howard Mueller (JHM) strain of mouse hepatitis virus (JHMV) provides an excellent example of how chemokines aid in host defense as well as contribute to disease. Intracranial inoculation of the CNS of susceptible mice with JHMV results in an acute encephalomyelitis characterized by widespread dissemination of virus throughout the parenchyma. Virus-specific T cells are recruited to the CNS, and control viral replication through release of antiviral cytokines and cytolytic activity. Sterile immunity is not acquired, and virus will persist primarily in white matter tracts leading to chronic neuroinflammation and demyelination. Chemokines are expressed and contribute to defense as well as chronic disease by attracting targeted populations of leukocytes to the CNS. The T cell chemoattractant chemokine CXCL10 (interferon-inducible protein 10 kDa, IP-10) is prominently expressed in both stages of disease, and serves to attract activated T and B lymphocytes expressing CXC chemokine receptor 3 (CXCR3), the receptor for CXCL10. Functional studies that have blocked expression of either CXCL10 or CXCR3 illuminate the important role of this signaling pathway in host defense and neurodegeneration in a model of viral-induced neurologic disease.

**Keywords:** CXCL10, coronavirus, CNS infection

## JHMV-Induced Acute Encephalomyelitis

**T**HE NEUROATTENUATED J2.2v-1 John Howard Mueller (JHM) strain of mouse hepatitis virus (JHMV), a well-characterized laboratory strain derived from a mAb escape mutant from the highly lethal JHM-DL virus, can cause severe encephalomyelitis and demyelination in adult mice (9). Central nervous system (CNS) infection with JHMV provides well-accepted models for (i) viral-induced encephalomyelitis, (ii) evaluating molecular and cellular mechanisms that oversee neuroinflammation, (iii) defining mechanisms of viral-induced immune-mediated demyelination, and (iv) employing this model to assess therapeutic approaches to enhance remyelination within the context of persistent viral infection of the CNS.

In response to intracranial (i.c.) infection of susceptible mice with JHMV, the virus rapidly spreads, infecting res-

ident cells of the CNS (92). Within 24 h, JHMV penetrates into the parenchyma, and infects and replicates in astrocytes, oligodendrocytes, and microglia (15,92) (Fig. 1A). Brain viral titers peak between days 5 and 7 postinfection (p.i.), but decline below level of detection by plaque assay (~100 plaque-forming unit/g tissue) between 10 and 14 days p.i. (69) (Fig. 2). Importantly, while both the innate and adaptive immune responses effectively control CNS viral replication, sterile immunity is not achieved, and viral antigen and RNA persist within the CNS (52). Viral persistence results in chronic neuroinflammation leading to an immune-mediated demyelinating disease with clinical and histologic similarities to multiple sclerosis (MS) (Fig. 3A).

In response to JHMV infection of the CNS, there is a rapid increase in the expression of proinflammatory cytokines and chemokines along with matrix-metalloproteinases


<sup>1</sup>Department of Pathology, University of Utah School of Medicine, Salt Lake City, Utah.

<sup>2</sup>Department of Molecular Biology and Biochemistry, University of California, Irvine, Irvine, California.

<sup>3</sup>Immunology, Inflammation and Infectious Disease Initiative, University of Utah School of Medicine, Salt Lake City, Utah.

<sup>4</sup>Neuroscience Initiative, University of Utah School of Medicine, Salt Lake City, Utah.

\*Present address: Department of Immunology and Microbiology, The Scripps Research Institute, La Jolla, California.




**FIG. 1.** JHMV infection of the CNS induces rapid expression of CXCL10. *In situ* hybridization showing distribution of (A) viral RNA and (B) CXCL10 mRNA in brains of MHV-infected mice. Two sequential sagittal sections of paraffin-embedded brain from infected mice at indicated time points were probed with  $^{35}\text{S}$ -labeled antisense riboprobes specific to either JHMV or CXCL10. Signal was detected by autoradiography after a 5-day exposure to film. The probes used for each section are indicated. Note the strict colocalization of CXCL10 mRNA with viral RNA at days 2 and 7 postinfection. (C) GFAP-positive astrocyte (purple) cells express CXCL10 mRNA in the CNS of JHMV-infected mice. Combined immunohistochemistry for GFAP and *in situ* hybridization for CXCL10 mRNA were performed on the brain of a mouse following infection. Astrocytes and their processes are stained purple, and are identified as being positive for CXCL10 mRNA expression at day 7 p.i. by overlaying silver grains (arrows). Original magnification,  $\times 400$  (35). CNS, central nervous system; JHMV, John Howard Mueller strain of mouse hepatitis virus; mRNA, messenger RNA; p.i., postinfection.

(MMPs) (5,28,104,105). Both  $\text{IFN-}\alpha$  and  $\text{IFN-}\beta$  are expressed early, and elegant studies by Bergmann and colleagues (30) have implicated an important role of these cytokines in host defense by demonstrating an increase in viral spread and mortality in JHMV-infected  $\text{IFNAR}^{-/-}$  mice. In addition, administration of type I interferons impedes viral spread throughout the CNS, further supporting an important role of these cytokines in host defense (58,75). Type I interferons have also been suggested to augment

MHC class I expression arguing for a role in host defense through increased antigen presentation to T cells (1).

Neutrophils, natural killer (NK) cells, and monocyte/macrophages rapidly migrate to the CNS in response to JHMV infection (Fig. 2) (51,82,97,105). Neutrophils and monocyte/macrophages contribute to the permeabilization of the blood-brain barrier (BBB) through secretion of MMPs, and this subsequently promotes infiltration of virus-specific T cells into the CNS (28,72,101,104,105). JHMV




**FIG. 2.** JHMV infection of the CNS invokes rapid infiltration of defined immune cell subsets. Cartoon depiction of immune response following i.c. infection of the CNS of susceptible C57BL/6 with JHMV. Cellular components of the innate immune response, for example, neutrophils, macrophages, and NK cells are rapidly mobilized, and migrate to the CNS and contribute to opening the blood-brain barrier and controlling viral replication. Infiltrating CD4<sup>+</sup> and CD8<sup>+</sup> T cells reduce viral titers below level of detection through IFN- $\gamma$  secretion and cytolytic activity. Neutralizing virus-specific antibody is required to suppress viral recrudescence during chronic disease. i.c., intracranial; NK, natural killer.

infection of IL-15 knockout mice, which lack functional NK cells, is able to effectively control viral replication, arguing that NK cells are not required for host defense (106).

JHMV-specific CD4<sup>+</sup> and CD8<sup>+</sup> T cells expand to viral antigens presented within draining cervical lymph nodes, and

traffic into the CNS through a permeable BBB (105). Antiviral effector mechanisms associated with viral clearance within the CNS include the elevated expression of MHC class I and MHC class II on antigen-presenting cells (APCs), after secretion of IFN- $\gamma$  by both CD4<sup>+</sup> and CD8<sup>+</sup> T cells as well as perforin-mediated cytotoxicity of astrocytes and microglia by virus-specific CD8<sup>+</sup> T cells (40,61,69). Within the context of the JHMV model, CD8<sup>+</sup> T cell expansion and antiviral effector function are enhanced through CD4<sup>+</sup> T cells (67). Further support for the role of CD4<sup>+</sup> T cells in enhancing antiviral CD8<sup>+</sup> T cell function is provided through studies in which CD4<sup>+</sup> T cells were depleted resulting in reduced CD8<sup>+</sup> T cell expression of IFN- $\gamma$  and granzyme B combined with elevated CD8<sup>+</sup> T cell apoptosis (67). These findings support earlier studies (80,103), demonstrating that CD4<sup>+</sup> T cells play a crucial role in both enhancing peripheral activation of CD8<sup>+</sup> T cells and prolonging their antiviral function within the CNS; IL-21 has been suggested to be a critical factor in controlling these specific events (67).

Oligodendrocytes infected with JHMV appear resilient to lytic effects of CD8<sup>+</sup> T cells but are able to respond to IFN- $\gamma$  secreted from virus-specific T cells and control viral replication through this mechanism (19,41,49,61). More recently, microglia have been shown to be important in host defense following JHMV infection of the CNS. Wheeler *et al.* (96) demonstrated increased morbidity/mortality associated with impaired antiviral effector responses by T cells following targeted deletion of microglia. These findings highlight that microglia are able to shape both innate and adaptive immune responses following infection with a neurotropic virus. With regard to B cells and their role in host defense following JHMV infection of the CNS, neutralizing JHMV-specific antibody is detected during chronic disease and is critical in preventing viral recrudescence (40,54,68,70) (Fig. 2).



**FIG. 3.** Persistent JHMV infection results in an immune-mediated demyelinating disease. (A) Cartoon depiction of viral persistence within the CNS and demyelination following i.c. infection of C57BL/6 mice with JHMV. Viral titers within the CNS peak between 5 and 7 days p.i., and then decline below levels of detection as a result of infiltrating virus-specific T cells. Sterile immunity is not achieved, and viral RNA/antigen can be detected out to 1 year p.i. Robust immune-mediated demyelination occurs as a result of viral persistence resulting in chronic neuroinflammation. (B) Representative *in situ* hybridization showing viral RNA (virus-specific <sup>35</sup>S-labeled antisense riboprobes) present within a spinal cord white matter tract; sequential spinal cord section stained with LFB/H&E, showing that viral persistence results in immune cell infiltration into white matter tracts accompanied by myelin damage. H&E, hematoxylin and eosin; LFB, luxol fast blue.



More recently, Perlman and colleagues have provided important insight into the functional role of regulatory T cells (Tregs) during acute JHMV-induced CNS disease (2,102). Tregs are detected within the CNS at the same time as effector CD4<sup>+</sup> T cells, indicating that the emergence and accumulation of both populations of cells are on a similar timeline following viral infection. Further, virus-specific Tregs express both IFN- $\gamma$  and IL-10 suggesting immune regulatory capacities mediated through cytokines secreted following antigen stimulation. Indeed, virus-specific Tregs dampen proliferation of virus-specific effector CD4<sup>+</sup> T cells, and depletion of Tregs increases mortality (2,102). These data suggest that within the context of acute JHMV-induced neurologic disease, Tregs limit immunopathological CNS disease without negatively impacting viral clearance (2).


### JHMV-Induced Demyelination

Infection of susceptible mice with JHMV results in a chronic immune-mediated demyelinating disease making this an excellent and well-accepted model for the human

demyelinating disease MS (6,7,36,39,50,60). Virus persists within the CNS, and *in situ* hybridization reveals viral RNA colocalizing with areas of demyelination in spinal cords of mice at day 35 p.i. with virus (Fig. 3A, B). A hallmark feature of JHMV infection of the CNS is characterized by viral spread into the spinal cord, with astrocytes and oligodendroglia being primary targets of infection and persistence. As a result, animals develop demyelinating lesions within the brain and spinal cord that are associated with clinical manifestations, including awkward gait and hindlimb paralysis.

Staining of JHMV-infected spinal cords with either luxol fast blue (LFB) or toluidine blue reveals demyelinating lesions concentrated within the anterior funiculus and lateral white matter columns of the spinal cord (Fig. 4A, B) (92). In addition, electron microscopic analysis of spinal cords from JHMV-infected mice reveals the extensive loss of myelin surrounding axons (Fig. 4C, D).

Axonopathy within the white matter tracts of the spinal cord is present as observed through the use of the SMI-32 staining or Bielschowsky's silver impregnation stain and



**FIG. 4.** JHMV-induced demyelination and axonal damage. Toluidine blue stained spinal cord sections from (A) control (day 0, D0) and (B) day 28 (D28) postinfection. Demyelination is spread throughout ventral funiculus and lateral white matter columns with notable loss of toluidine blue staining. Electron microscopy reveals extensive loss of myelin sheath at (D) day 28 p.i. compared with (C) noninfected control mice in which thick myelin sheaths are present. Boxed areas in (A, B) indicate regions analyzed for electron microscopic analysis. Focal axonal degeneration occurs in the ventral side of JHMV-infected Thy1-YFP mouse (F) spinal cords when compared with control (E) spinal cords at day 7. 2-Photon time-lapse images (times marked in min:sec) depicting absence of FAD in a noninfected Thy1-YFP spinal cord, scale bars in (E)=20  $\mu$ m (21).

initial observations suggested that this occurred concomitantly with demyelination, whereas axonal degeneration has been argued to precede oligodendrocyte dysregulation in MS (11,12). Indeed, our laboratory has recently employed 2-photon (2P) microscopy to visualize axonal damage in response to JHMV infection of Thy1-YFP mice in which medium-to-large caliber axons fluoresce yellow. Using this approach, we were able to detect axonal damage occurring as early as 7 days p.i. with virus, further supporting the notion that axonopathy can precede demyelination in this model (Fig. 4E, F) (21).

Current evidence suggests that demyelination in JHMV-infected mice is not the result of induction of an autoimmune response against neuroantigens, that is, epitope spreading, as has recently been reported to occur during Theiler's virus-induced demyelination (55,56). However, transfer of T cells from JHMV-infected animals into naïve recipients results in demyelination (95). More recently, Stohlman and colleagues (71) clearly demonstrated the presence of APCs capable of activating self-reactive (SR) T cells in JHMV-infected mice. SR T cell accumulation within the CNS of infected mice was shown to peak in mice persistently infected with JHMV; yet, these cells were not retained arguing for minimal pathologic function. In addition, a recent report has suggested that infection with mouse hepatitis virus strain A59 promotes activation of autoreactive T cells specific to myelin basic protein, although the contributions of these cells to demyelination remain to be fully defined (25).

Oligodendrocytes are an important viral reservoir during chronic JHMV-induced disease (15,92). Nonetheless, viral-induced lysis of oligodendrocytes is not considered a primary mechanism contributing to demyelination, as evidenced by JHMV infection of immunodeficient mice (lacking thymically-educated T and B lymphocytes), resulting in widespread viral replication within oligodendrocytes with very limited demyelination (97). Moreover, adoptive transfer of splenocytes from JHMV-immunized immunocompetent mice into immunodeficient mice infected i.c. with JHMV results in robust demyelination, implicating T cells as mediators of white matter damage (29,93,97).

Early studies from our laboratory demonstrated that JHMV-infected CD4<sup>-/-</sup> or CD8<sup>-/-</sup> mice develop demyelination demonstrating the importance of both T cell subsets in augmenting demyelination, yet CD4<sup>+</sup> T cells may have a more important role compared with CD8<sup>+</sup> T cells (37). CD4<sup>+</sup> T cells secrete the chemokine CCL5, a potent chemoattractant for inflammatory macrophages, and we have shown that this is a mechanism that contributes to demyelination in JHMV-infected mice (37). IFN- $\gamma$  release by CD8<sup>+</sup> T cells also contributes to macrophage migration and accumulation within the CNS that subsequently enhance demyelination (64). Activated CD4<sup>+</sup> T cells not specific to defined viral antigens, for example, bystander CD4<sup>+</sup> T cells have also been shown to contribute to demyelination in JHMV-infected mice (26). Although activated CD4<sup>+</sup> T cells are thought to amplify demyelination, in part, through recruitment of macrophages, these cells clearly exert a protective role through IFN- $\gamma$ -mediated control of viral replication and/or additional undefined mechanisms (63,81). Macrophages have been shown to be important in development of demyelinating lesions within spinal cord white matter during chronic JHMV infection

(16,97). Furthermore, antibody-mediated neutralization of the chemokine CCL5 or genetic ablation of its receptor *Ccr5* is associated with reduced macrophage infiltration correlating with a reduction in demyelination (17,18).

Adding additional insight into how T cells contribute to either disease or defense are studies from Trandem *et al.* (85), showing that adoptive transfer of Tregs to JHMV-infected mice attenuates clinical disease severity, and this is associated with dampened neuroinflammation and demyelination. Clearly, T cell infiltration into the CNS of mice persistently infected with JHMV is important in the pathogenesis of disease, although a unifying mechanism(s) attributed to how these cells contribute to disease progression as well as protection remains elusive.

### The Chemokine CXCL10 and JHMV-Induced Acute Encephalomyelitis

Chemokines, small (8–10 kDa) proteins expressed by almost all nucleated cell types, are divided into four sub-families based upon the number and spacing of conserved cysteine residues present within the amino terminus of the protein. Chemokine function is controlled through often promiscuous signaling through seven transmembrane G-protein-coupled receptors. While initially characterized as important in inflammation by targeting distinct leukocyte populations, chemokines are now considered critical mediators of a variety of biological processes, including development, tissue homeostasis, and coordinated immune responses during viral infection.

The human CXCL10/IP-10 (interferon-inducible protein 10 kDa) was originally cloned and characterized following IFN- $\gamma$  treatment of the human monocyte-like U937 in 1985 by Luster *et al.* (48). The mouse ortholog, originally dubbed cytokine response gene-2 (*crg-2*), was subsequently cloned and characterized in 1990 by Vanguri and Farber (89). The molecular and biochemical characterization of CXCL10 are outside the scope of this review, yet there are numerous articles detailing these specific biological aspects of this chemokine related to apoptosis (31,74), cell growth, and proliferation (38,53), as well as regulating angiostasis (99).

CXCL10 is a member of the non-ELR CXC chemokine along with CXCL9 and CXCL11, and these three chemokine ligands all bind to the surface receptor CXC chemokine receptor 3 (CXCR3) that is expressed on numerous different cell types. Binding of CXCL10 to CXCR3 expressed by cells of the immune system has been shown to influence migration/homing of macrophages, dendritic cells, NK cells, and activated T cell subsets to areas of inflammation (24,44,47). Initially described as potentially important in attracting T cells to psoriatic plaques (20), CXCL10 has subsequently been shown to be expressed in numerous human inflammatory diseases (24,32,44,47). In addition, CXCL10 is expressed in response to microbial infection, and is important in attracting targeted CXCR3-positive leukocytes to sites of infection that help control/eliminate the invading pathogen (88).


We became interested in host factors governing neuroinflammation in response to JHMV infection of the CNS. Previously, numerous cytokines had been shown to be increased in response to CNS infection, yet it was unclear

whether chemokines were expressed (62). Using a RNase protection assay (RPA) targeting chemokines, we demonstrated that transcripts encoding a number of different chemokines are rapidly synthesized in response to JHMV infection of the CNS (35). Of these, the chemokine CXCL10 is the predominant transcript detected at both acute and chronic stages of disease arguing for a potentially important role in both host defense and disease. *In situ* hybridization of CXCL10 transcripts revealed strict colocalization of CXCL10 messenger RNA (mRNA) transcripts with viral transcripts (Fig. 1A, B), arguing that soluble factors released from infected cells, for example, type I interferons may enhance CXCL10 expression.

We have determined that resident glial cells including astrocytes (Fig. 1C) as well as inflammatory macrophage/microglia express CXCL10 within the CNS of JHMV-infected mice (35). The early and dominant expression of CXCL10 following CNS infection by JHMV argued for a potential role as a key sentinel molecule in host defense. In support of this notion, treatment of infected mice with an anti-CXCL10-neutralizing antibody resulted in increased mortality and impaired ability to control JHMV replication that correlated with reduced levels of IFN- $\gamma$ -producing T cells within the CNS (45). Therefore, these results argued

that early expression of CXCL10 aided in host defense by attracting CXCR3-positive virus-specific T cells. These findings were further supported by subsequent studies employing JHMV infection of germline CXCL10<sup>-/-</sup> mice (13) that resulted in decreased entry of IFN- $\gamma$ -positive T cells into the CNS and reduced ability for JHMV replication. These findings indicated that blocking CXCL10 signaling, through use of either neutralizing antibody or genetic ablation, reduced activated virus-specific T cell entry into the CNS.

Interestingly, we demonstrated through flow cytometry for staining of CXCR3 and intracellular IFN- $\gamma$  following stimulation with virus-specific peptides that >90% of these virus-specific T cells expressed the CXCL10 receptor (79). However, CXCL10 neutralization selectively reduced accumulation and/or retention of virus-specific CD4<sup>+</sup> T cells to the CNS, yet exhibited a milder effect on virus-specific CD8<sup>+</sup> T cells (79). Furthermore, administration of anti-CXCR3 antibody to JHMV-infected mice reduced CD4<sup>+</sup> T cell infiltration, while CD8<sup>+</sup> T cell trafficking was not dramatically affected (78). The selective effect of anti-CXCR3 treatment on CD4<sup>+</sup> T cells was not the result of either reduced proliferation or modulation in chemokine receptor gene expression. Therefore, CXCR3 signaling has a



**FIG. 5.** MHV-CXCL10 and MHV have genetic similarity. Both viruses were generated by a recombination reaction with the thermolabile N gene deletion (designated by *asterisk*) mutant MHV-Alb4 and mRNA generated from a transcription reaction using plasmids that encode from upstream of gene 4 to the 3' end of MHV-CXCL10 and MHV. (A) The recombination reaction for MHV results in a recombinant that is genetically identical with the WT virus. MHV-CXCL10 is identical with MHV except that gene 4 is replaced by the coding sequence for CXCL10. (B) CXCL10<sup>-/-</sup> mice i.c. infected with MHV-CXCL10 exhibit 100% survival, whereas only 60% of MHV-infected mice survive to day 12 p.i. (C) Treatment of MHV-CXCL10-infected mice with an anti-CXCL10-neutralizing Ab results in significantly increased (\* $p \leq 0.05$ ) clinical scores compared with treatment with an isotype control Ab (data shown are presented as mean  $\pm$  standard error of the mean) (89). E, E protein (small envelope protein); HE, hemagglutinin-esterase; M, membrane protein; N, nucleocapsid protein; S, surface protein; UTR, 3' untranslated region; WT, wild type.

nonredundant role in T cell subset trafficking in response to viral infection, and argue that differential signals are required for trafficking and retention of virus-specific CD4<sup>+</sup> and CD8<sup>+</sup> T cells in response to JHMV CNS infection.

As an additional method to assess the importance of CXCL10 in host defense against JHMV-induced neurologic disease, we generated a recombinant virus strain of MHV capable of expressing CXCL10 (86,91). The CXCL10-expressing recombinant of MHV (MHV-CXCL10) was generated through targeted recombination using a reverse genetic approach (86). In addition, an isogenic wild-type control virus was constructed in the same manner (86). For both viruses, the exogenous gene was inserted into open reading frame (ORF)4 of the MHV-A59 parental virus (Fig. 5A). Notably, the A59 strain of MHV is capable of replicating in both the CNS and the liver following i.c. inoculation, allowing us the opportunity to explore whether the protective effects of CXCL10 are restricted to the CNS. Importantly, MHV ORF4 encodes for a nonstructural protein that is not essential for growth in tissue culture or within the mouse CNS (59,100).

Inclusion of CXCL10 into the genome of MHV did not alter virus-specific RNA synthesis or virus-specific proteins, and resulted in secretion of CXCL10 in tissue culture (86). In addition, *in vitro* growth kinetics of the CXCL10-engineered virus did not alter viral replication as compared with the isogenic control virus (86). To determine whether CXCL10 expression derived from the recombinant MHV-CXCL10 resulted in enhanced protection from disease, CXCL10<sup>-/-</sup> mice were i.c. injected with either MHV-CXCL10 or control recombinant virus, MHV. MHV infection resulted in ~40% mortality out to day 12 p.i. (Fig. 5B). In marked contrast, 100% of mice infected with MHV-CXCL10 survived until day 12 p.i. (Fig. 5B). Our previous studies indicate that localized expression of CXCL10 within virally infected tissues is important in host defense, and peripheral expression of CXCL10 in noninfected tissues does not dramatically impact the immune response. In support of this notion, CXCL10 transcripts in CXCL10<sup>-/-</sup> mice infected with MHV-CXCL10 were selectively expressed in the CNS and liver, yet transcripts were absent in CXCL10<sup>-/-</sup> mice infected with control virus. Not surprisingly, CXCL10<sup>-/-</sup> mice infected with MHV-CXCL10 showed reduced viral titers within the brains and livers, and this correlated with increased T cell accumulation within these tissues compared with control mice. This protection from viral-induced CNS and liver disease in MHV-CXCL10-infected mice was dependent upon CXCL10 derived from the recombinant virus as treatment of anti-CXCL10 virus ameliorated these effects (Fig. 5C) (91).

Given that CXCL10 signaling has been implicated in coordinating both effector T cell generation and trafficking, we wanted to determine if CXCL10 expression following JHMV infection was important in attracting T cells into the CNS or in contributing to antiviral effector function. We have determined that MHV infection of CXCL10<sup>+/+</sup> or CXCL10<sup>-/-</sup> mice results in comparable levels of T cell activation and similar numbers of virus-specific CD4<sup>+</sup> and CD8<sup>+</sup> T cells (77). We did not detect any differences in T cell proliferation, IFN- $\gamma$  secretion by virus-specific T cells, or CD8<sup>+</sup> T cell cytolytic activity. Analysis of chemokine receptor expression on CD4<sup>+</sup> and CD8<sup>+</sup> T cells obtained

from MHV-immunized CXCL10<sup>+/+</sup> and CXCL10<sup>-/-</sup> mice revealed comparable levels of CXCR3 and CCR5, which are capable of responding to ligands CXCL10 and CCL5, respectively. Adoptive transfer of splenocytes acquired from MHV-immunized CXCL10<sup>-/-</sup> mice into MHV-infected RAG1<sup>-/-</sup> mice resulted in T cell infiltration into the CNS, reduced viral burden, and demyelination comparable with RAG1<sup>-/-</sup> recipients of immune CXCL10<sup>+/+</sup> splenocytes. Collectively, these data imply that CXCL10 functions primarily as a T cell chemoattractant and does not significantly influence T cell effector response following JHMV infection (77).


While T cells clearly have an important role in controlling JHMV replication within the CNS during acute disease, antibody and B cells have a critical role in preventing viral recrudescence in persistently infected mice (54,68,70). Given the importance of antibody-secreting cells (ASCs) in suppressing re-emergence of virus, understanding how these cells migrate into the CNS is critical with regard to understanding host defense mechanisms associated with viral persistence within the CNS.

To this end, Bergmann and colleagues (65,87) have shown that ASCs express CXCR3 arguing for an important role in signaling through this receptor, and allowing these cells to migrate and accumulate within the CNS of JHMV-infected mice in which ligands CXCL9 and CXCL10 are expressed. A definitive role for CXCL10 in attracting CXCR3-positive ASCs into the CNS was confirmed through experiments in which either CXCL10<sup>-/-</sup> or CXCL9<sup>-/-</sup> mice were infected with JHMV and virus-specific antibody within the CNS evaluated (66). Phares *et al.* (66) clearly showed that ASC recruitment to the CNS of infected CXCL10<sup>-/-</sup> mice, but not CXCL9<sup>-/-</sup> mice, was dramatically impaired, thus highlighting that CXCL10 is critical for ASC recruitment. In addition to attracting ASCs to the CNS, CXCL10 was required for parenchymal entry.

### CXCL10 and JHMV-Induced Demyelination

We have previously determined that CXCL10 is associated with demyelinating lesions in mice persistently infected with MHV (Fig. 6A) (35). To determine the role of CXCL10 in contributing to demyelination in mice persistently infected with JHMV, experimental animals were treated with anti-CXCL10 or control antisera beginning on day 12 p.i., which represents a time in which demyelination is established and neurologic deficits such as hindlimb paralysis are evident. Blocking CXCL10, but not CXCL9, resulted in a dramatic reduction in clinical disease severity as animals exhibited an almost complete restoration of motor skills. Importantly, clinical disease returned when we stopped anti-CXCL10 injection, further supporting an important role of CXCL10 in contributing to clinical disease (46).

We were also able to show that the muted clinical disease in anti-CXCL10-treated mice correlated with a targeted reduction in CD4<sup>+</sup> T cells and macrophages entering the CNS as well as muted expression of IFN- $\gamma$  and the macrophage chemoattractant chemokine CCL5. Furthermore, analysis of demyelination by toluidine blue staining of spinal cord sections revealed that mice treated with control sera displayed numerous inflammatory foci and robust demyelination



**FIG. 6.** Antibody targeting of CXCL10 in mice persistently infected with JHMV reduces demyelination and increases remyelination. (A) CXCL10 mRNA transcripts were detected by *in situ* hybridization in white matter tracts of demyelinating spinal cords at day 35 p.i. (A) CXCL10-positive cells (arrows) adjacent to demyelinating lesions. (B) Spinal cord section in which the sense control probe for CRG-2 was used. No positive cells were detected. Original magnification  $\times 400$  (35). Toluidine blue-stained transverse section of an (C) anti-CXCL10-treated mouse, showing that the region of demyelination is well defined and limited to the ventral column, whereas in (D) control-treated animals lesions extend throughout the ventral and lateral columns. (E) Electron micrograph of an anti-CXCL10-treated mouse showing axons within the ventral column with thin myelin sheaths (denoted by M and arrow) surrounding axon (a) characteristic of remyelination. (F) Electron micrograph of a control mouse, showing axons (a) within the ventral column with no evidence of remyelination (46).

throughout the ventral, lateral, and dorsal columns (Fig. 6D). In contrast, demyelination was limited to the ventral column in mice treated with anti-CXCL10, supporting the observation that progression of disease is impeded (Fig. 6C). Removal of anti-CXCL10 treatment correlated with a marked increase in the severity of demyelination. Evaluation of electron micrographs from anti-CXCL10-treated and control animals showed evidence of remyelination as indicated by a thin myelin sheath surrounding axons, whereas the majority of axons in control mice were entirely demyelinated (Fig. 6E, F).


The potential role of CXCL10 in contributing to demyelination in JHMV-infected mice by attracting inflammatory T cells and macrophages into the CNS was supported by additional studies, showing that demyelination was reduced in JHMV-infected CXCL10<sup>-/-</sup> mice (13) as well as in infected animals treated with anti-CXCR3 antisera (78). In both instances, the reduction in demyelination correlated with reduced CD4<sup>+</sup> T cell and macrophage infiltration. These findings argue that blocking CXCL10 signaling results in a reduction in white matter damage by specifically inhibiting CD4<sup>+</sup> T cells gaining access to the CNS and secreting IFN- $\gamma$  that increases expression of the macrophage chemoattractant chemokine CCL5.

Interestingly, CXCL10 is increased within the cerebrospinal fluid and CNS lesions of MS patients (76), suggesting that this may be a relevant target for therapeutic intervention. Early reports using experimental autoimmune encephalomyelitis (EAE), an autoimmune-mediated neuroinflammatory disease, indicated that antibody targeting of CXCL10 blocked CD4<sup>+</sup> T cell recruitment to the CNS, resulting in diminished clinical disease severity (14). However, subsequent studies contested these findings, and demonstrated that blocking CXCL10 either made disease worse (73) or had no effect (8). More recently, Pleasure and colleagues (57) employed a unique transgenic model in which CXCL10 was selectively ablated in astrocytes, and showed dampened disease onset that correlated with reduced CD4<sup>+</sup> T cell entry and demyelination. Collectively, these diverse findings in different preclinical animal models of MS emphasize that the model employed may dictate experimental outcome when evaluating how CXCL10 expression influences chronic neuroinflammation and demyelination.

#### CXCL10 and Oligodendroglia Biology

Exposure of cultured oligodendrocyte progenitor cells (OPCs) to IFN- $\gamma$  restricts proliferation and differentiation, as





**FIG. 7. CXCL10 treatment results in OPC apoptosis.** (A) Secreted CXCL10 protein levels in supernatant from OPC cultures treated with IFN- $\gamma$  (10, 50, and 100 U/mL—48 h) were measured by ELISA. (B) Treatment of OPC cultures for 6 days with CXCL10 (10 ng/mL) showed a significant increase (\* $p$  < 0.05; \*\*\* $p$  < 0.0001) in TUNEL positive cells when compared with untreated cultures; values are expressed as mean  $\pm$  standard deviation. (C) Western blotting of proteins isolated from OPC-enriched cultures obtained from either *CXCR3*<sup>+/+</sup> or *CXCR3*<sup>-/-</sup> mice confirms that CXCR3 is expressed in WT cultures. (D) MTT assay showing cell death following 6 days of treatment of *CXCR3*<sup>-/-</sup> or WT OPC cultures with either IFN- $\gamma$  or CXCL10. Cell death is significantly (\*\* $p$  < 0.0001;  $n$  = 3 different experiments) reduced in *CXCR3*<sup>-/-</sup> cultures compared with WT cultures (82). CXCR3, CXC chemokine receptor 3; ELISA, enzyme-linked immunosorbent assay; OPC, oligodendrocyte progenitor cell.

well as triggers apoptosis (3,4,10,22,23,27,43,90,94). Moreover, overexpression of IFN- $\gamma$  within the CNS of transgenic mice results in severe behavioral deficits associated with deleterious consequences on oligodendrocytes that correlate with hypomyelination. These studies highlight the potential detrimental effect of sustained IFN- $\gamma$  expression by inflammatory leukocytes infiltrating into the CNS (34,42). During chronic inflammatory diseases such as MS, OPCs/oligodendrocytes are exposed to numerous inflammatory cytokines/chemokines that create a hostile and damaging environment. Therefore, it is important to evaluate how these cells are protected from the damaging effects of IFN- $\gamma$  signaling.

We have examined the mechanisms by which IFN- $\gamma$  mediates apoptosis of cultured OPCs, and found that IFN- $\gamma$  induces CXCL10 expression in cultured OPCs and contributes to apoptosis through a caspase-dependent mechanism (Fig. 7A, B) (84). Cultured OPCs express CXCR3, and cultures derived from either *CXCR3*<sup>+/+</sup> or *CXCR3*<sup>-/-</sup> mice exhibited reduced sensitivity to either IFN- $\gamma$ - or CXCL10-induced apoptosis (Fig. 7C, D). Moreover, signaling through the CXC chemokine receptor 2 (CXCR2) through engagement with ligand CXCL1 restricts both IFN- $\gamma$ - and CXCL10-mediated apoptosis associated with limiting cleavage of caspase 3 and increased expression of the antiapoptotic Bcl2 protein. Therefore, we would argue that

in addition to contributing to demyelinating diseases through attraction of CXCR3-bearing lymphocytes, CXCL10 may have a more direct role in white matter damage through promoting oligodendrocyte loss through induction of oligodendroglia.

This increased susceptibility of OPCs to IFN- $\gamma$ /CXCL10-induced apoptosis is not restricted to mice as we have also determined that treatment of human embryonic stem cell-derived OPCs with either IFN- $\gamma$  or CXCL10 results in increased apoptosis through a caspase 3-mediated effect (83).

### Concluding Remarks

Studies over the past 20 years from our laboratory and others have helped shape our understanding of the functional role of CXCL10 in host defense and disease in response to JHMV infection of the CNS. Using either antibody targeting or genetic silencing of CXCL10, it has been determined that early expression of CXCL10 is beneficial as it serves to attract CXCR3-positive T cells into the CNS that subsequently aid in controlling viral replication. Equally important is the demonstration that ASCs respond to CXCL10 expression in the CNS to enter the parenchyma and suppress viral replication through secretion of virus-specific antibody. Conversely, sustained expression of CXCL10 also contributes to JHMV-induced demyelination through attraction of

CD4<sup>+</sup> T cells that amplifies neuroinflammation through IFN- $\gamma$ -mediated expression of other chemokines.

Importantly, subsequent studies by other investigators have demonstrated that CXCL10 is important in host defense against other neurotropic viruses, including Herpes Simplex Virus-1 (HSV-1) (98) and West Nile Virus (WNV) (33). Although much is known about CXCL10 and how it shapes inflammation in acute and chronic diseases following viral infection of the CNS, there are undoubtedly a number of additional questions that need to be addressed with regard to how the CXCL10: CXCR3 signaling pathway influences glial biology and repair in response to viral-induced neurologic disease.

### Acknowledgments

This work was funded by the National Institutes of Health (NIH) R01NS041249 and NIH R01NS091939, as well as support from the Ray and Tye Noorda Foundation.

### Author Disclosure Statement

No competing financial interests exist.

### References

- Akwa Y, Hassett DE, Eloranta ML, *et al.* Transgenic expression of IFN-alpha in the central nervous system of mice protects against lethal neurotropic viral infection but induces inflammation and neurodegeneration. *J Immunol* 1998;161:5016–5026.
- Anghelina D, Zhao J, Trandem K, *et al.* Role of regulatory T cells in coronavirus-induced acute encephalitis. *Virology* 2009;385:358–367.
- Baerwald KD, and Popko B. Developing and mature oligodendrocytes respond differently to the immune cytokine interferon-gamma. *J Neurosci Res* 1998;52:230–239.
- Balabanov R, Strand K, Kemper A, *et al.* Suppressor of cytokine signaling 1 expression protects oligodendrocytes from the deleterious effects of interferon-gamma. *J Neurosci* 2006;26:5143–5152.
- Bender SJ, and Weiss SR. Pathogenesis of murine coronavirus in the central nervous system. *J Neuroimmune Pharmacol* 2010;5:336–354.
- Bergmann CC, Lane TE, and Stohlman SA. Coronavirus infection of the central nervous system: host-virus stand-off. *Nat Rev Microbiol* 2006;4:121–132.
- Buchmeier MJ, and Lane TE. Viral-induced neurodegenerative disease. *Curr Opin Microbiol* 1999;2:398–402.
- Byrne FR, Winters A, Brankow D, *et al.* An antibody to IP-10 is a potent antagonist of cell migration in vitro and in vivo and does not affect disease in several animal models of inflammation. *Autoimmunity* 2009;42:171–182.
- Cheever FS, Daniels JB, Pappenheimer AM, *et al.* A murine virus (Jhm) causing disseminated encephalomyelitis with extensive destruction of myelin: I. isolation and biological properties of the virus. *J Exp Med* 1949;90:181–194.
- Chew LJ, King WC, Kennedy A, *et al.* Interferon-gamma inhibits cell cycle exit in differentiating oligodendrocyte progenitor cells. *Glia* 2005;52:127–143.
- Dandekar AA, Wu GF, Pewe L, *et al.* Axonal damage is T cell mediated and occurs concomitantly with demyelination in mice infected with a neurotropic coronavirus. *J Virol* 2001;75:6115–6120.
- Das Sarma J, Kenyon LC, Hingley ST, *et al.* Mechanisms of primary axonal damage in a viral model of multiple sclerosis. *J Neurosci* 2009;29:10272–10280.
- Dufour JH, Dziejman M, Liu MT, *et al.* IFN- $\gamma$ -inducible protein 10 (IP-10; CXCL10)-deficient mice reveal a role for IP-10 in effector T cell generation and trafficking. *J Immunol* 2002;168:3195–3204.
- Fife BT, Kennedy KJ, Paniagua MC, *et al.* CXCL10 (IFN-gamma-inducible protein-10) control of encephalitogenic CD4<sup>+</sup> T cell accumulation in the central nervous system during experimental autoimmune encephalomyelitis. *J Immunol* 2001;166:7617–7624.
- Fleming JO, Trousdale MD, el-Zaatari FA, *et al.* Pathogenicity of antigenic variants of murine coronavirus JHM selected with monoclonal antibodies. *J Virol* 1986;58:869–875.
- Fleury HJ, Sheppard RD, Bornstein MB, *et al.* Further ultrastructural observations of virus morphogenesis and myelin pathology in JHM virus encephalomyelitis. *Neuropathol Appl Neurobiol* 1980;6:165–179.
- Glass WG, Hickey MJ, Hardison JL, *et al.* Antibody targeting of the CC chemokine ligand 5 results in diminished leukocyte infiltration into the central nervous system and reduced neurologic disease in a viral model of multiple sclerosis. *J Immunol* 2004;172:4018–4025.
- Glass WG, Liu MT, Kuziel WA, *et al.* Reduced macrophage infiltration and demyelination in mice lacking the chemokine receptor CCR5 following infection with a neurotropic coronavirus. *Virology* 2001;288:8–17.
- Gonzalez JM, Bergmann CC, Ramakrishna C, *et al.* Inhibition of interferon-gamma signaling in oligodendroglia delays coronavirus clearance without altering demyelination. *Am J Pathol* 2006;168:796–804.
- Gottlieb AB, Luster AD, Posnett DN, *et al.* Detection of a gamma interferon-induced protein IP-10 in psoriatic plaques. *J Exp Med* 1988;168:941–948.
- Greenberg ML, Weinger JG, Matheu MP, *et al.* Two-photon imaging of remyelination of spinal cord axons by engrafted neural precursor cells in a viral model of multiple sclerosis. *Proc Natl Acad Sci U S A* 2014;111: E2349–E2355.
- Grinspan JB, Reeves MF, Coulaloglou MJ, *et al.* Re-entry into the cell cycle is required for bFGF-induced oligodendroglial dedifferentiation and survival. *J Neurosci Res* 1996;46:456–464.
- Grinspan JB, Stern JL, Franceschini B, *et al.* Trophic effects of basic fibroblast growth factor (bFGF) on differentiated oligodendroglia: a mechanism for regeneration of the oligodendroglial lineage. *J Neurosci Res* 1993;36: 672–680.
- Groom JR, and Luster AD. CXCR3 ligands: redundant, collaborative and antagonistic functions. *Immunol Cell Biol* 2011;89:207–215.
- Gruslin E, Moisan S, St-Pierre Y, *et al.* Transcriptome profile within the mouse central nervous system and activation of myelin-reactive T cells following murine coronavirus infection. *J Neuroimmunol* 2005;162:60–70.
- Haring JS, and Perlman S. Bystander CD4 T cells do not mediate demyelination in mice infected with a neurotropic coronavirus. *J Neuroimmunol* 2003;137:42–50.
- Horiuchi M, Itoh A, Pleasure D, *et al.* MEK-ERK signaling is involved in interferon-gamma-induced death of oligodendroglial progenitor cells. *J Biol Chem* 2006;281: 20095–20106.

28. Hosking MP, Liu L, Ransohoff RM, *et al.* A protective role for ELR+ chemokines during acute viral encephalomyelitis. *PLoS Pathog* 2009;5:e1000648.
29. Houtman JJ, and Fleming JO. Dissociation of demyelination and viral clearance in congenitally immunodeficient mice infected with murine coronavirus JHM. *J Neurovirol* 1996;2:101–110.
30. Ireland DD, Stohlman SA, Hinton DR, *et al.* Type I interferons are essential in controlling neurotropic coronavirus infection irrespective of functional CD8 T cells. *J Virol* 2008;82:300–310.
31. Janette G, Rafael G, Ellika S, *et al.* Neutralization of the chemokine CXCL10 reduces apoptosis and increases axon sprouting after spinal cord injury. *J Neurosci Res* 2006;84:724–734.
32. Karin N, and Razon H. Chemokines beyond chemoattraction: CXCL10 and its significant role in cancer and autoimmunity. *Cytokine* 2018;109:24–28.
33. Klein RS, Lin E, Zhang B, *et al.* Neuronal CXCL10 directs CD8<sup>+</sup> T-cell recruitment and control of West Nile virus encephalitis. *J Virol* 2005;79:11457–11466.
34. LaFerla FM, Sugarman MC, Lane TE, *et al.* Regional hypomyelination and dysplasia in transgenic mice with astrocyte-directed expression of interferon-gamma. *J Mol Neurosci* 2000;15:45–59.
35. Lane TE, Asensio VC, Yu N, *et al.* Dynamic regulation of alpha- and beta-chemokine expression in the central nervous system during mouse hepatitis virus-induced demyelinating disease. *J Immunol* 1998;160:970–978.
36. Lane TE, and Buchmeier MJ. Murine coronavirus infection: a paradigm for virus-induced demyelinating disease. *Trends Microbiol* 1997;5:9–14.
37. Lane TE, Liu MT, Chen BP, *et al.* A central role for CD4(+) T cells and RANTES in virus-induced central nervous system inflammation and demyelination. *J Virol* 2000;74:1415–1424.
38. Lasagni L, Francalanci M, Annunziato F, *et al.* An alternatively spliced variant of CXCR3 mediates the inhibition of endothelial cell growth induced by IP-10, Mig, and I-TAC, and acts as functional receptor for platelet factor 4. *J Exp Med* 2003;197:1537–1549.
39. Libbey JE, Lane TE, and Fujinami RS. Axonal pathology and demyelination in viral models of multiple sclerosis. *Discov Med* 2014;18:79–89.
40. Lin MT, Hinton DR, Marten NW, *et al.* Antibody prevents virus reactivation within the central nervous system. *J Immunol* 1999;162:7358–7368.
41. Lin MT, Stohlman SA, and Hinton DR. Mouse hepatitis virus is cleared from the central nervous systems of mice lacking perforin-mediated cytolysis. *J Virol* 1997;71:383–391.
42. Lin W, Kemper A, Dupree JL, *et al.* Interferon-gamma inhibits central nervous system remyelination through a process modulated by endoplasmic reticulum stress. *Brain* 2006;129:1306–1318.
43. Lin W, Kunkler PE, Harding HP, *et al.* Enhanced integrated stress response promotes myelinating oligodendrocyte survival in response to interferon-gamma. *Am J Pathol* 2008;173:1508–1517.
44. Liu L, Callahan MK, Huang D, *et al.* Chemokine receptor CXCR3: an unexpected enigma. *Curr Top Dev Biol* 2005;68:149–181.
45. Liu MT, Chen BP, Oertel P, *et al.* The T cell chemoattractant IFN-inducible protein 10 is essential in host defense against viral-induced neurologic disease. *J Immunol* 2000;165:2327–2330.
46. Liu MT, Keirstead HS, and Lane TE. Neutralization of the chemokine CXCL10 reduces inflammatory cell invasion and demyelination and improves neurological function in a viral model of multiple sclerosis. *J Immunol* 2001;167:4091–4097.
47. Luster AD. Chemokines—chemotactic cytokines that mediate inflammation. *N Engl J Med* 1998;338:436–445.
48. Luster AD, Unkeless JC, and Ravetch JV. Gamma-interferon transcriptionally regulates an early-response gene containing homology to platelet proteins. *Nature* 1985;315:672–676.
49. Malone KE, Stohlman SA, Ramakrishna C, *et al.* Induction of class I antigen processing components in oligodendroglia and microglia during viral encephalomyelitis. *Glia* 2008;56:426–435.
50. Marro BS, Blanc CA, Loring JF, *et al.* Promoting remyelination: utilizing a viral model of demyelination to assess cell-based therapies. *Expert Rev Neurother* 2014;14:1169–1179.
51. Marro BS, Grist JJ, and Lane TE. Inducible expression of CXCL1 within the central nervous system amplifies viral-induced demyelination. *J Immunol* 2016;196:1855–1864.
52. Marten NW, Stohlman SA, and Bergmann CC. Role of viral persistence in retaining CD8(+) T cells within the central nervous system. *J Virol* 2000;74:7903–7910.
53. Maru SV, Holloway KA, Flynn G, *et al.* Chemokine production and chemokine receptor expression by human glioma cells: role of CXCL10 in tumour cell proliferation. *J Neuroimmunol* 2008;199:35–45.
54. Matthews AE, Weiss SR, Shlomchik MJ, *et al.* Antibody is required for clearance of infectious murine hepatitis virus A59 from the central nervous system, but not the liver. *J Immunol* 2001;167:5254–5263.
55. McMahon EJ, Bailey SL, Castenada CV, *et al.* Epitope spreading initiates in the CNS in two mouse models of multiple sclerosis. *Nat Med* 2005;11:335–339.
56. Miller SD, Vanderlugt CL, Begolka WS, *et al.* Persistent infection with Theiler's virus leads to CNS autoimmunity through epitope spreading. *Nat Med* 1997;3:1133–1136.
57. Mills Ko E, Ma JH, Guo F, *et al.* Deletion of astroglial CXCL10 delays clinical onset but does not affect progressive axon loss in a murine autoimmune multiple sclerosis model. *J Neuroinflammation* 2014;11:105.
58. Minagawa H, Takenaka A, Mohri S, *et al.* Protective effect of recombinant murine interferon beta against mouse hepatitis virus infection. *Antiviral Res* 1987;8:85–95.
59. Ontiveros E, Kuo L, Masters PS, *et al.* Inactivation of expression of gene 4 of mouse hepatitis virus strain JHM does not affect virulence in the murine CNS. *Virology* 2001;289:230–238.
60. Pachner AR. Experimental models of multiple sclerosis. *Curr Opin Neurol* 2011;24:291–299.
61. Parra B, Hinton DR, Marten NW, *et al.* IFN-gamma is required for viral clearance from central nervous system oligodendroglia. *J Immunol* 1999;162:1641–1647.
62. Pearce BD, Hobbs MV, McGraw TS, *et al.* Cytokine induction during T-cell-mediated clearance of mouse hepatitis virus from neurons in vivo. *J Virol* 1994;68:5483–5495.
63. Pewe L, Haring J, and Perlman S. CD4 T-cell-mediated demyelination is increased in the absence of gamma in-

- terferon in mice infected with mouse hepatitis virus. *J Virol* 2002;76:7329–7333.
64. Pewe L, and Perlman S. Cutting edge: CD8 T cell-mediated demyelination is IFN-gamma dependent in mice infected with a neurotropic coronavirus. *J Immunol* 2002; 168:1547–1551.
  65. Phares TW, Marques CP, Stohlman SA, *et al.* Factors supporting intrathecal humoral responses following viral encephalomyelitis. *J Virol* 2011;85:2589–2598.
  66. Phares TW, Stohlman SA, Hinton DR, *et al.* Astrocyte-derived CXCL10 drives accumulation of antibody-secreting cells in the central nervous system during viral encephalomyelitis. *J Virol* 2013;87:3382–3392.
  67. Phares TW, Stohlman SA, Hwang M, *et al.* CD4 T cells promote CD8 T cell immunity at the priming and effector site during viral encephalitis. *J Virol* 2012;86:2416–2427.
  68. Ramakrishna C, Bergmann CC, Atkinson R, *et al.* Control of central nervous system viral persistence by neutralizing antibody. *J Virol* 2003;77:4670–4678.
  69. Ramakrishna C, Stohlman SA, Atkinson RA, *et al.* Differential regulation of primary and secondary CD8<sup>+</sup> T cells in the central nervous system. *J Immunol* 2004;173:6265–6273.
  70. Ramakrishna C, Stohlman SA, Atkinson RD, *et al.* Mechanisms of central nervous system viral persistence: the critical role of antibody and B cells. *J Immunol* 2002;168: 1204–1211.
  71. Savarin C, Bergmann CC, Gaignage M, *et al.* Self-reactive CD4<sup>+</sup> T cells activated during viral-induced demyelination do not prevent clinical recovery. *J Neuroinflammation* 2015;12:207.
  72. Savarin C, Stohlman SA, Atkinson R, *et al.* Monocytes regulate T cell migration through the glia limitans during acute viral encephalitis. *J Virol* 2010;84:4878–4888.
  73. Shosaku N, Tshikatu K, Hiroyuki Y, *et al.* Neutralization of IFN-inducible protein 10/CXCL10 exacerbates experimental autoimmune encephalomyelitis. *Eur J Immunol* 2002;32:1784–1791.
  74. Sidahmed AME, León AJ, Bosinger SE, *et al.* CXCL10 contributes to p38-mediated apoptosis in primary T lymphocytes in vitro. *Cytokine* 2012;59:433–441.
  75. Smith AL, Barthold SW, and Beck DS. Intranasally administered alpha/beta interferon prevents extension of mouse hepatitis virus, strain JHM, into the brains of BALB/cByJ mice. *Antiviral Res* 1987;8:239–245.
  76. Sørensen TL, Tani M, Jensen J, *et al.* Expression of specific chemokines and chemokine receptors in the central nervous system of multiple sclerosis patients. *J Clin Invest* 1999;103:807–815.
  77. Stiles LN, Hardison JL, Schaumburg CS, *et al.* T cell antiviral effector function is not dependent on CXCL10 following murine coronavirus infection. *J Immunol* 2006; 177:8372–8380.
  78. Stiles LN, Hosking MP, Edwards RA, *et al.* Differential roles for CXCR3 in CD4<sup>+</sup> and CD8<sup>+</sup> T cell trafficking following viral infection of the CNS. *Eur J Immunol* 2006;36:613–622.
  79. Stiles LN, Liu MT, Kane JAC, *et al.* CXCL10 and trafficking of virus-specific T cells during coronavirus demyelination. *Autoimmunity* 2009;42:484–491.
  80. Stohlman SA, Bergmann CC, Lin MT, *et al.* CTL effector function within the central nervous system requires CD4<sup>+</sup> T cells. *J Immunol* 1998;160:2896–2904.
  81. Stohlman SA, Hinton DR, Parra B, *et al.* CD4 T cells contribute to virus control and pathology following central nervous system infection with neurotropic mouse hepatitis virus. *J Virol* 2008;82:2130–2139.
  82. Thirion Gand Coutelier J-P. Production of protective gamma interferon by natural killer cells during early mouse hepatitis virus infection. *J Gen Virol* 2009;90:442–447.
  83. Tirotta E, Kirby LA, Hatch MN, *et al.* IFN-gamma-induced apoptosis of human embryonic stem cell derived oligodendrocyte progenitor cells is restricted by CXCR2 signaling. *Stem Cell Res* 2012;9:208–217.
  84. Tirotta E, Ransohoff RM, and Lane TE. CXCR2 signaling protects oligodendrocyte progenitor cells from IFN-gamma/CXCL10-mediated apoptosis. *Glia* 2011;59:1518–1528.
  85. Trandem K, Anghelina D, Zhao J, *et al.* Regulatory T cells inhibit T cell proliferation and decrease demyelination in mice chronically infected with a coronavirus. *J Immunol* 2010;184:4391–4400.
  86. Trifilo MJ, and Lane TE. The CC chemokine ligand 3 regulates CD11c<sup>+</sup>CD11b<sup>+</sup>CD8alpha<sup>-</sup> dendritic cell maturation and activation following viral infection of the central nervous system: implications for a role in T cell activation. *Virology* 2004;327:8–15.
  87. Tschen SI, Stohlman SA, Ramakrishna C, *et al.* CNS viral infection diverts homing of antibody-secreting cells from lymphoid organs to the CNS. *Eur J Immunol* 2006;36:603–612.
  88. Van Raemdonck K, Van den Steen PE, Liekens S, *et al.* CXCR3 ligands in disease and therapy. *Cytokine Growth Factor Rev* 2015;26:311–327.
  89. Vanguri P, and Farber JM. Identification of CRG-2. An interferon-inducible mRNA predicted to encode a murine monokine. *J Biol Chem* 1990;265:15049–15057.
  90. Vartanian T, Li Y, Zhao M, *et al.* Interferon-gamma-induced oligodendrocyte cell death: implications for the pathogenesis of multiple sclerosis. *Mol Med* 1995;1:732–743.
  91. Walsh KB, Edwards RA, Romero KM, *et al.* Expression of CXC chemokine ligand 10 from the mouse hepatitis virus genome results in protection from viral-induced neurological and liver disease. *J Immunol* 2007;179:1155–1165.
  92. Wang FI, Hinton DR, Gilmore W, *et al.* Sequential infection of glial cells by the murine hepatitis virus JHM strain (MHV-4) leads to a characteristic distribution of demyelination. *Lab Invest* 1992;66:744–754.
  93. Wang FI, Stohlman SA, and Fleming JO. Demyelination induced by murine hepatitis virus JHM strain (MHV-4) is immunologically mediated. *J Neuroimmunol* 1990;30:31–41.
  94. Wang Y, Ren Z, Tao D, *et al.* STAT1/IRF-1 signaling pathway mediates the injurious effect of interferon-gamma on oligodendrocyte progenitor cells. *Glia* 2010;58:195–208.
  95. Watanabe R, Wege H, and ter Meulen V. Adoptive transfer of EAE-like lesions from rats with coronavirus-induced demyelinating encephalomyelitis. *Nature* 1983; 305:150–153.
  96. Wheeler DL, Sariol A, Meyerholz DK, *et al.* Microglia are required for protection against lethal coronavirus encephalitis in mice. *J Clin Invest* 2018;128:931–943.
  97. Wu GF, and Perlman S. Macrophage infiltration, but not apoptosis, is correlated with immune-mediated demyelination following murine infection with a neurotropic coronavirus. *J Virol* 1999;73:8771–8780.
  98. Wuest TR, and Carr DJ. Dysregulation of CXCR3 signaling due to CXCL10 deficiency impairs the antiviral response to herpes simplex virus 1 infection. *J Immunol* 2008;181:7985–7993.

99. Yang J, and Richmond A. The angiostatic activity of interferon-inducible protein-10/CXCL10 in human melanoma depends on binding to CXCR3 but not to glycosaminoglycan. *Mol Ther* 2004;9:846–855.
100. Yokomori K, and Lai MM. Mouse hepatitis virus S RNA sequence reveals that nonstructural proteins ns4 and ns5a are not essential for murine coronavirus replication. *J Virol* 1991;65:5605–5608.
101. Yong VW, Zabad RK, Agrawal S, *et al.* Elevation of matrix metalloproteinases (MMPs) in multiple sclerosis and impact of immunomodulators. *J Neurol Sci* 2007;259:79–84.
102. Zhao J, Zhao J, Fett C, *et al.* IFN-gamma- and IL-10-expressing virus epitope-specific Foxp3(+) T reg cells in the central nervous system during encephalomyelitis. *J Exp Med* 2011;208:1571–1577.
103. Zhou J, Hinton DR, Stohlman SA, *et al.* Maintenance of CD8<sup>+</sup> T cells during acute viral infection of the central nervous system requires CD4<sup>+</sup> T cells but not interleukin-2. *Viral Immunol* 2005;18:162–169.
104. Zhou J, Stohlman SA, Atkinson R, *et al.* Matrix metalloproteinase expression correlates with virulence following neurotropic mouse hepatitis virus infection. *J Virol* 2002;76:7374–7384.
105. Zhou J, Stohlman SA, Hinton DR, *et al.* Neutrophils promote mononuclear cell infiltration during viral-induced encephalitis. *J Immunol* 2003;170:3331–3336.
106. Zuo J, Stohlman SA, Hoskin JB, *et al.* Mouse hepatitis virus pathogenesis in the central nervous system is independent of IL-15 and natural killer cells. *Virology* 2006;350:206–215.

Address correspondence to:

*Dr. Thomas E. Lane*

*Department of Pathology*

*University of Utah School of Medicine*

*Salt Lake City, UT 84112*

*E-mail: tom.lane@utah.path.edu*