

Accepted Manuscript

Circulation of Alphacoronavirus, Betacoronavirus and Paramyxovirus in Hipposideros bat species in Zimbabwe

Mathieu Bourgarel, Davies M. Pfukenyi, Vanina Boué, Loïc Talignani, Ngoni Chiweshe, Fodé Diop, Alexandre Caron, Gift Matope, Dorothée Missé, Florian Liégeois



PII: S1567-1348(18)30007-8
DOI: <https://doi.org/10.1016/j.meegid.2018.01.007>
Reference: MEEGID 3377
To appear in: *Infection, Genetics and Evolution*
Received date: 28 September 2017
Revised date: 8 January 2018
Accepted date: 9 January 2018

Please cite this article as: Mathieu Bourgarel, Davies M. Pfukenyi, Vanina Boué, Loïc Talignani, Ngoni Chiweshe, Fodé Diop, Alexandre Caron, Gift Matope, Dorothée Missé, Florian Liégeois, Circulation of Alphacoronavirus, Betacoronavirus and Paramyxovirus in Hipposideros bat species in Zimbabwe. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Meegid(2017), <https://doi.org/10.1016/j.meegid.2018.01.007>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Short communication

Circulation of *Alphacoronavirus*, *Betacoronavirus* and *Paramyxovirus* in *Hipposideros* bat species in Zimbabwe.

Mathieu Bourgarel^{1,2}, Davies M. Pfukenyi³, Vanina Boué⁴, Loïc Talignani⁴, Ngoni Chiweshe¹, Fodé Diop⁴, Alexandre Caron^{1,5,6}, Gift Matope³, Dorothée Missé⁴ and Florian Liégeois^{4,*}.

¹ CIRAD, UMR ASTRE, RP-PCP, Harare, Zimbabwe.

² ASTRE, Univ. Montpellier, CIRAD, INRA, Montpellier, France.

³ University of Zimbabwe, Faculty of veterinary sciences, Harare, Zimbabwe

⁴ MIVEGEC, IRD, CNRS, Univ. Montpellier, Montpellier, France.

⁵ CIRAD, UMR ASTRE, RP-PCP, Maputo, Mozambique

⁶ Faculdade de Veterinária, Universidade Eduardo Mondlane, Maputo, Mozambique

Bourgarel Mathieu: CIRAD-ASTRE, 6 Lanark Road – Harare, Zimbabwe;

mathieu.bourgarel@cirad.com

Chiweshe Ngoni: CIRAD-ASTRE, 6 Lanark Road – Harare, Zimbabwe; chiweshengoni@gmail.com

Caron Alexandre: CIRAD-ASTRE, Faculdade de Veterinária - Universidade Eduardo Mondlane Av. de Moçambique Km. 1,5 - Caixa Postal 257, Maputo 01009, Moçambique; alexandre.caron@cirad.fr

Pfukenyi Davies M.: Faculty of Veterinary Science, University of Zimbabwe, P.O. Box MP167, Mt Pleasant Harare, Zimbabwe ; dpfukas@gmail.com

Matope Gift: Faculty of Veterinary Science, University of Zimbabwe, P.O. Box MP167, Mt Pleasant Harare, Zimbabwe ; giftmatope@gmail.com

Boué Vanina: MIVEGEC, IRD, CNRS, Univ. Montpellier, France: sveltina@yahoo.fr

Talignani Loïc: MIVEGEC, IRD, CNRS, Univ. Montpellier, France: loic.talignani@ird.fr

Diop Fode: MIVEGEC, IRD, CNRS, Univ. Montpellier, France: fode.diop@ird.fr

Dorothee Missé: MIVEGEC, IRD, CNRS, Univ. Montpellier, France: dorothee.misse@ird.fr

*Corresponding Author:

Florian Liégeois

IRD / UMR 224, MIVEGEC

911 avenue Agropolis

34394 Montpellier

E-Mail: florian.liegeois@ird.fr

Nucleotide sequence accession number:

The new *Coronavirus* and *Paramyxovirus* sequences reported in this study are available in GenBank under the following accession numbers: BtCov-Zim001Mab, MG000865; BtCov-Zim015Mab, MG000866; BtCov-Zim019Mab, MG000867; BtCov-Zim021, Mab MG000868; BtCov-Zim037Mab, MG000869; BtCov-Zim040Mab, MG000870; BtCov-Zim034Mab, MG000871; BtCov-Zim035Mag, MG000872; BtPV-Zim026Mag, MG000873.

Abstract

Bats carry a great diversity of zoonotic viruses with a high-impact on human health and livestock. Since the emergence of new coronaviruses and paramyxoviruses in humans (e.g. Severe Acute Respiratory Syndrome Coronavirus (SARS-CoV) and Nipah virus), numerous studies clearly established that bats can maintain some of these viruses. Improving our understanding on the role of bats in the epidemiology of the pathogens they harbour is necessary to prevent cross-species spill over along the wild/domestic/human gradient. In this study, we screened bat faecal samples for the presence of *Coronavirus* and *Paramyxovirus* in two caves frequently visited by local people to collect manure and/or to hunt bats in Zimbabwe. We amplified partial *RNA-dependent RNA polymerase* genes of *Alpha* and *Betacoronavirus* together with the partial *polymerase* gene of *Paramyxovirus*. Identified coronaviruses were related to pathogenic human strains and the paramyxovirus belonged to the recently described *Jeilongvirus* genus. Our results highlighted the importance of monitoring virus circulation in wildlife, especially bats, in the context of intense human-wildlife interfaces in order to strengthen prevention measures among local populations and to implement sentinel surveillance in sites with high zoonotic diseases transmission potential.

Keywords: Bat, *Coronavirus*, *Paramyxovirus*, Phylogeny, Emerging Infectious Diseases, Zimbabwe

Bats comprise nearly 1200 species and constitute $\approx 20\%$ of living mammal species and are distributed on all continents except Antarctic, Arctic and a few islands (Simmons, 2005). Due to their unique (only flying mammals) and diverse lifestyles, bats differ from other sylvatic disease mammalian reservoirs and are predisposed for the acquisition and maintenance of viruses (Hayman et al., 2013). During the past two decades, bats (*Chiroptera*) have been identified as the reservoir host of a number of high-impact zoonotic viruses known to induce highly lethal diseases in humans and domestic animals (Brook and Dobson, 2015). They have been associated with emerging *Paramyxovirus* (Nipah and hendra viruses), *Coronavirus* (MERS-CoV and SARS-CoV) and *Filovirus* (Ebola and Marburg viruses) (Smith and Wang, 2013) which attracted global attention due to their severity and/or large-scale spread. Those emergences have been caused by the ever-increasing interfaces between domestic animals, people and bat communities created by current global and human changes (Brierley et al., 2016). Human activities that increase exposure to bats induce new and more infectious contacts between species and promote the spill over of unknown pathogens from bats to other animals. The identification of the reservoir species is key for the control of these emerging infectious diseases in order to prevent/manage practices at risk of pathogens spill over.

Although numerous studies have been implemented on bat-borne viruses around the world, large gaps still exist concerning the viral diversity among *Chiroptera* especially in some regions that attracted little disease research until now. The Republic of Zimbabwe is situated southern Africa in the subtropical zone and has an exceptional great diversity of wildlife. To date more than 60 bat species have been recorded in Zimbabwe (Monadjem, 2010). Accordingly, Zimbabwe represents a potential hot spot for future emergence of microorganisms from bats that can transmit

infections to humans and livestock (Morse et al., 2012). Many cases of rabies, anthrax, African swine fever and foot and mouth diseases have been recorded in the last 20 years. Furthermore, *Lyssavirus* were demonstrated in bats (Duvenhage virus) and in cats and dogs (Mokola Virus, Lagos bat virus) (Bingham et al., 2001; Foggin, 1982). In the 1970s, a traveller who passed through Zimbabwe was probably infected with the *Marburg* virus after visiting the Chinhoyi caves about 135 km northwest of Harare, capital of Zimbabwe (Peterson et al., 2006). We report here the first evidence of circulation of *Coronavirus* and *Paramyxovirus* in *Hipposideros* bat species in Zimbabwe.

Between June 2016 and February 2017, 99 and 146 faecal samples were respectively collected in two caves (Fig. 1) regularly visited by local people to collect bat guano used as fertilizer. Each cave was visited twice at two different periods. Two square meters plastic sheets were laid down in the caves, underneath the bat colonies for overnight (five plastic sheets per cave). Faeces were collected from each plastic sheet at a rate of \approx 6 grams of pooled faeces in 15 ml tube with 6 ml of homemade RNA stabilization solution (Pol Scientific, 1999). Samples were stored at -80°C until laboratory analyses.

Bat species were identified by *Cytochrome b* amplification (Kocher et al., 1989) and sequencing after DNA extraction using Qiamp DNA stool (Qiagen S.A, Courtaboeuf, France). *Cytochrome b* sequences were then compared to available bat sequences in the GenBank database using *Basic Local Alignment Search Tool* (BLAST) program and species were confirmed by phylogenetic analysis (supplementary material, Fig 1S,). Only bats from *Hipposideros spp.*, representing two distinct colonies, were identified. To date, two different *Hipposideros* bat species have been reported in Zimbabwe; *Hipposideros caffer* and *Hipposideros vittatus* (Monadjem,

2010). Our samples were closer to *Hip. caffer* than any other *Hipposideros spp* (supplementary material, Fig 1S,).

RNA extraction was carried out from all faecal samples collected. Briefly, two sample tubes from the same plastic sheet were pooled and transferred in a 50 ml tube with 20 ml of PBS 1X then vigorously mixed. All together we made 73 (51 in June 2016 and 22 in February 2017) pools from Mabura cave and 50 (35 in June 2016 and 15 in February 2017) pools from Magweto cave respectively. Tubes were centrifuged at 4500 rpm for 10 min. Supernatant was filtered using gauze in order to eliminate faecal matter and transferred in fresh tubes then re-centrifuged at 4500 rpm for 10 min. Supernatant was filtered through a 0.2 μm filter to remove eukaryotic and bacterial sized particles. Seven millilitres of filtered samples were centrifuged at 250,000 g for 2.5 h at 4°C. The pellets were re-suspended in 600 μl H₂O molecular grade and 150 μl were used to extract RNA using NucleoSpin® RNA Kit (Macherey-Nagel, France) according to the manufacturer's protocol. The 123 RNA samples extracted from the pools were then reverse transcribed using random hexamers and screened for *Coronavirus* (CoV) and *Paramyxovirus* (ParV) as previously described employing a pan-coronavirus and pan-paramyxovirus nested RT-PCR directed against partial polymerase *RNA-dependent RNA polymerase (RdPd)* and *polymerase* gene sequences, respectively (Chu et al., 2011; Tong et al., 2008). PCR products (415 bp for CoV and 531 bp for ParV) were agarose gel purified (GeneClean Turbo Kit, MP Biomedicals, France) and directly sequenced in both 5' and 3' directions using cycle sequencing and dye terminator methodologies (Eurofins, Germany). Overlapping sequences were assembled into contiguous sequences using SEQMAN DNASTAR software (lasergene, DNASTAR, Inc., Madison, WI, USA). Partial non-concatenated nucleic acid sequences of the new *Coronavirus* and

Paramyxovirus as well as from *Cytochrome B* were aligned using MEGA 7 (Kumar et al., 2016), with minor manual adjustments. Sites that could not be unambiguously aligned were excluded and divergent regions were excluded from subsequent analyses. Phylogenies were inferred using both Bayesian methods and Maximum Likelihood (ML) method implemented in MrBayes v3.2.6 and in PhyML respectively (Guindon et al., 2010; Ronquist et al., 2012). Mr Bayes ran for four million generations for *Coronavirus RdRp* and *Paramyxovirus polymerase* genes, respectively, with a 10% burn-in. Bayesian parameters were examined with the Tracer program (Tracer). Convergence diagnostic for the Estimated sample Size (ESS) values and Potential Scale Reduction Factor (PSRF) were >500 and equal to 1 respectively. In ML method, the reliability of branching orders was tested using the bootstrap approach (1000 replicates). The suited evolution model (GTR + Γ_4 + I for *Coronavirus* and *Cytochrome B*, and GTR + Γ_4 for *Paramyxovirus*) was selected by Akaike's Information criterion (AIC) using Topali software (Milne et al., 2009). From both phylogenetic analyses, similar tree topologies were obtained (data not shown). Identities analyses were done using ClustalX (Larkin et al., 2007).

We characterized *Alphacoronavirus* in Mabura cave as well as *Betacoronavirus* and *Paramyxovirus* in Magweto cave from roundleaf bats, which was the only bat genus observed in the two visited caves at the time of our samplings. Our new *Alphacoronavirus* formed a well sustained specific sub-clade close to the human *Coronavirus* 229E strain (HCoV-229E) (Fig 2) that circulates in human population worldwide and mostly causes mild respiratory disease (Perlman, 2013). This close relationship is confirmed by a high percentage (95%) of amino acid identities (Supplementary Material, Table S1). Interestingly, our BtCoV 229E related strains are distinct to those identified in *Hip. caffer rufer* from Ghana (Pfefferle et al.,

2009). Our results are in accord with the recently suggested long evolutionary history of 229E-related CoV in old world hipposiderid bats (Corman et al., 2015). Nonetheless it is unclear whether bats directly transmitted this virus to human or if an intermediate host was involved in the transmission chain such as demonstrated for SARS-CoV and MERS-CoV (Smith and Wang, 2013).

In Mabura cave, during our first visit during the cold dry season in June 2016 we collected faeces from three plastic sheets and Bat 229-E like virus was amplified from samples issued from each plastic sheet suggesting an important circulation of this virus in the bat colony. Interestingly, no viruses were amplified from the second sampling in this cave during the rainy season in February 2017. Nonetheless, during the second visit we observed a consequent diminution of bats present in the cave and our sampling was lower than expected. This could be due to *Hipposideros* spp. seasonal movement. Besides, the absence of *Alphacoronavirus* could also be due to temporal variation in virus shedding in bats (Plowright et al., 2015).

In Magweto cave we amplified *Betacoronavirus* from only one pooled sample (Fig 2). It could be due to a low circulation of this virus in the bat colony. Phylogenetic analyses showed that this new virus formed a specific clade with betacoronaviruses isolated in Asia and Africa (Gouilh et al., 2011; Pfefferle et al., 2009; Quan et al., 2010) with 90% to 87% of amino acid identities (Supplementary material, Table S1) and together they formed a sister clade with the described SARS-CoV strains with 77% of amino acid identities (Fig 2, Supplementary material, Table S1). The SARS-CoV related (SARS-CoVr) sister clade is well sustained and our new Bt SARS-CoVr strain is positioned at the root of this clade. This finding could strengthen the African origin hypothesis of SARS-like group (Pfefferle et al., 2009; Quan et al., 2010). Nonetheless, this hypothesis is controversial and, in order to disentangle the Bt

SARS-CoVr origin, future studies should focus on Hipposideridae as well as on Rhinolophidae and Rhinonycteridae since these three bat families diverged from a common ancestor, which potentially hosted the ancestor of SARS-related COVs (Foley et al., 2015; Gouilh et al., 2011). Additionally, SARS-CoVr have been characterised from these three bat families (Pfefferle et al., 2009; Smith et al., 2016; Wu et al., 2016).

SARS-CoV emerged at the beginning of 21st century following a human transmission by an intermediary host, a palm civet, in China. More than 8000 human infections were reported around the world with a case fatality rate of up to 10% (Smith and Wang, 2013). To date several studies evidenced different bat species as potential SARS and SARS-like CoV reservoirs worldwide (Li et al., 2005).

In addition, in the same cave we amplified a *Paramyxovirus* closer to bat *Paramyxovirus* (77 to 87% of amino acid identities) related to the putative *Jeilongvirus* genus (Fig 3, Supplementary material Table S1) than other *Paramyxovirus* lineages. To date, the pathogenic potential of the viruses from this genus is currently unknown. However, the *Beilong* virus was discovered on human kidney cell lines and neutralizing antibodies against *J* virus have been detected in rodents, pigs and humans (Audsley et al., 2016). In addition, bat viruses belonging to the related-*Jeilongvirus* genus were widely detected in China and more recently in Luxembourg in Europe (Pauly et al., 2017). Altogether, these data highlight the need for further studies on the zoonotic potential of these viruses.

Although *Coronavirus* and *Paramyxovirus* have been widely described in bats around the world (Anthony et al., 2017; Drexler et al., 2012), our results pointed out the need to widen viral screening in under-investigated countries particularly when the country has considerable potential as a hot spot for emerging infectious diseases

(Morse et al., 2012). Our study focused on two caves in Zimbabwe with an important bat-human interface throughout guano harvesting and/or bats poaching. Non-invasive sampling provides a rapid approach to target site of interest for in-depth studies on virus prevalence in bats and temporal variation in virus shedding in bats (viral ecology) and provides a first risk assessment of the transmission of bat-borne pathogens to humans. Finally, our study will enable, in agreement with the local health authorities, to carry out a specific communication within the local populations on the risk of contamination and how to prevent it.

Acknowledgements

We thank Billy Butete and Cavin Mandina for their field assistance. We thank Estelle Ména for her technical assistance. This work was supported by grants from the Agence Nationale de la Recherche (grant ANR-14-CE14-0029). We thank the Research Council of Zimbabwe for approving this study (research registration certificate N° 03006) and the Hurungwe Rural District council and the Zibagwe Rural District Council for their assistance and facilitation. We thank the Animal Research Ethics Committee of Zimbabwe for their approval (ref number 002/2017). This work was conducted within the framework of the Research Platform “Production and Conservation in Partnership” (RP-PCP).

Declaration of interest

We declare that we have no conflicts of interest.

References

Anthony, S.J., Johnson, C.K., Greig, D.J., Kramer, S., Che, X., Wells, H., Hicks, A.L., Joly, D.O., Wolfe, N.D., Daszak, P., Karesh, W., Lipkin, W.I., Morse, S.S., Consortium, P., Mazet, J.A.K., Goldstein, T., 2017. Global patterns in coronavirus diversity. *Virus Evol* 3, vex012.

Audsley, M.D., Marsh, G.A., Lieu, K.G., Tachedjian, M., Joubert, D.A., Wang, L.F., Jans, D.A., Moseley, G.W., 2016. The immune evasion function of J and Beilong virus V proteins is distinct from that of other paramyxoviruses, consistent with their inclusion in the proposed genus Jeilongvirus. *J Gen Virol* 97, 581-592.

Bingham, J., Javangwe, S., Sabeta, C.T., Wandeler, A.I., Nel, L.H., 2001. Report of isolations of unusual lyssaviruses (rabies and Mokola virus) identified retrospectively from Zimbabwe. *J S Afr Vet Assoc* 72, 92-94.

Brierley, L., Vonhof, M.J., Olival, K.J., Daszak, P., Jones, K.E., 2016. Quantifying Global Drivers of Zoonotic Bat Viruses: A Process-Based Perspective. *Am Nat* 187, E53-64.

Brook, C.E., Dobson, A.P., 2015. Bats as 'special' reservoirs for emerging zoonotic pathogens. *Trends Microbiol* 23, 172-180.

Chu, D.K., Leung, C.Y., Gilbert, M., Joyner, P.H., Ng, E.M., Tse, T.M., Guan, Y., Peiris, J.S., Poon, L.L., 2011. Avian coronavirus in wild aquatic birds. *J Virol* 85, 12815-12820.

Corman, V.M., Baldwin, H.J., Tateno, A.F., Zerbinati, R.M., Annan, A., Owusu, M., Nkrumah, E.E., Maganga, G.D., Oppong, S., Adu-Sarkodie, Y., Vallo, P., da Silva Filho, L.V., Leroy, E.M.,

Thiel, V., van der Hoek, L., Poon, L.L., Tschapka, M., Drosten, C., Drexler, J.F., 2015. Evidence for an Ancestral Association of Human Coronavirus 229E with Bats. *J Virol* 89, 11858-11870.

Drexler, J.F., Corman, V.M., Muller, M.A., Maganga, G.D., Vallo, P., Binger, T., Gloza-Rausch, F., Cottontail, V.M., Rasche, A., Yordanov, S., Seebens, A., Knornschild, M., Oppong, S., Adu Sarkodie, Y., Pongombo, C., Lukashev, A.N., Schmidt-Chanasit, J., Stocker, A., Carneiro, A.J., Erbar, S., Maisner, A., Fronhoffs, F., Buettner, R., Kalko, E.K., Kruppa, T., Franke, C.R., Kallies, R., Yandoko, E.R., Herrler, G., Reusken, C., Hassanin, A., Kruger, D.H., Matthee, S., Ulrich, R.G., Leroy, E.M., Drosten, C., 2012. Bats host major mammalian paramyxoviruses. *Nat Commun* 3, 796.

Foggin, C.M., 1982. Atypical rabies virus in cats and a dog in Zimbabwe. *Vet Rec* 110, 338.

Foley, N.M., Thong, V.D., Soisook, P., Goodman, S.M., Armstrong, K.N., Jacobs, D.S., Puechmaille, S.J., Teeling, E.C., 2015. How and why overcome the impediments to resolution: lessons from rhinolophid and hipposiderid bats. *Mol Biol Evol* 32, 313-333.

Gouilh, M.A., Puechmaille, S.J., Gonzalez, J.P., Teeling, E., Kittayapong, P., Manuguerra, J.C., 2011. SARS-Coronavirus ancestor's foot-prints in South-East Asian bat colonies and the refuge theory. *Infect Genet Evol* 11, 1690-1702.

Guindon, S., Dufayard, J.F., Lefort, V., Anisimova, M., Hordijk, W., Gascuel, O., 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst Biol* 59, 307-321.

Hayman, D.T., Bowen, R.A., Cryan, P.M., McCracken, G.F., O'Shea, T.J., Peel, A.J., Gilbert, A., Webb, C.T., Wood, J.L., 2013. Ecology of zoonotic infectious diseases in bats: current knowledge and future directions. *Zoonoses Public Health* 60, 2-21.

Kocher, T.D., Thomas, W.K., Meyer, A., Edwards, S.V., Paabo, S., Villablanca, F.X., Wilson, A.C., 1989. Dynamics of mitochondrial DNA evolution in animals: amplification and sequencing with conserved primers. *Proc Natl Acad Sci U S A* 86, 6196-6200.

Kumar, S., Stecher, G., Tamura, K., 2016. MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol* 33, 1870-1874.

Larkin, M.A., Blackshields, G., Brown, N.P., Chenna, R., McGettigan, P.A., McWilliam, H., Valentin, F., Wallace, I.M., Wilm, A., Lopez, R., Thompson, J.D., Gibson, T.J., Higgins, D.G., 2007. Clustal W and Clustal X version 2.0. *Bioinformatics* 23, 2947-2948.

Li, W., Shi, Z., Yu, M., Ren, W., Smith, C., Epstein, J.H., Wang, H., Crameri, G., Hu, Z., Zhang, H., Zhang, J., McEachern, J., Field, H., Daszak, P., Eaton, B.T., Zhang, S., Wang, L.F., 2005. Bats are natural reservoirs of SARS-like coronaviruses. *Science* 310, 676-679.

Milne, I., Lindner, D., Bayer, M., Husmeier, D., McGuire, G., Marshall, D.F., Wright, F., 2009. TOPALi v2: a rich graphical interface for evolutionary analyses of multiple alignments on HPC clusters and multi-core desktops. *Bioinformatics* 25, 126-127.

Monadjem, A., P. J. Taylor, AND M. C. Schoeman, 2010. *Bats of Southern and Central Africa: A Biogeographic and Taxonomic Synthesis*. Book, Wits University Press.

Morse, S.S., Mazet, J.A., Woolhouse, M., Parrish, C.R., Carroll, D., Karesh, W.B., Zambrana-Torrel, C., Lipkin, W.I., Daszak, P., 2012. Prediction and prevention of the next pandemic zoonosis. *Lancet* 380, 1956-1965.

Pauly, M., Pir, J.B., Loesch, C., Sausy, A., Snoeck, C.J., Hubschen, J.M., Muller, C.P., 2017. Novel Alphacoronaviruses and Paramyxoviruses Cocirculate with Type 1 and Severe Acute Respiratory System (SARS)-Related Betacoronaviruses in Synanthropic Bats of Luxembourg. *Appl Environ Microbiol* 83.

Perlman, P.S.M.a.S., 2013. Coronaviridae. In *Fields Virology* (Sixth Ed.), Lippincott Williams & Wilkins, Philadelphia., 825-858.

Peterson, A.T., Lash, R.R., Carroll, D.S., Johnson, K.M., 2006. Geographic potential for outbreaks of Marburg hemorrhagic fever. *Am J Trop Med Hyg* 75, 9-15.

Pfefferle, S., Oppong, S., Drexler, J.F., Gloza-Rausch, F., Ipsen, A., Seebens, A., Muller, M.A., Annan, A., Vallo, P., Adu-Sarkodie, Y., Kruppa, T.F., Drosten, C., 2009. Distant relatives of severe acute respiratory syndrome coronavirus and close relatives of human coronavirus 229E in bats, Ghana. *Emerg Infect Dis* 15, 1377-1384.

Plowright, R.K., Eby, P., Hudson, P.J., Smith, I.L., Westcott, D., Bryden, W.L., Middleton, D., Reid, P.A., McFarlane, R.A., Martin, G., Tabor, G.M., Skerratt, L.F., Anderson, D.L., Cramer, G., Quammen, D., Jordan, D., Freeman, P., Wang, L.F., Epstein, J.H., Marsh, G.A., Kung'u, N.Y., McCallum, H., 2015. Ecological dynamics of emerging bat virus spillover. *Proc Biol Sci* 282, 20142124.

Quan, P.L., Firth, C., Street, C., Henriquez, J.A., Petrosov, A., Tashmukhamedova, A., Hutchison, S.K., Egholm, M., Osinubi, M.O., Niezgodna, M., Ogunkoya, A.B., Briese, T., Rupprecht, C.E., Lipkin, W.I., 2010. Identification of a severe acute respiratory syndrome coronavirus-like virus in a leaf-nosed bat in Nigeria. *MBio* 1.

Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D.L., Darling, A., Höhna, S., Larget, B., Liu, L., Suchard, M.A., Huelsenbeck, J.P., 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Syst Biol* 61, 539-542.

Pol Scientific, 1999. <http://www.protocol-online.org/prot/Protocols/RNAlater-3999.html>.

Simmons, N.B., 2005. Evolution. An Eocene big bang for bats. *Science* 307, 527-528.

Smith, C.S., de Jong, C.E., Meers, J., Henning, J., Wang, L., Field, H.E., 2016. Coronavirus Infection and Diversity in Bats in the Australasian Region. *Ecohealth* 13, 72-82.

Smith, I., Wang, L.F., 2013. Bats and their virome: an important source of emerging viruses capable of infecting humans. *Curr Opin Virol* 3, 84-91.

Tong, S., Chern, S.W., Li, Y., Pallansch, M.A., Anderson, L.J., 2008. Sensitive and broadly reactive reverse transcription-PCR assays to detect novel paramyxoviruses. *J Clin Microbiol* 46, 2652-2658.

Tracer, 2003. v1.6. Available online: <http://tree.bio.ed.ac.uk/software/tracer/>.

Wu, Z., Yang, L., Ren, X., Zhang, J., Yang, F., Zhang, S., Jin, Q., 2016. ORF8-Related Genetic Evidence for Chinese Horseshoe Bats as the Source of Human Severe Acute Respiratory Syndrome Coronavirus. *J Infect Dis* 213, 579-583.

ACCEPTED MANUSCRIPT

Figure 1. Geographical distribution of bats faecal samples collection sites.

Blues circles represent the caves where bat faecal samples were collected. The name of the caves as well as the GPS location is noted next to the circle. Number of faeces collected per site is shown in brackets.

Figure 2. Phylogenetic analysis of partial *RNA-dependent RNA polymerase (RdRp)* of the newly identified *Alphacoronavirus* and *Betacoronavirus* sequences from Zimbabwe. New partial *RdRp* (415bp) CoV sequences are represented in bold and were compared to previously identified *Alphacoronavirus* and *Betacoronavirus* available in the GenBank. Accession numbers are showed before the strain name. Only Bayesian posterior probabilities are showed. Asterisks at nodes represent posterior probability $\geq 90\%$. Scale bars indicate the number of base substitutions per site.

Figure 3. Phylogenetic analysis of partial *polymerase* gene of the newly identified *Paramyxovirus* (ParV) sequence from Zimbabwe. New partial *pol* (531bp) ParV sequences are represented in bold and were compared to previously identified *Paramyxovirus* available in the GenBank. Accession numbers are showed before the strain name. Only Bayesian posterior probabilities are showed. Asterisks at nodes represent posterior probability $\geq 90\%$. Scale bars indicate the number of base substitutions per site

Figure 1S, supplementary material:

Phylogenetic analysis of partial *CytB* sequences (550 bp). New *CytB* sequences obtained were compared to representative bat family *CytB* sequences available in

the GenBank. Only Bayesian posterior probabilities are showed. Asterisks at nodes represent posterior probability $\geq 85\%$. Scale bars indicate the number of base substitutions per site. New *CytB* sequences are represented by the red triangle. The following sequences have been used for this analysis: KX467590.1 *Miniopterus schreibersii*, KX467591.1 *Miniopterus magnater*, KX467593.1 *Tylonycteris pachypus*, KX467595.1 *Hypsugoal aschanicus*, KX467596.1 *Nyctalus plancyi*, KX467597.1 *Pipistrellus abramus*, KX467598.1 *Murina leucogaster*, KX467601.1 *Myotis laniger*, KX467611.1 *Myotis davidii*, KX467606.1 *Myotis blythii*, KX467607.1 *Myotis chinensis*, KX467608.1 *Myotis pequinius*, KX467609.1 *Myotis bombinus*, KX467605.1 *Myotis formosus*, KX467610.1 *Myotis altarium*, KX467612.1 *Myotis frater*, KX467599.1 *Myotis ricketti*, KX467600.1 *Myotis adversus*, KX467604.1 *Myotis fimbriatus*, KX467602.1 *Myotis daubentonii*, KX467603.1 *Myotis macrodactylus*, KF218429.1 *Taphozous nudiventris*, AY591536.1 *Otomops martiensseni*, AY591537.1 *Otomops martiensseni*, KX467574.1 *Megaderma lyra*, KX467575.1 *Rhinolophus pusillus*, KX467576.1 *Rhinolophus lepidus*, KX467579.1 *Rhinolophus macrotis*, KX467580.1 *Rhinolophus marshalli*, KX467582.1 *Rhinolophus rex*, KX467578.1 *Rhinolophus ferrumequinum*, KX467583.1 *Rhinolophus sinicus*, KX467577.1 *Rhinolophus affinis*, KX467581.1 *Rhinolophus pearsonii*, KX467584.1 *Hipposideros cineraceus*, KX467585.1 *Hipposideros pomona*, FJ347979.1 *Hipposideros caffer*, FJ347980.1 *Hipposideros caffer*, KX467587.1 *Hipposideros armiger*, KX467588.1 *Hipposideros larvatus*, KX467589.1 *Hipposideros pratti*, KX467586.1 *Aselliscus stoliczkanus*, KT583805.1 *Hipposideros vittatus*, KT583806.1 *Hipposideros vittatus*, KT583801.1 *Hipposideros gigas*, KT583817.1 *Hipposideros commersoni*, DQ005796.1 *Triaenops persicus*, DQ005797.1 *Triaenops persicus*, KX823312.1 *Megaerops niphanae*, KX823313.1 *Cynopterus sp.*, KX823314.1 *Cynopterus sp.*, KX823317.1 *Sphaeria sblanfordi*, KX822888.1 *Eidolon helvum*, KX822889.1 *Eidolon helvum*, KX823318.1 *Macroglossus sobrinus*, KX823319.1 *Eonycteris spelaea*, KX822938.1 *Rousettus aegyptiacus*, KX822939.1 *Rousettus aegyptiacus*, KX823315.1 *Rousettus leschenaultii*, KX823311.1 *Plerotes anchietae*, KX823011.1 *Hypsignathus monstrosus*, KX823012.1 *Hypsignathus monstrosus*, KX823087.1 *Epomops franqueti*, KX823088.1 *Epomops franqueti*, KX823233.1 *Epomops buettikoferi*, KX823234.1 *Epomops*

buettikoferi, KX822985.1 *Nanonycteris veldkampii*, KX822986.1 *Nanonycteris veldkampii*,
KX823306.1 *Epomops dobsonii*, KX823307.1 *Epomophorus anelli*, KX823308.1 *Epomophorus*
crypturus, KX823309.1 *Epomophorus labiatus*, KX823310.1 *Epomophorus minimus*, KX822798.1
Epomophorus gambianus, KX822810.1 *Micropteropus pusillus*, KX822797.1 *Epomophorus*
gambianus, KX822809.1 *Micropteropus pusillus*.

ACCEPTED MANUSCRIPT

Circulation of *Alphacoronavirus*, *Betacoronavirus* and *Paramyxovirus* in *Hipposideros* bat species in Zimbabwe.

Highlights

- *Coronavirus* and *Paramyxovirus* circulate in *Hipposideros* bat species in Zimbabwe.
- Importance of widening viral screening in under-investigated countries.
- Sentinel surveillance in sites with high zoonotic transmission potential.



Figure 1




Figure 2




Figure 3