

Journal Pre-proof

Sero-prevalence, cross-species infection and serological determinants of prevalence of Bovine Coronavirus in Cattle, Sheep and Goats in Ghana

Vitus Burimuah, Augustina Sylverken, Michael Owusu, Philip El-Duah, Richmond Yeboah, Jones Lamptey, Yaw Oppong Frimpong, Olivia Agbenyega, Raphael Folitse, William Tasiame, Benjamin Emikpe, Eddie-Williams Owiredu, Samuel Oppong, Yaw Adu-Sarkodie, Christian Drosten



PII: S0378-1135(19)31150-2

DOI: <https://doi.org/10.1016/j.vetmic.2019.108544>

Reference: VETMIC 108544

To appear in: *Veterinary Microbiology*

Received Date: 27 September 2019

Revised Date: 2 December 2019

Accepted Date: 2 December 2019

Please cite this article as: Burimuah V, Sylverken A, Owusu M, El-Duah P, Yeboah R, Lamptey J, Frimpong YO, Agbenyega O, Folitse R, Tasiame W, Emikpe B, Owiredu E-Williams, Oppong S, Adu-Sarkodie Y, Drosten C, Sero-prevalence, cross-species infection and serological determinants of prevalence of Bovine Coronavirus in Cattle, Sheep and Goats in Ghana, *Veterinary Microbiology* (2019), doi: <https://doi.org/10.1016/j.vetmic.2019.108544>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2019 Published by Elsevier.

Sero-prevalence, cross-species infection and serological determinants of prevalence of Bovine Coronavirus in Cattle, Sheep and Goats in Ghana

Vitus Burimuah^{1,2}, Augustina Sylverken^{3,4}, Michael Owusu^{4,5}, Philip El-Duah^{1,4,6}, Richmond Yeboah^{1,4}, Jones Lamptey^{1,4}, Yaw Oppong Frimpong^{4,7}, Olivia Agbenyega⁸, Raphael Folitse², William Tasiame^{2,6}, Benjamin Emikpe², Eddie-Williams Owiredu⁹, Samuel Oppong¹⁰, Yaw Adu-Sarkodie¹, Christian Drosten^{6*} Christian.drosten@charite.de

1. Department of Clinical Microbiology, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana; vitus7uk@yahoo.co.uk (VB); elduahphilip9@gmail.com (PED); yasax@hotmail.co.uk (YAS)
2. School of Veterinary Medicine, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana; vitus7uk@yahoo.co.uk (VB); raphfolitse@yahoo.com (RF)
3. Department of Theoretical and Applied Biology, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana; annan@kccr.de
4. Kumasi Centre for Collaborative Research in Tropical Medicine, Kumasi, Ghana; annan@kccr.de (AS); owusumichael-gh@hotmail.com (MO); elduahphilip9@gmail.com (PED); Yeboahrichmond82@yahoo.com (RY); jlampzey80@gmail.com (JL); oppongfrimpong1@gmail.com (YOF)
5. Department of Medical Laboratory Technology, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana; owusumichael-gh@hotmail.com
6. Institute of Virology, Charite, Universitätsmedizin Berlin, Germany; elduahphilip9@gmail.com (PED); christian.drosten@charite.de (CD)
7. Department of Animal Science, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana; oppongfrimpong1@gmail.com
8. Department of Agroforestry, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana; olivia_agbenyega@yahoo.com
9. Department of Molecular Medicine, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana; eddiewilliams.owiredu@gmail.com
10. Department of Wildlife and Range Management, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana; kobbyoppong@yahoo.com

Address correspondence to

Christian Drosten, Charité – Universitätsmedizin Berlin, Institute of Virology, Charitéplatz 1, 10117 Berlin, Germany. E-mail:

Highlights

- Bovine coronavirus has considerable seroprevalence in cattle across Ghana
- Sheep and goats are kept without strict separation from cattle and show seropositivity against bovine coronavirus
- Bovine coronavirus seroprevalence is positively correlated with large farm size
- Highest bovine coronavirus seroprevalence was found in Ghana's Northern Province with prevailing arid climate

Abstract

Cattle, goats and sheep are dominant livestock species in sub-Saharan Africa, with sometimes limited information on the prevalence of major infectious diseases. Restrictions due to notifiable epizootics complicate the exchange of samples in surveillance studies and suggest that laboratory capacities should be established domestically. Bovine Coronavirus (BCoV) causes mainly enteric disease in cattle. Spillover to small ruminants is possible. Here we established BCoV serology based on a recombinant immunofluorescence assay for cattle, goats and sheep, and studied the seroprevalence of BCoV in these species in four different locations in the Greater Accra, Volta, Upper East, and Northern provinces of Ghana. The whole sampling and testing was organized and conducted by a veterinary school in Kumasi, Ashanti Region of Ghana. Among sampled sheep (n=102), goats (n=66), and cattle (n=1495), the seroprevalence rates were 25.8%, 43.1% and 55.8%. For cattle, seroprevalence was significantly higher on larger farms (82.2% vs 17.8%, comparing farms with >50 or <50 animals; p=0.027). Highest prevalence was seen in the Northern province with dry climate, but no significant trend following the north-south gradient of sampling sites was detected. Our study identifies a considerable seroprevalence for BCoV in Ghana and provides further support for the spillover of BCoV to small ruminants in settings with mixed husbandry and limited separation between species.

Keywords: Bovine Coronavirus, Africa, Ghana, Global Health

Introduction

Cattle, goats and sheep are among the major livestock species in Ghana. The present numbers in 2017 based on the Food and Agriculture Organization (FAO) animal production database are estimated to range around 1.76, 6.4, and 4.6 million cattle, goats and sheep in the country, respectively. Among livestock, only

chicken outnumber these species (74 million). While disease surveillance is in place, there are knowledge gaps concerning the laboratory-based prevalence of some major livestock diseases. Among these is Bovine Coronavirus (BCoV) that affects cattle and other livestock species including horses and camels.

BCoV is an enveloped plus strand RNA virus that belongs to the genus *Betacoronavirus* [1-3]. While different strains may have some antigenic variability, all strains elicit cross-reactive seropositivity and thus form a single serotype [4, 5]. The virus is an important livestock pathogen causing effects on animal welfare as well as the economy [6]. It causes diarrhea and respiratory disease in calves, as well as winter dysentery in adult cattle [7, 8]. Transmission of BCoV is mainly through respiratory or fecal-oral routes [4], infecting the respiratory (nasal, tracheal, and lung) and intestinal (villi and crypts of the ileum and colon) epithelial cells [9]. When infected with BCoV, within-herd transmission is generally rapid and infected animals display diverse clinical signs including diarrhea with or without blood, fever, and respiratory signs, which range from none to severe [4, 7].

In many African countries including Ghana, livestock species live in close contact and animals serve diverse purposes such as transportation, draught power, fuel, clothing and as a source of meat and milk. Husbandry practices do not involve the same standards of species separation and hygiene as in other parts of the world. Close and sustained interaction between different animals as well as between animals and humans pose a risk of interspecies spillover of pathogens.

BCoV is characteristically a cattle virus. However, reports indicate BCoV infections also occur in small ruminants. Previous studies in Australia [10], New Zealand [11], Chile [12], and Scotland [13] reported BCoV infection in small ruminants. Eisa and Mohamed, 2014 also detected BCoV antigens in goats [14] whereas Trávén *et al.*, detected BCoV antibodies in sheep [15]. Recently, Gumusova *et al.*, have also detected BCoV antibodies in goats [16].

Studies regarding the prevalence of BCoV and its associated risk factors are however limited in Africa, and none have been conducted in Ghana. This study evaluated the sero-prevalence of BCoV infection and assessed its associated risk factors among cattle, sheep, and goats in Ghana.

Materials and Methods

Study design and area

This study employed a cross-sectional design and was conducted between January, 2015 to December, 2018 in five districts in four regions of Ghana. Ghana is located in the west of Africa, sharing borders with Togo to the east, Cote d'Ivoire to the west, Burkina Faso to the north and the Gulf of Guinea, to the south and lies on latitude 7.9528 and longitude -1.0307. Ghana has a tropical climate with an average annual temperature of about 26°C and the annual rainfall of 736.6 mm/29". Agriculture dominates the economy of Ghana and extensive farming practices in the country increase the livestock-wildlife-human interface

Ethics approval

This study was approved by the Wildlife division of the Ghana Forestry Commission (Approval number: AO4957).

Study population, sampling strategy and data collection

A total of 1,498 animals aged ≥ 6 months, comprising 1,328 cattle, 104 sheep and 66 goats were included in the study. Animals aged < 6 months were excluded due to the possibility of detecting maternal antibodies.

Sampling was done using a simple two-stage cluster sampling technique. The Regional Veterinary Officers of the Ministry of Food and Agriculture (MOFA), Ghana in the selected regions were contacted for information on animal populations in their respective regions prior to the study. The list provided served as the sampling frame. Prior to the study, a survey was carried out and an inclusion criteria of animal population (cattle, sheep and goats) $\geq 1,000$ for districts to be eligible for selection for the study was upheld. As a result, five (5) districts that fulfilled the criteria were randomly selected. Secondly, farms within these districts with herd size ≥ 100 animals were randomly selected. All cattle, sheep, and goats within the selected farms were included in the study. If a district meets the first criteria, but the individual farms fail to meet the second criteria, farms which were very close to each other were pooled together.

Sites included were: Bongo district in the Upper East; Savelugu and Wale wale in Northern; Ada West in Greater Accra, and North Tongu in Volta. The map with the locations for sampling is shown in **Fig 1**.

Validated questionnaire was used to obtain data on possible risk factors of BCoV. Data collected include: age, sex, dietary changes, parturition, and lactation status of female animals. Additionally, the body score, presence of ectoparasites, and signs of infections such as fever, diarrhoea, respiratory distress, neurological disorder, and icterus were also assessed.



Fig 1. The map showing the locations for sampling.

Sample collection, processing, and analysis.

Ten milliliters (10 ml) of blood was collected through jugular puncture from each animal after disinfection of the site with 70% using vacutainer tubes (Becton Dickinson, NJ, USA) with needles (18 gauge). In the field, the blood was allowed to clot before transportation to the Veterinary laboratory in the district. At the district laboratory, the samples were spun for 10 min at 1,500 rpm to obtain the sera. The sera were transferred into three separate aliquots in cryotubes for each animal. The tubes were subsequently placed in liquid nitrogen to minimize antibody degradation. These processes were undertaken under sterile conditions. Upon obtaining representative samples per district, the frozen samples were transported in a cold-chain to the Kumasi Centre for Collaborative Research (KCCR) for long term storage at -70°C prior to laboratory analysis. Sample collection and preparation in each district and transportation to the KCCR lab took an average of 5-7 days.

During laboratory analysis, frozen sera were thawed at room temperature, vortexed and aliquots of 100 μl of each sample were prepared. The aliquots were incubated at 56°C for 30 minutes in warm water prior to recombinant immunofluorescence assay (IFA) as previously described [17-19]. Briefly, Vero B4 cells were co-transfected with pCG1 plasmids bearing Human coronavirus OC43 spike proteins. After overnight transfection, cells were harvested by treatment with trypsin to detach them in a cell culture incubator at 37°C and re-suspended in Dulbecco's Modified Eagle Medium (DMEM) (Invitrogen, USA) in 10% Foetal

Calf Serum (FCS). Aliquots of cells were pelleted at 300 x g for 5 minutes and washed twice with 1 ml phosphate buffered saline (PBS). Fifty microliters (50 μ l) of cell pellets were spotted on 12 well slides by dispensing and immediately aspirating, allowing 2 seconds interval between spotting. The cells were fixed using ice cold acetone/methanol (1:1), dried at ambient temperature, and kept at 4°C after drying for 20 minutes. To conduct the assay, 45 μ l of protein-free blocking solution (Roti®-Block, Carl Roth, Karlsruhe, Germany) was first added to each of the 12 spotted fields on the slide and incubated at room temperature in a humid box for 30 minutes, followed by rinsing with 1X tween-free PBS. After inactivation at 56 °C for 30 min, sera to be tested were diluted 1:100 in a 1X concentration of the protein-free blocking solution. Subsequently, 30 μ l of the diluted sera was applied on each of the spotted area and incubated at 37°C for 1 hour in a humid chamber, followed by rinsing with 0.1% tween in 1X PBS. Secondary antibody detection was done by the Alexa488 fluorescent reporter-conjugated goat anti-bovine, donkey anti-sheep, and donkey anti-goat IgG antibodies for cattle, sheep, and goat BCoV IgG respectively.

Test evaluation was done by microscopic examination under a fluorescent microscope and a positive outcome was determined by bright green cytoplasmic fluorescence as shown in Fig. 2.




Fig. 2. Depiction of typical BCoV IgG positive outcome (Panel A) against a negative outcome (Panel B). Cell nuclei were stained with DAPI and are shown as dark blue and the bright green impressions around the nuclei represent fluorescent antibody-antigen complexes.

Statistical Analysis

Data were presented as frequencies (percentages) and Chi square test was used to test for association where applicable. Univariate and multivariate logistic regression analysis were performed to determine the possible factors associated with BCoV sero-positivity for cattle, sheep, and goats. A p-value < 0.05 was considered statistically significant. All statistical analyses were performed using IBM Statistical Package for

the Social Sciences (SPSS) software 25 (SPSS Inc., Chicago, IL, USA), and GraphPad Prism 7 version 7.04 (GraphPad Software, Inc., La Jolla, California USA).

Results

The proportions of sheep, goat, and cattle were 4.4%, 6.8%, and 88.8% respectively. There were more adults than weaners (74.9% vs 25.1%) and more female animals (71.2%) compared to male animals (28.8%), with 22.5% of the females being lactating females at the time of the study. Majority of the animals had high rectal temperature (95.6%) and 4.4% had physical signs of disease of which 2.7% was diarrhea, 1.7% was respiratory distress, and 0.1% of the animals were icteric. None of the animals had neurological disorders. Additionally, a higher proportion of the animals had ectoparasites (93.9%), were thin (76.5%) and have not had any dietary change (98.9%) (**Table 1**).

The sero-prevalence of BCoV in the entire animal population was 53.6%. Upon stratification by sheep, goats, and cattle, the prevalence was 25.8%, 43.1% and 55.8%, respectively. Cattles had significantly higher prevalence of BCoV compared to sheep and goats. Among the entire animal population, seropositivity of BCoV was significantly associated with farms with ≥ 50 animals (75.9% vs 24.1%, $p < 0.0001$). Upon stratification by type of animal, this effect seemed to be explained by cattle (82.2% vs 17.8%, $p = 0.027$) but not sheep and goats that are normally kept in smaller groups (**Table 2**). The sero-prevalence of BCoV was highest in the Northern region followed by the Volta region (**Table 3**). Even though our sampling sites formed a north-south gradient, there was no latitude-dependent trend in seroprevalence.

There was no statistically significant association between the possible risks factors assessed and BCoV sero-positivity among all animals with the exception of dietary change, where a significantly lower odd of BCoV was observed among cattle with recent dietary change [OR=0.08, 95% CI (0.01-0.61), $p = 0.015$] (**Table 4**). Multivariate logistic regression identified both effect to be independent (farm size [OR= 1.39, 95% CI (1.04-1.87), $p = 0.025$]; dietary change [OR=0.07, 95% CI (0.01-0.58), $p = 0.013$]) in cattle (**Table 5**).

Discussion

BCoV is a ubiquitous infection and BCoV-specific antibodies have been detected in cattle populations in numerous countries [20-23]. BCoV shares recent common ancestry with human coronavirus OC43 (HCoV-OC43) [24] and the two are serologically closely related to the extent that HCoV-OC43 is often used as a proxy in serological testing as previously described where specific proteins of BCoV were not available for serological testing or when National legislation restricts the use of certain livestock pathogens [25, 26]. In most African countries such as Ghana, diverse farm animals live in close contact that poses a risk of cross-species infection. Indeed, we found seropositivity against BCoV not only among cattle (55.8%) but also among sheep and goats at 25.8% and 43.1% prevalence rates, respectively. This prevalence was

predominant among animals from the Volta region of Ghana. Cattle presented with significantly higher prevalence of BCoV compared to sheep and goats.

Varying prevalence rates of BCoV have been reported in different countries. A study by Alkan *et al.* reported prevalence ranging from 4.4-100.0% among cattle in Turkey [27]. Another study by Gumusova *et al.* in northern Turkey reported a sero-prevalence of 98.43% in cattle. Yavru *et al.* [23] and O'Connor *et al.* [28] reported a sero-prevalence of 94% among 184 calves and their mothers in Burdur, Turkey and 90% among 852 animals from 3 Ontario feedlots, respectively based on enzyme linked immunosorbent assay (ELISA) method.

Bidokhti *et al.* [29], Hasoksuz *et al.* [30], and Yildirim *et al.* [31] also reported a sero-prevalence of 82-86%, 54.5%, and 26.3%, respectively among cattle. The discrepancies in the prevalence rates compared to that of this present study could be attributed, at least in part, to differences in geographical location, different management systems, source population size, method employed for BCoV antibody detection, and the samples size used in the different investigations. In addition, the higher prevalence rate among cattle could be due to the fact that a higher proportion of the animals were cattle as well as the tropism of BCoV to cattle

Few studies have been conducted to evaluate the prevalence of BCoV among small ruminants. The most recent study was conducted by Gumusova *et al.* in 2007 in Turkey. In their study, they evaluated the sero-prevalence of BCoV in goats by employing commercially available competitive ELISA kits and reported a BCoV sero-prevalence of 41.1% [16]. In a previous study, Eisa and Mohamed reported detection of BCoV antigens in goats [14]. Additionally, Tråvén *et al.*, in a study among 218 sheep from 40 flocks in different parts of Sweden, reported that 19% of the sheep were positive for BCoV antibodies [15]. Prior to these recent studies, there had been reports of BCoV infection in small ruminants in Australia [10], New Zealand [11], Chile [12], and Scotland [13]. Our finding in sheep and goats, thus, provides update information of spillover of BCoV from cattle.

Granted the contagious nature of BCoV, it is imperative that factors that influence exposure and the determinants of BCoV infection be identified which would assist in the development of apt control and preventive measures against BCoV and other infectious diseases.

Though there was no statistically significant association between the possible risks factors assessed and BCoV sero-positivity, we found BCoV sero-positivity to be significantly associated with farms with higher cattle density. This finding is in harmony with studies by Beaudeau *et al.* [32], Hägglund *et al.* [33], and Ohlson *et al.* [34] who reported that large herd size is a risk factor for BCoV infections in dairy cattle. This may be due to poor biosecurity especially among farms with larger herd size in Ghana, and also due to the close contact between animals in these farms which could potentiate the transmission of BCoV compared to farms with small herd size [32, 34].

Infectious diseases surveillance can be greatly enhanced by research studies, as often there is close collaboration between governmental and academic institutions. Restrictions in the movement of samples

to prevent the spread of notifiable livestock diseases create a demand for domestic laboratory capacities. Through this study we hope to demonstrate the value of capacity building in field-based research.

Acknowledgement

This work was supported by Deutsche Forschungsgemeinschaft under a Grant to Y. A. S. and C. D. (DR 772/12-1).

Journal Pre-proof

References

1. Yang D, Leibowitz JL. The structure and functions of coronavirus genomic 3' and 5' ends. *Virus research*. 2015;206:120-33.
2. Oma VS, Tråvén M, Alenius S, Myrnel M, Stokstad M. Bovine coronavirus in naturally and experimentally exposed calves; viral shedding and the potential for transmission. *Virology journal*. 2016;13(1):100.
3. Pfefferle S, Oppong S, Drexler JF, Gloza-Rausch F, Ipsen A, Seebens A, et al. Distant relatives of severe acute respiratory syndrome coronavirus and close relatives of human coronavirus 229E in bats, Ghana. *Emerging infectious diseases*. 2009;15(9):1377.
4. Clark M. Bovine coronavirus. *British Veterinary Journal*. 1993;149(1):51-70.
5. El-Ghorr A, Snodgrass D, Scott F, Campbell I. A serological comparison of bovine coronavirus strains. *Archives of virology*. 1989;104(3-4):241-8.
6. Lathrop SL, Wittum TE, Brock KV, Loerch SC, Perino LJ, Bingham HR, et al. Association between infection of the respiratory tract attributable to bovine coronavirus and health and growth performance of cattle in feedlots. *American journal of veterinary research*. 2000;61(9):1062-6.
7. Boileau MJ, Kapil S. Bovine coronavirus associated syndromes. *Veterinary Clinics: Food Animal Practice*. 2010;26(1):123-46.
8. Ksiazek TG, Erdman D, Goldsmith CS, Zaki SR, Peret T, Emery S, et al. A novel coronavirus associated with severe acute respiratory syndrome. *New England journal of medicine*. 2003;348(20):1953-66.
9. Park S, Kim G, Choy H, Hong Y, Saif L, Jeong J, et al. Dual enteric and respiratory tropisms of winter dysentery bovine coronavirus in calves. *Archives of virology*. 2007;152(10):1885-900.
10. Pass D, Penhale W, Wilcox G, Batey R. Intestinal coronavirus-like particles in sheep with diarrhoea. *The Veterinary Record*. 1982;111:106-7.
11. Durham P, Stevenson B, Farquharson B. Rotavirus and coronavirus associated diarrhoea in domestic animals. *New Zealand Veterinary Journal*. 1979;27:30-2.
12. Reinhardt G, Zamora J, Tadich N, Polette M, Aguilar M, Riedermann S, et al. Diagnosis of coronavirus in sheep in Valdivia province, X region, Chile. *Archivos de Medicina Veterinaria*. 1995;27:129-32.
13. Snodgrass D, Herring J, Reid H, Scott F, Gray E. Virus infections in cattle and sheep in Scotland 1975-1978. *The Veterinary Record*. 1980;106:193-5.
14. Eisa M, Mohamed A. Role of enteric pathogens in enteritis in lambs, goat kids and children and their zoonotic importance. *Vet Med J Giza*. 2004;52:41-59.
15. Tråvén M, Carlsson U, Lundén A, Larsson B. Serum antibodies to bovine coronavirus in Swedish sheep. *Acta Vet Scand*. 1999;40:69-74.
16. Gumusova O, Yazici Z, Albayrak H, Çakiroglu D. First report of bovine rotavirus and bovine coronavirus seroprevalance in goats in Turkey. *Vet glasnik*. 2007;61:75-9.
17. Hoyer BJ, Munster VJ, Nishiura H, Klaassen M, Fouchier RA. Surveillance of wild birds for avian influenza virus. *Emerg Infect Dis*. 2010;16(12):1827.
18. El-Duah P, Sylverken A, Owusu M, Yeboah R, Lamptey J, Frimpong YO, et al. Potential Intermediate Hosts for Coronavirus Transmission: No Evidence of Clade 2c Coronaviruses in Domestic Livestock from Ghana. *Tropical medicine and infectious disease*. 2019;4(1):34.
19. Reusken C, Ababneh M, Raj V, Meyer B, Eljarah A, Abutarbush S, et al. Middle East respiratory syndrome coronavirus (MERS-CoV) serology in major livestock species in an affected region in Jordan, June to September 2013. 2013.

20. Hasoksuz M, Hoet AE, Loerch SC, Wittum TE, Nielsen PR, Saif LJ. Detection of respiratory and enteric shedding of bovine coronaviruses in cattle in an Ohio feedlot. *Journal of Veterinary Diagnostic Investigation*. 2002;14(4):308-13.
21. Kapil S, Trent AM, Goyal SM. Excretion and persistence of bovine coronavirus in neonatal calves. *Archives of virology*. 1990;115(1-2):127-32.
22. Lathrop SL, Wittum TE, Loerch SC, Perino LJ, Saif LJ. Antibody titers against bovine coronavirus and shedding of the virus via the respiratory tract in feedlot cattle. *American journal of veterinary research*. 2000;61(9):1057-61.
23. Yavru S, Yapici O, Kale M, Sahinduran S, Pehlivanoglu F, Albay MK, et al. Bovine coronavirus (BoCV) infection in calves with diarrhoea and their dams. *Acta Scientiae Veterinariae*. 2016;44:1-7.
24. Vijgen L, Keyaerts E, Lemey P, Maes P, Van Reeth K, Nauwynck H, et al. Evolutionary history of the closely related group 2 coronaviruses: porcine hemagglutinating encephalomyelitis virus, bovine coronavirus, and human coronavirus OC43. *Journal of virology*. 2006;80(14):7270-4.
25. Reusken CB, Haagmans BL, Müller MA, Gutierrez C, Godeke G-J, Meyer B, et al. Middle East respiratory syndrome coronavirus neutralising serum antibodies in dromedary camels: a comparative serological study. *The Lancet infectious diseases*. 2013;13(10):859-66.
26. Meyer B, Müller MA, Corman VM, Reusken CB, Ritz D, Godeke GJ, et al. Antibodies against MERS coronavirus in dromedaries, United Arab Emirates, 2003 and 2013. *Emerg Infect Dis*. 2014;20(4):552-9.
27. Alkan F, Bilge-Dağalp S, Can Şahna K, Özgünlük İ. Sığırlarda coronavirus enfeksiyonunun epidemiyolojisi. *Ankara Üniv Vet Fak Derg*. 2003;50:59-64.
28. O'Connor A, Martin SW, Nagy E, Menzies P, Harland R. The relationship between the occurrence of undifferentiated bovine respiratory disease and titer changes to bovine coronavirus and bovine viral diarrhoea virus in 3 Ontario feedlots. *Canadian Journal of Veterinary Research*. 2001;65(3):137.
29. Bidokhti MR, Tråvén M, Fall N, Emanuelson U, Alenius S. Reduced likelihood of bovine coronavirus and bovine respiratory syncytial virus infection on organic compared to conventional dairy farms. *The Veterinary Journal*. 2009;182(3):436-40.
30. Hasoksuz M, Kayar A, Dodurka T, Ilgaz A. Detection of respiratory and enteric shedding of bovine coronaviruses in cattle in Northwestern Turkey. *Acta Veterinaria Hungarica*. 2005;53(1):137-46.
31. Yildirim Y, Dagalp SB, Tan MT, Kalaycioglu AT. Seroprevalence of the rotavirus and corona virus infections in cattle. *Journal of Animal and Veterinary Advances*. 2008;7(10):1320-3.
32. Beaudeau F, Björkman C, Alenius S, Frössling J. Spatial patterns of bovine corona virus and bovine respiratory syncytial virus in the Swedish beef cattle population. *Acta Veterinaria Scandinavica*. 2010;52(1):33.
33. Häggglund S, Svensson C, Emanuelson U, Valarcher J, Alenius S. Dynamics of virus infections involved in the bovine respiratory disease complex in Swedish dairy herds. *The Veterinary Journal*. 2006;172(2):320-8.
34. Ohlson A, Heuer C, Lockhart C, Tråvén M, Emanuelson U, Alenius S. Risk factors for seropositivity to bovine coronavirus and bovine respiratory syncytial virus in dairy herds. *Veterinary Record*. 2010;167(6):201-7.

Table 1. Baseline characteristics of the entire animal population

Variables	Total (n=1495)	Sheep (n=66)	Goat (n=102)	Cattle (n=1327)
Type of animals				
Sheep	66 (4.4)	-	-	-
Goat	102 (6.8)	-	-	-
Cattle	1327 (88.8)	-	-	-
Age of animals				
Weaner	375 (25.1)	7(10.6)	15(14.7)	353(26.6)
Adult	1120 (74.9)	59(89.4)	87(85.3)	974(73.4)
Sex of animals				
Male	430 (28.8)	4(6.1)	13(12.7)	413(31.1)
Female	1065 (71.2)	62(93.9)	89(87.3)	914(68.9)
<i>Lactating females</i>	336 (22.5)	1(1.5)	8(7.8)	327(24.6)
Rectal temperature				
Normal	204 (13.6)	16(24.2)	32(31.4)	204(13.6)
High	1291 (86.4)	50(75.8)	70(68.6)	1291(86.4)
Signs of disease				
No	1430 (95.6)	61(92.4)	92(90.2)	1276(96.2)
Yes	66 (4.4)	5(7.6)	10(9.8)	51(3.8)
Diarrhea	41 (2.7)	0(0.0)	7(6.9)	34(2.7)
Respiratory distress	25 (1.7)	4(6.1)	3(2.9)	18(1.4)
Icterus	1 (0.1)	1(1.5)	0(0.0)	0(0.0)
Neurological disorder	0 (0.0)	0(0.0)	0(0.0)	0(0.0)
Presence of ectoparasites				
No	91 (6.1)	10(15.2)	10(9.8)	71(5.4)

Yes	1404 (93.9)	56(84.8)	92(90.2)	1256(94.6)
Body scoring				
Emaciated	15 (1.0)	1(1.5)	0(0.0)	14(1.1)
Thin	1144 (76.5)	65(98.5)	93(91.2)	986(74.3)
Normal	330 (22.1)	0(0.0)	9(8.8)	321(24.2)
Moderately fat	6 (0.4)	0(0.0)	0(0.0)	6(0.5)
Dietary changes				
No	1479 (98.9)	65(98.5)	98(96.1)	1316(98.9)
Yes	16 (1.1)	1(1.5)	4(3.9)	16(1.1)

Normal temperature for cattle: 37.8-39.5 °C; Normal temperature for sheep and goats: 38.5-40.5 °C

For cattle, sheep and goats: Age of weaner: 6 months to 1 year; Age of adult: >1 year

Table 2. Sero-prevalence of BCoV and its association with farm density

Variables	Sero-prevalence	Farm with < 50 animals	Farm with ≥ 50 animals	p-value
Total				<0.0001
Positive	801(53.6)	193(24.1)	608(75.9)	
Negative	694(46.4)	241(34.7)	453(65.3)	
Sheep				NA
Positive	17(25.8)	17(100.0)	0(0.0)	
Negative	49(74.2)	49(100.0)	0(0.0)	
Goat				NA
Positive	44(43.1)	44(100.0)	0(0.0)	
Negative	58(56.9)	58(100.0)	0(0.0)	
Cattle				0.027
Positive	740(55.8)	132(17.8)	608(82.2)	
Negative	587(44.2)	134(22.8)	453(77.2)	

Table 3. The sero-prevalence of BCoV among the entire animal population stratified by regions

Region	Negative	Positive	p-value
Upper East	191(54.7)	158(45.3)	0.002
Greater Accra	142(47.3)	158(52.7)	
Volta	198(43.3)	259(56.7)	
Northern	163(41.9)	226(58.1)	

Table 4. Possible risk factors for BCoV sero-positivity

Variables	Sheep		Goat		Cattle	
	OR(95% CI)	p-value	OR(95% CI)	p-value	OR(95% CI)	p-value
Age of animals						
Weaner	1		1		1	
Adult	2.23(0.25-20.02)	0.474	0.32(0.10-1.02)	0.054	1.18(0.93-1.51)	0.175
Sex of animals						
Male	1		1		1	
Female	0.32(0.04-2.47)	0.274	1.25(0.38-4.12)	0.716	1.16(0.92-1.46)	0.219
Lactating females	9.00(0.35-231.83)	0.185	2.35(0.53-10.42)	0.261	1.25(0.97-1.62)	0.080
Rectal temperature						
Normal	1		1		1	
High	0.47(0.14-1.58)	0.223	0.55(0.24-1.29)	0.171	0.91(0.65-1.28)	0.606
Signs of disease						
No	1		1		1	
Yes	0.70(0.07-6.77)	0.761	0.30(0.06-1.48)	0.138	1.14(0.65-2.01)	0.654
Diarrhea	-	NA	0.20(0.02-1.74)	0.145	1.14(0.57-2.27)	0.716
Respiratory distress	0.96(0.09-9.89)	0.972	0.65(0.06-7.42)	0.730	0.99(0.39-2.53)	0.986
Icterus	0.92(0.04-23.75)	0.962	-	NA	-	NA
Neurological disorder	-	NA	-	NA	-	NA
Presence of ectoparasites						
No	1		1		1	
Yes	0.27(0.07-1.10)	0.068	1.15(0.31-4.37)	0.833	0.86(0.53-1.40)	0.555
Body scoring						
Normal	-	NA	1		1	
Emaciated	1		-		1.06(0.36-3.12)	0.919
Thin	0.11(0.004-2.86)	0.185	0.94(0.24-3.74)	0.934	0.99(0.77-1.28)	0.098
Moderately fat	-	NA	-		1.59(0.29-8.79)	0.597

Dietary changes						
No	1		1		1	
Yes	9.00(0.35-231.83)	0.185	0.14(0.01-2.60)	0.185	0.08(0.01-0.61)	0.015

Table 5. Multivariate analysis of potential risk factors of BCoV sero-positivity in cattle

Variables	aOR (95% CI)	p-value
Farm density		
<50 animals	1	
≥ 50 animals	1.39(1.04-1.87)	0.025
Body scoring		
Normal	1	
Emaciated	1.01(0.34-3.00)	0.983
Thin	0.91(0.70-1.20)	0.509
Moderately fat	1.78(0.32-9.97)	0.513
Dietary changes		
No	1	
Yes	0.07(0.01-0.58)	0.013
Lactating females	1.27(0.98-1.65)	0.067

Journal Pre-proof